

Hypothesis Testing

Mark Lunt

Arthritis Research UK Epidemiology Unit
University of Manchester



18/10/2011

Introduction

- We saw last week that we can never know the population parameters without measuring the entire population.
- We can, however, make inferences about the population parameters from random samples.
- Last week, we saw how we can create a confidence interval, within which we are reasonably certain the population parameter lies.
- This week, we will see a different type of inference: can we be certain that the parameter **does not** take a particular value ?

Hypothesis Testing

- Start with an interesting hypothesis e.g. Drug A lowers blood pressure more than Drug B.
- Form the opposite (**null**) hypothesis e.g. Drug A and Drug B have the same effect on blood pressure.
- Assess the weight of evidence **against** the **null** hypothesis.
- If there is sufficient evidence against the null hypothesis, it can be rejected.

The Null Hypothesis

- Simplest acceptable model.
- If the null hypothesis is true, the world is uninteresting.
- Must be possible to express numerically (“test statistic”).
- Sampling distribution of test statistic must be known.

Formulating a hypothesis test

Interpreting a hypothesis test

Common types of hypothesis test

Power calculations

Hypothesis tests and confidence intervals

Components of Hypothesis test

Test statistics

Examples

The Alternative Hypothesis

- “Null Hypothesis is untrue”
- Covers any other possibility.
- May be one-sided, if effect in opposite direction is as uninteresting as the null hypothesis

One and Two-sided tests

- Good example: χ^2 test.
 - χ^2 test measures difference between expected and observed frequencies
 - Only unusually large differences are evidence against null hypothesis.
- Bad example: clinical trial
 - A drug company may only be interested in how much better its drug is than the competition.
 - Easier to get a significant difference with a one-sided test.
 - The rest of the world is interested in differences in either direction, want to see a two-sided test.
- One-sided tests are rarely justified

Test Statistic

- Null hypothesis distribution must be known.
 - Expected value if null hypothesis is true.
 - Variation due to sampling error (standard error) if null hypothesis is true.
- From this distribution, probability of any given value can be calculated.
- Can be a mean, proportion, correlation coefficient, regression coefficient etc.

Normally Distributed Statistics

- Many test statistics can be considered normally distributed, if sample is large enough.
- If the test statistic T has mean μ and standard error σ , then $\frac{T - \mu}{\sigma}$ has a normal distribution with mean 0 and standard error 1.
- We do not know σ , we only have estimate s .
- If our sample is of size n , $\frac{T - \mu}{s}$ has a t-distribution with $n - 1$ d.f.
- Hence the term “t-test”.
- If $n \geq 100$, a normal distribution is indistinguishable from the t-distribution.
- Extreme values less unlikely with a t -distribution than a normal distribution.

Formulating a hypothesis test

Interpreting a hypothesis test

Common types of hypothesis test

Power calculations


Hypothesis tests and confidence intervals

Components of Hypothesis test

Test statistics

Examples

Test statistic: Normal distribution



Formulating a hypothesis test

Interpreting a hypothesis test

Common types of hypothesis test

Power calculations


Hypothesis tests and confidence intervals

Components of Hypothesis test

Test statistics

Examples

T-distribution and Normal Distribution



Non-Normally Distributed Statistics

- Statistics may follow a distribution other than the normal distribution.
 - χ^2
 - Mann-Whitney U
- Many will be normally distributed in large enough samples
- Tables can be used for small samples.
- Can be compared to quantiles of their own distribution

Formulating a hypothesis test

Interpreting a hypothesis test

Common types of hypothesis test

Power calculations


Hypothesis tests and confidence intervals

Components of Hypothesis test

Test statistics

Examples

Test Statistic: χ^2_{10}



Example 1: Height and Gender

Null hypothesis On average, men and women are the same height

Alternative Hypothesis One gender tends to be taller than the other.

Test Statistic Difference in mean height between men and women.

One-Sided Hypotheses

- Men are taller than women
- Women are taller than men

Example 2: Drinks preferences

Null hypothesis Equal numbers of people prefer Coke and Pepsi

Alternative Hypothesis Most people prefer one drink to the other

Test Statistic Several possibilities:

- Difference in proportions preferring each drink
- Ratio of proportions preferring each drink

One-Sided Hypotheses

- More people prefer Coke
- More people prefer Pepsi

The p -value

- Probability of obtaining a value of the test statistic at least as extreme as that observed, *if the null hypothesis is true*.
- Small value \Rightarrow data obtained was unlikely to have occurred under null hypothesis
- Therefore, null hypothesis is probably not true.
- It is **not** the probability that the null hypothesis is true.

Interpreting the p -value

- $0 \leq p \leq 1$
- Large p (≥ 0.2 , say) \Rightarrow no evidence against null hypothesis
- $p \leq 0.05 \Rightarrow$ there is some evidence against null hypothesis
- Effect is “statistically significant at the 5% level”
- 0.05 is an arbitrary value: 0.045 is very little different from 0.055.
- Smaller $p \Rightarrow$ more evidence

Factors Influencing *p*-value

- **Effect Size:** a big difference is easier to find than a small difference.
- **Sample Size:** The more subjects, the easier to find a difference
- Always report actual *p*-values, not $p < 0.05$ or $p > 0.05$
- *NS* is unforgivable
- “No significant difference” can mean “no difference in population” or “Sample size was too small to be certain”

Getting it Wrong

- There are two ways to get it wrong:
 - The null hypothesis is true, we conclude that it isn't (Type I error).
 - The null hypothesis is not true, we conclude that it is (Type II error).

Type I Error (α)

- Null hypothesis is true, but there is evidence against it.
- 1 time in 20 that the null hypothesis is true, a statistically significant result at the 5% level will be obtained.
- The smaller the p -value, the less likely we are making a type I error.
- Testing several hypotheses at once increases the probability that at least one of them will be incorrectly found to be statistically significant.
- Several corrections are available for “Multiple Testing”, *Bonferroni's* is the most commonly used, easiest and least accurate.

Type II Error (β)

- Null hypothesis is not true, but no evidence against it in our sample.
- Depends on study size: small studies less likely to detect an effect than large ones
- Depends on effect size: large effects are easier to detect than small ones
- **Power** of a study = $1 - \beta$ = Probability of detecting a given effect, if it exists.

Testing \bar{x}

- Can compare \bar{x} to a hypothetical value (e.g. 0).
- Sometimes called “One-sample t-test”.
- Test statistic $T = \frac{\bar{x} - \mu}{S.E.(x)}$.
- Compare T to a t-distribution on $n - 1$ d.f.

Testing \bar{x} : Example

- The following data are uterine weights (in mg) for a sample of 20 rats. Previous work suggests that the mean uterine weight for the stock from which the sample was drawn was 24mg. Does this sample confirm that suggestion ?
- 9, 14, 15, 15, 16, 18, 18, 19, 19, 20, 21, 22, 22, 24, 24, 26, 27, 29, 30, 32
- $\bar{x} = 21.0$
- S.D.(x) = 5.912

Testing \bar{x} : Solution

$$\begin{aligned} S.E.(x) &= \frac{5.912}{\sqrt{20}} \\ &= 1.322 \end{aligned}$$

$$\begin{aligned} T &= \frac{\bar{x} - 24.0}{S.E.(x)} \\ &= \frac{21.0 - 24.0}{1.322} \\ &= -2.27 \end{aligned}$$

- Comparing -2.27 to a t-distribution on 19 degrees of freedom gives a p -value of 0.035
- I.e if the stock had a mean uterine weight of 24mg, and we took repeated random samples, less than 4 times in 100 would a sample have such a low mean weight.

One-Sample t-test in Stata

```
. ttest x = 24
```

One-sample t test

Variable	Obs	Mean	Std. Err.	Std. Dev.	[95% Conf. Interval]	
x	20	21	1.321881	5.91163	18.23327	23.76673

Degrees of freedom: 19

Ho: mean(x) = 24

Ha: mean < 24
t = -2.2695
P < t = 0.0175

Ha: mean != 24
t = -2.2695
P > |t| = 0.0351

Ha: mean > 24
t = -2.2695
P > t = 0.9825

The Unpaired (two-sample) T-Test

- For comparing two means
- If we are comparing x in a group of size n_x and y in a group of size n_y ,
 - Null hypothesis is $\bar{x} = \bar{y}$
 - Alternative hypothesis is $\bar{x} \neq \bar{y}$
 - Test statistic

$$T = \frac{\bar{y} - \bar{x}}{\text{S.E. of } (\bar{y} - \bar{x})}$$

- T is compared to a t distribution on $n_x + n_y - 2$ degrees of freedom
- You may need to test (`sdtest`) whether the standard deviation is the same in the two groups.
- If not, use the option `unequal`.

Two-Sample t-test in Stata

```
. ttest nurseht, by(sex)
```

Two-sample t test with equal variances

Group	Obs	Mean	Std. Err.	Std. Dev.	[95% Conf. Interval]	
female	227	159.774	.4247034	6.398803	158.9371	160.6109
male	175	172.9571	.5224808	6.911771	171.9259	173.9884
combined	402	165.5129	.4642267	9.307717	164.6003	166.4256
diff		-13.18313	.6666327		-14.49368	-11.87259

Degrees of freedom: 400

Ho: mean(female) - mean(male) = diff = 0

Ha: diff < 0
 t = -19.7757
 P < t = 0.0000

Ha: diff != 0
 t = -19.7757
 P > |t| = 0.0000

Ha: diff > 0
 t = -19.7757
 P > t = 1.0000

Comparing Proportions

- We wish to compare $p_1 = \frac{a}{n_1}$, $p_2 = \frac{b}{n_2}$
- Null hypothesis: $\pi_1 = \pi_2 = \pi$
- Standard error of $p_1 - p_2 = \sqrt{\pi(1 - \pi)\left(\frac{1}{n_1} + \frac{1}{n_2}\right)}$
- Estimate π by $p = \frac{a+b}{n_1+n_2}$
- $\frac{p_1 - p_2}{\sqrt{p(1-p)\left(\frac{1}{n_1} + \frac{1}{n_2}\right)}}$ can be compared to a standard normal distribution

Comparing Proportions in Stata

```
. cs back_p sex
```


	sex		
	Exposed	Unexposed	Total
Cases	637	445	1082
Noncases	1694	1739	3433
Total	2331	2184	4515
Risk	.2732733	.2037546	.2396456
	Point estimate		[95% Conf. Interval]
Risk difference	.0695187		.044767 .0942704
Risk ratio	1.341188		1.206183 1.491304
Attr. frac. ex.	.2543926		.1709386 .329446
Attr. frac. pop	.1497672		

chi2(1) = 29.91 Pr>chi2 = 0.0000			

Sample Size


- Given:
 - Null hypothesis value
 - Alternative hypothesis value
 - Standard error
 - Significance level (generally 5%)
- Calculate:
 - Power to reject null hypothesis for given sample size
 - Sample size to give chosen power to reject null hypothesis

Power Calculations Illustrated: 1




Shaded area = probability of type I error

Power Calculations Illustrated: 2




- $H_0: \bar{x} = 0, S.E.(x) = 1$
- $H_A: \bar{x} = 3, S.E.(x) = 1$
- Power = 85%

Power Calculations Illustrated: 3



- $H_0: \bar{x} = 0, S.E.(x) = 1$
- $H_A: \bar{x} = 4, S.E.(x) = 1$
- Power = 98%

Power Calculations Illustrated: 4



- $H_0: \bar{x} = 0, S.E.(x) = 0.77$
- $H_A: \bar{x} = 3, S.E.(x) = 0.77$
- Power = 97%

Power Calculations in Stata

- `sampsi`
- Only for differences between two groups
- Difference in proportion or mean of normally distributed variable
- Can calculate sample size for given power, or power for given sample size
- Does not account for matching
- Need hypothesised proportion or mean & SD in each group

Parameters in Sample Size Calculation

- Power
- Significance level
- Mean (proportion) in each group
- SD in each group
 - Missing SD \Rightarrow proportion

sampsi syntax for sample size

```
sampsi #1 #2, [ratio() sd1() sd2() power()]  
where
```

- `ratio` Ratio of the size of group 2 to size of group 1 (defaults to 1).
- `sd1` Standard deviation in group 1 (not given for proportions).
- `sd2` Standard deviation in group 2 (not given for proportions). Assumed equal to `sd1` if not given).
- `power` Desired power as a probability (i.e. 80% power = 0.8). Default is 90%.

sampsi for power

```
sampsi #1 #2, [ratio() sd1() sd2() n1() n2() ]  
where
```

- `ratio` Ratio of the size of group 2 to size of group 1 (defaults to 1).
- `sd1` Standard deviation in group 1 (not given for proportions).
- `sd2` Standard deviation in group 2 (not given for proportions). Assumed equal to `sd1` if not given).
- `n1` Size of group 1
- `n2` Size of group 2

sampsi examples

- Sample size needed to detect a difference between 25% prevalence in unexposed and 50% prevalence in exposed:
`sampsi 0.25 0.5`
- Sample size needed to detect a difference in mean of 100 in group one and 120 in group 2 if the standard deviation is 20 and group 2 is twice as big as group 1
`sampsi 100 120, sd1(20) ratio(2)`

Hypothesis Tests and Confidence Intervals

- Hypothesis tests about means and proportions are closely related to the corresponding confidence intervals for the mean and proportion.
 - Null value outside 95% confidence interval \Rightarrow null hypothesis rejected at 5% significance level.
- Confidence intervals convey more information and are to be preferred.
- Dichotomising at $p = 0.05$ ignores lots of information.
- There is a movement to remove “archaic” hypothesis tests from epidemiology literature, to be replaced by confidence intervals.
- If there are several groups being compared, a single hypothesis test is possible, several confidence intervals would be required.

Hypothesis Tests vs. Confidence Intervals

Outcome	Exposed	
	No	Yes
No	6	6
Yes	2	5

- $p = 0.37$
- OR = 2.5, 95% CI = (0.3, 18.3)

Outcome	Exposed	
	No	Yes
No	600	731
Yes	200	269

- $p = 0.36$
- OR = 1.1, 95% CI = (0.9, 1.4)