

Inference in the Presence of Redundant Moment Conditions
and the Impact of Government Health Expenditure on Health
Outcomes in England¹

Martyn Andrews Obbey Elamin Alastair R. Hall²

University of Manchester

Kostas Kyriakoulis Matthew Sutton

North Carolina State University

University of Manchester

Final Version, June 2014

¹This work was performed as part of a project entitled “Towards Improved Inferences in Health Economics Analyses Using Moment-Based Econometric Methods” supported by the National Institute of Health Research (NIHR) Research Methods Funding Scheme.

²Corresponding author: Economics, School of Social Sciences, University of Manchester, Manchester M13 9PL, UK. Email: alastair.hall@manchester.ac.uk

Abstract

In his 1999 paper with Breusch, Qian and Wyhowski in the *Journal of Econometrics*, Peter Schmidt introduced the concept of “redundant” moment conditions. Such conditions arise when estimation is based on moment conditions that are valid and can be divided into two sub-sets: one that identifies the parameters and another that provides no further information. Their framework highlights an important concept in the moment-based estimation literature namely, that not all valid moment conditions need be informative about the parameters of interest. In this paper, we demonstrate the empirical relevance of the concept in the context of the impact of government health expenditure on health outcomes in England. Using a simulation study calibrated to this data, we perform a comparative study of the finite performance of inference procedures based on Generalized Method of Moment (GMM) and info-metric (IM) estimators. The results indicate that the properties of GMM procedures deteriorate as the number of redundant moment conditions increases; in contrast the IM methods provide reliable point estimators but the performance of associated inference techniques based on first order asymptotic theory, such as confidence intervals and overidentifying restriction tests, deteriorates as the number of redundant moment conditions increases. However, for IM methods, it is shown that bootstrap procedures can provide reliable inferences; we illustrate such methods when analysing the impact of government health expenditure on health outcomes in England.

Key words: Generalized Method of Moments, Info-metric estimation, Empirical Likelihood, Exponential Tilting

1 Introduction

The introduction by Lars Hansen (Hansen 1982) of Generalized Method of Moments (GMM) offered a method for obtaining estimators of the parameters of economic models based on the information in population moment conditions. So long as this information is both valid and (strongly) identifies the parameters, Hansen (1982) established the consistency and asymptotic normality of the estimator, and proposed a variant known as the “two-step” GMM estimator which is asymptotically efficient in the class of semi-parametric estimators based on the population moment condition in question, see Chamberlain (1987).

In practice, the underlying economic/statistical model typically implies an array of possible moment conditions, and it has been recognized that the choice of which to use impacts on the comparative statistical properties of the resulting estimator. In essence, moment conditions contain differing amounts of information about the parameters of interest. To pursue this point further, we restrict attention to the class of population moment conditions associated with generalized instrumental variables (IV) estimation that is, in which the moment condition states the orthogonality of vector of instruments to a model residual. This is because this class of moment conditions is the most commonly encountered in econometrics (and is the type involved in our empirical analysis below).

A lot of attention has focused on the two extreme cases, namely optimal instruments and weak instruments. Hansen (1985) characterized the asymptotic efficiency bound for IV, and, since then, various papers have examined how to construct so-called optimal instruments that achieve this bound in certain cases of interest.¹ However, a drawback to their use is that the construction of the optimal instrument can be complicated and may require additional assumptions about the data generation process beyond those implied by the economic model; this often proves a significant limitation and the use of optimal instruments is not common in empirical practice. At the other extreme is the weak instrument case. Following the insight in Nelson and Startz (1990), Staiger and Stock (1997) demonstrated that the standard first order statistical analysis of Hansen breaks down if the instrument is weak;

¹For references see the survey in Newey (1993) or Hall (2005)[Ch. 7.2].

that is, the population moment condition provides insufficient information to (strongly) identify the parameters.² Driven by a number of high profile empirical examples, the problem of inference in the presence of weak instruments has received a lot of attention in the literature.³

However, while both these extremes are of interest, they are not the only information scenarios of relevance in empirical applications. In his 1999 *Journal of Econometrics* paper with Breusch, Qian and Wyhowski, Peter Schmidt introduced the concept of redundant moments—or instruments—which represents an important information scenario that, in some sense, lies in between the two extremes described above. This covers the situation in which a subset of the instruments, z_1 say, lead to moment conditions that strongly identify the parameters and the remainder, z_2 say, provide no additional information. In such circumstances, Hansen’s (1982) analysis still applies and implies that (suitably scaled) the estimator has a limiting normal distribution with mean zero and variance V , say, whether estimation is based on z_1 alone or z_1, z_2 that is, given z_1 is included the addition of z_2 has no impact on the first order asymptotic properties of the estimator; in this case z_2 is said to be *redundant* given z_1 . While this result implies no (first order) asymptotic cost to the inclusion of redundant instruments, we show below, using second order asymptotics, that the finite sample properties of IV are adversely affected by the inclusion of redundant moment conditions.

The concept of redundancy, as originally stated, is occasionally criticised for being unrealistically strict in the sense that z_2 provides *no* additional information beyond that in z_1 . However, this seems pedantic to us: the key insight is to realize that there are situations where some instruments provide identification and most of the information, and the remainder of the instruments provide very little, for which redundancy, as defined above, is just the limit case.⁴

²Also see Stock and Wright (2000) for an analysis of the consequence of weak identification in nonlinear models.

³For a review of the weak instrument literature see Stock, Wright, and Yogo (2002) and Hall (2005)[Ch. 8.2].

⁴For example, the ideas can equivalently be expressed using the concept of *near*-redundancy as in Hall,

In this paper we illustrate these ideas using an important empirical example in which exactly this type of structure is present. For any economy, a key policy question is the extent to which the level of government expenditure on health influences the populations' health. Even though there are surprisingly few estimates of the elasticity of health outcomes with respect to government health expenditure in the literature, estimates of this parameter have been found by regressing (log) mortality on (log) health expenditures, typically exploiting cross-section variations in both variables by region/county/state etc. In this paper, the data are from England (in 2005–06) and the unit of observation is a so-called Primary Care Trust (PCT), of which there are 152. However, expenditures are correlated with the regression error because expenditure is determined by a funding rule that involves four key variables, one of which, a composite need index, is endogenous. As a result, OLS estimation is inappropriate as it leads to inconsistent estimators of the elasticity. However, instrumental variables estimation is feasible because the three other key variables in the funding rule are arguably exogenous and can be used as instruments. Given the construction of the funding rule, these three instruments are expected to be important determinants of expenditure. In addition to these three variables, it is possible to include other instruments, such as variables that are related to the needs index but not to mortality, which tend to be of lesser importance in the determination of expenditure. The “funding rule instruments” are expected to strongly identify the parameters of interest and contain most of the information, whereas the remaining instruments—which we refer to as “secondary”—are likely to add some information but are relatively less important.

The key question for a policy maker is to estimate the elasticity as precisely as possible—especially when using IV estimation—and so directly leads to the issue of which instruments to use in the estimation: if just the funding rule instruments then information from the secondary instruments has been left out at the potential cost of increased asymptotic variance; however, if the secondary instruments are also included but provide so little extra information that they are effectively redundant then their inclusion may adversely affect

Inoue, Jana, and Shin (2008). Also see this paper for a comparison of near-redundancy and weak identification.

finite sample properties. Or put another way, do the secondary instruments provide sufficient additional information that it is beneficial to include them in a sample of size of 150? We explore this issue via a simulation study calibrated to our empirical example, with “stronger instruments” and “less informative instruments” proxying respectively the roles of the funding rule and secondary instruments. We find evidence that the inclusion of the “less informative” instruments does have an adverse effect. This raises the question of how to proceed. One option is to use GMM with just the stronger instruments but our simulation evidence suggests coverage probabilities of confidence intervals based on asymptotic theory may be significantly less than nominal levels in a sample size of 150. Therefore, we explore the use of a member of the class of info-metric (IM) estimators that are argued to have better finite sample properties than GMM.⁵ This class includes both Empirical Likelihood (EL) and Exponential Tilting (ET) estimators. We therefore also explore the performance of these IM estimators in our simulation study. While we find that IM estimators provide more reliable point estimates than GMM, we also find that first order asymptotic theory provides a poor approximation to the coverage probabilities of confidence intervals and rejection frequencies of model specification tests. However, we find these problems can be remedied by employing a bootstrap procedure proposed by Brown and Newey (2002) based on the probabilities obtained as part of the IM estimation. We contrast the inferences based on GMM and IM, and find significant differences in the elasticity of interest. Because the elasticity can be interpreted as the “cost per life”, if our estimates are taken at face value, this has important policy implications.

Section 2 provides the econometric analysis. In Section 3 we describe issues surrounding the estimation of the elasticity of interest using English data, together with some background describing how funding is allocated in England. Section 4 reports the simulation study. In Section 5, we return to our empirical example. Section 6 concludes.

⁵IM estimators can be characterised as Generalized Empirical Likelihood (GEL) estimators (Smith 1997).

2 Moment based inference and redundancy

To formally define redundancy, it is necessary to first present the first order asymptotic distribution of the moment based estimators. The model in our empirical study below fits the following generic linear specification:

$$y_t = x_t' \theta_0 + u_t, \quad t = 1, 2, \dots, T, \quad (1)$$

where y_t is the dependent variable, an observed scalar; x_t is a $(p \times 1)$ vector of observed explanatory (or regressor) variables; u_t is the unobserved error term. The t subscript indicates the observations pertain to the t^{th} member of the sample, and T denotes the sample size. The parameters of interest are denoted by the $p \times 1$ vector θ_0 .

As noted above, IV involves the use of a set of variables as “instruments”; these are denoted by z_t , a $(q \times 1)$ vector of instruments. We assume these instruments are valid in the sense that they are orthogonal to the error so that the following population moment condition holds

$$E[z_t u_t(\theta_0)] = 0, \quad (2)$$

where $u_t(\theta) = y_t - x_t' \theta$. For IV to work, it must be the case (amongst other things) that there are at least as many instruments as parameters, and so we assume $q \geq p$. For ease of presentation, we assume all variables, $v_t = (x_t', u_t, z_t')'$ are independently and identically distributed.

There are a number of ways in which the information in (2) can be exploited to produce estimators of θ_0 . As discussed in the Introduction, we focus on IV estimators (Two Stage Least Squares and the Generalized Method of Moments) and the class of IM estimators (EL and ET). Below we describe both the methods and also their statistical properties, with particular emphasis on the impact of redundant moment conditions on the latter.

2.1 GMM estimation

The GMM estimator based on (2) is defined to be:

$$\hat{\theta}_T = \operatorname{argmin}_{\theta \in \Theta} Q_T(\theta),$$

where

$$Q_T(\theta) = g_T(\theta)' W_T g_T(\theta),$$

$g_T(\theta) = T^{-1} \sum_{t=1}^T z_t(y_t - x_t'\theta)$, and W_T is known as the “weighting matrix”.

For the method to work, the weighting matrix needs to satisfy certain restrictions.⁶ As it is the most common choice in practice, we focus on the two-step GMM estimator that is, $W_T \xrightarrow{P} W = \{\operatorname{Var}[z_t u_t]\}^{-1}$ because this is the optimal choice in the sense that yields the minimum asymptotic variance. Notice that if u_t is conditionally homoscedastic given z_t , then the optimal choice can be obtained using $W_T = (T^{-1} \sum_{t=1}^T z_t z_t')^{-1}$ in which case GMM equals 2SLS. However, if u_t is conditionally heteroscedastic given z_t , then 2SLS is inefficient in large samples relative to the two-step GMM estimator, providing the motivation for using two-step GMM rather than just 2SLS to implement IV.

First order asymptotic analysis

To develop our analysis, we impose a number of conditions. The first of these is to partition the instrument vector into two parts: $z_t = [z'_{1,t}, z'_{2,t}]'$. We then impose the following assumption.

Assumption 1 (i) $(u_t, x'_t, z'_t)'$ are independently and identically distributed and y_t is generated via (1); (ii) $E[z_t u_t] = 0$, and $\operatorname{rank}\{E[x_t z'_{1,t}]\} = p$; (iii) $\operatorname{Var}[u_t | z_t] = \sigma^2$.

Assumption 1(ii) states that z_t are both valid and relevant instruments. Assumption 1(iii) states that the errors are conditionally homoscedastic (given the instruments) as a consequence of which 2SLS and optimal GMM are the same. Under this assumption (and certain

⁶ W_T must be a positive semi-definite matrix that converges in probability to a positive definite matrix of constants; see, for example, Hall (2005)[Chap.1].

other regularity conditions), it can be shown that⁷

$$T^{1/2}(\hat{\theta}_T - \theta_0) \xrightarrow{d} N(0, V_\theta), \quad (3)$$

where $V_\theta = \sigma^2(M_{xz}M_{zz}^{-1}M_{zx})^{-1}$ and $M_{ab} = E[a_t b_t']$ for any random vectors a_t, b_t .

We now define the concept of redundancy; note that although stated in terms of GMM it is equivalently defined for the IM estimators discussed below because they have the same first order asymptotic distribution as two-step GMM. To this end, let $\hat{\theta}_{1,T}$ be the two-step GMM estimator of θ_0 based on $E[z_{1,t}u_t(\theta_0)] = 0$ and assume $T^{1/2}(\hat{\theta}_{1,T} - \theta_0) \xrightarrow{d} N(0, V_\theta^{(1)})$. (Notice that Assumption 1(ii) implies θ_0 is identified by $E[z_{1,t}u_t(\theta_0)] = 0$.)

Definition 1 $E[z_{2,t}u_t(\theta_0)] = 0$ is said to be redundant for estimation of θ_0 given $E[z_{1,t}u_t(\theta_0)] = 0$ if $V_\theta = V_\theta^{(1)}$.

Or equivalently, the instruments $z_{2,t}$ are redundant given $z_{1,t}$ if the asymptotic variance of the estimator is the same whether either $z_{1,t}$ or z_t are used as instruments. In other words, the inclusion of $z_{2,t}$ does not contribute to the asymptotic precision of the estimation. By construction, if $z_{2,t}$ is *non*-redundant given $z_{1,t}$, then $V_\theta^{(1)} - V_\theta$ is positive semi-definite and the inclusion of $z_{2,t}$ improves the asymptotic precision.

As might be expected, it is possible to relate redundancy to the properties of the relationship between the endogenous regressors and instruments. To demonstrate this, we focus on the case that arises in our empirical examples, namely where there is one parameter of interest. Accordingly we partition $x_t = [w_t, c_t']'$ where w_t is the scalar endogenous regressor and c_t is the vector of controls, and then also partition θ conformably as $\theta = (\alpha, \phi)'$. Thus α is the scalar parameter of interest and ϕ is the $(p-1) \times 1$ vector of parameters on the control variables. Let $\hat{\alpha}_T$ be the GMM estimator of α_0 and V_α be the $(1,1)$ element of V_θ .⁸ We further partition $z_{1,t} = [c_t', h_t']'$, where h_t' are the identifying instruments, ie those not used as controls.

⁷For example, see Hall (2005)[Chap. 2.3].

⁸It follows from (3) that $T^{1/2}(\hat{\alpha}_T - \alpha_0) \xrightarrow{d} N(0, V_\alpha)$.

Proposition 1 *If Assumption 1 (and certain other regularity conditions) hold then: $z_{2,t}$ is redundant for estimation of α_0 iff $R_{w,z}^2 = R_{w,z_1}^2$, where R_{w,z_1}^2 is the population multiple correlation coefficient from the regression of w_t on $z_{1,t}$, $R_{w,z}^2$ is the population multiple correlation coefficient from the regression of w_t on z_t .*

The proof is given in the Appendix. Thus, $z_{2,t}$ is redundant given $z_{1,t}$ if it has no additional linear explanatory power for w_t above that of $z_{1,t}$. Note that if $z_{2,t}$ is non-redundant given $z_{1,t}$ then it must follow that $R_{w,z}^2 > R_{w,z_1}^2$.

We can state the first order asymptotic consequences of augmenting an existing set of instruments z_t , by the addition of one extra instrument denoted $z_{3,t}$. If $z_{3,t}$ is redundant given z_t then the inclusion of $z_{3,t}$ makes no difference, but if $z_{3,t}$ is non-redundant given z_t then the inclusion of $z_{3,t}$ improves precision. Either way, the inclusion of the additional instrument never hurts.

Second order asymptotic analysis

For this part of our analysis, we impose one additional assumption.

Assumption 2 (i) $E[u_t^3|z_t] = 0$; (ii) $E[u_t v_t|z_t] = \sigma_{uv} \neq 0$.

Here, v_t is the implied reduced-form error term. Part (i) of this assumption states that the errors are symmetrically distributed conditional on z_t ; part (ii) states that the covariance of u_t and v_t is non-zero. Using results in Newey and Smith (2004), we can show the following.

Proposition 2 *If Assumptions 1, 2 (and certain other regularity conditions) hold then:*

$$\text{bias}(\hat{\alpha}_T) = \frac{(q-p-1)\sigma_{uv}}{T\sigma_w^2} \left(\frac{1}{R_{w,z}^2 - R_{w,z_1}^2} \right). \quad (4)$$

Proposition 2 reveals that the second order bias depends on the number of instruments (q), the explanatory power (over that of the controls) of the instruments for w_t ($R_{w,z}^2 - R_{w,z_1}^2$), the covariance between u_t and v_t (σ_{uv}) and the sample size (T). This is the bias formula for 2SLS introduced by Nagar (1959).

To consider the implications of this second order asymptotic result for the issue of instrument selection in our examples, we again frame our discussion in terms of considering the consequences of augmenting an existing set of instruments z_t by the addition of one extra instrument denoted $z_{3,t}$. Inspection of the formula for $bias(\hat{\alpha}_T)$ in Proposition 2 it can be seen that introduction of an additional instrument impacts both the numerator (by increasing q) and the denominator (by increasing $R_{w,z}^2$). The outcome is therefore ambiguous except in one special case: if $z_{3,t}$ is redundant given z_t then the denominator of the bias term is unaffected by the introduction of $z_{3,t}$ but the numerator increases, meaning the bias must also increase. However, note that this bias disappears as T increases: thus, *ceteris paribus*, the larger the sample, the less the bias.

2.2 Info-metric estimation

Concerns about the finite sample performance of GMM have led to interest in alternative methods of estimation based on the information in moment conditions. Leading examples of such estimators are Empirical Likelihood (EL) (Qin and Lawless, 1994 or Owen, 2001) and Exponential Tilting (ET) (Kitamura and Stutzer, 1997). While EL and ET can be derived from distinct estimation principles, it has been recognized that they have a common structure that has led to development of generic approaches of which both are special cases. The two such generic approaches are Generalized Empirical Likelihood (GEL) (Smith 1997) and Info-metric methods (Golan 2006). We focus on the second approach.

Within the Info-metric approach, the population moment condition (pmc) is viewed as a constraint on true probability distribution of data. If \mathbf{M} is set of all probability measures

then the subset that satisfies pmc for a given θ is

$$\mathbf{P}(\theta) = \left\{ P \in \mathbf{M} : \int f(v, \theta) dP = 0 \right\},$$

and the set that satisfies the pmc for all possible values of θ is

$$\mathbf{P} = \cup_{\theta \in \Theta} \mathbf{P}(\theta).$$

Estimation is based on the principle of finding the value of θ that makes $\mathbf{P}(\theta)$ as close as possible to true distribution of data.

To operationalize this idea, we work with discrete distributions. Let $p_t = P(v = v_t)$ and $P = [p_1, p_2, \dots, p_T]$. Assuming no ties, the empirical distribution of the data is: $\hat{\mu}_t = T^{-1}$; let $\hat{\mu} = [\hat{\mu}_1, \dots, \hat{\mu}_T]$. The Info-metric (IM) estimator is then defined to be:

$$\hat{\theta}_{IM} = \arg \inf_{\theta} \rho_T(\theta, \hat{\mu})$$

where

$$\begin{aligned} \rho_T(\theta, \hat{\mu}) &= \inf_{\hat{P}} D(P \parallel \hat{\mu}), \\ \hat{\mathbf{P}}(\theta) &= \left\{ \hat{P} : p_t > 0, \sum_{t=1}^T p_t = 1, \sum_{t=1}^T p_t f(v_t, \theta) \right\}, \end{aligned}$$

and $D(\cdot \parallel \cdot)$ is a measure of distance. An interpretation of the estimator can be built up as follows. $\hat{\mathbf{P}}(\theta)$ is the set of all discrete distributions that satisfy the pmc for a given value of θ . $\rho_T(\theta, \hat{\mu})$ represents the shortest distance between any member of $\hat{\mathbf{P}}(\theta)$ and the empirical distribution for a particular value of θ . $\hat{\theta}_{IM}$ is the parameter value that makes this distance as small as possible over θ .

To implement the estimator, it is necessary to specify a distance measure. A popular

choice in this literature is the Cressie and Read (1984) distance measure, defined as

$$D_{CR}^{(\eta)}(\mathbf{p}||\mathbf{q}) = \frac{\eta}{1+\eta} \sum_{t=1}^T p_t \left\{ \left(\frac{p_t}{q_t} \right)^\eta - 1 \right\}$$

which is defined for $-\infty < \eta < \infty$. This distance measure nests EL and ET as special cases: $\lim_{\eta \rightarrow 0} D_{CR}^{(\eta)}(\cdot||\cdot)$ yields the ET optimand; $\lim_{\eta \rightarrow -1} D_{CR}^{(\eta)}(\cdot||\cdot)$ yields the EL optimand.

In terms of statistical properties, EL/ET are consistent and have same limiting distribution - and thus the same first order asymptotic properties - as optimal GMM. However, their second order asymptotic properties are different. Using the same set-up as before and the results in Newey and Smith (2004), we can show the second order bias properties of EL/ET are as follows.

Proposition 3 *If Assumptions 1, 2 (and certain other regularity conditions) hold then:*

$$bias(\hat{\alpha}_{IM}) = \frac{-\sigma_{uw}}{T\sigma_w^2} \left(\frac{1}{R_{w,z}^2 - R_{w,z_1}^2} \right) \quad (5)$$

A comparison of the results in Propositions 2 and 3 reveals that the denominator of the bias terms for GMM and IM estimators are the same but there is a crucial difference in the numerators: for GMM the numerator depends on the number of instruments, for IM it does not. So returning to the analysis of the consequence of including an extra instrument, the inclusion of $z_{3,t}$ never increases the absolute bias. So, for IM estimators, there are no potential negative consequences in terms of first or second order asymptotic properties from the inclusion of an additional instrument. This indicates that IM estimators can be expected to yield more reliable point estimators in moderate-sized samples.

3 The impact of government health expenditure on health outcomes in England

From a policy perspective, a key question is whether, and, if so, to what extent, the allocation of funding to public sector health agencies can impact on population health. Such evidence is required to inform decisions about appropriate levels of overall funding and questions of distribution, such as whether differential allocation of funds can contribute to the reduction of inequalities in population health between areas. The extent to which the level of government expenditure on health influences the population's health is particularly important when assessing the wisdom of the UK Government's decision, in 2000, to increase health expenditure to the EU average by 2006 (Appleby and Boyle 2000). It is also relevant for the measurement of public service productivity (ONS 2006).

As noted above, our empirical example comes from England. The National Health Service in England is financed almost entirely from national taxation. The Department of Health negotiates every year with HM Treasury over how much money the National Health Service can spend. The size of the budget in 2005-06 was £53.9 billion, which averaged at £1,097 per person.

The NHS is organised in geographical areas, with Primary Care Trusts taking responsibility for local administration and purchasing of services. These PCTs receive fixed annual budgets from central government and are required to meet their populations' expenditure needs on hospital and community-based services (including pharmaceuticals) and to improve their local population's health.

In England, a funding rule is used to allocate the overall budget to each PCT (DOH 2005). This funding rule creates shares of the overall budget for each PCT that reflect their population size, age and other measured need factors, and expected input prices. These target shares are used to calculate a "Distance From Target" (DFT) for each PCT, which measures the extent to which their actual share of the national budget last year differs from that indicated by their target share. All PCTs receive a minimum level of funding uplift

and the residual funds are then distributed on the basis of the Distance From Target, with the most under-target PCTs receiving the largest increases in budget.

A PCT's budget can therefore be expressed as:

$$\text{Budget per head} = (\text{National budget per head}) * (\text{Age Index}) * (\text{Additional Needs Index}) * (\text{Input Price Index}) * (\text{DFT Index})$$

in which each of the four index adjustments takes a mean value of one.

The data are sourced from government websites. The health measure is a directly age-standardised mortality rate for the period 2005-2007, expressed as deaths per 100,000 European Standard population.⁹ The funding variable is the 2005/6 allocations from the *Unified exposition book: 2003/04, 2004/05 & 2005/06 PCT revenue resource limits*.¹⁰ The formula adjustments are those for the Hospital and Community Health Services element of the formula, taken from Table 5.12 of the same exposition book. The “Distances From Target” are the closing figures for 2005/6 taken from Table 4.2 of the same exposition book. The population counts used to calculate the allocations per head are based on the 2004 Attribution Data Set scaled to Office for National Statistics population projections.

The sample consists of the 152 PCT's in England in 2005-06. In what follows, we estimate the following equation with these data:

$$\ln(H) = \beta \ln(E) + \text{controls} + u, \tag{6}$$

where $\ln(H)$ denotes the log of the mortality rate, $\ln(E)$ is the log of the allocation of health expenditure per head. The exact specification of our observed control variables is irrelevant at this stage. u denotes everything that is unobserved or not included in the model. The variable $\ln(E)$ is potentially endogenous because it is easy to see why expenditure levels might be a function of historical mortality (reverse causality) and because expenditure levels may reflect unobserved area-specific effects (unobserved heterogeneity). Indeed, both

⁹<https://indicators.ic.nhs.uk/webview/>

¹⁰http://webarchive.nationalarchives.gov.uk/+/www.dh.gov.uk/en/Managingyourorganisation/Financeandplanning/Allocations/DH_4000344

sources of endogeneity mean that OLS is biased upwards.

With panel data, we might be able to deal with the latter, but the former can only be addressed using IV. But can suitable instruments be found? For this example, such variables naturally occur because of the funding rule discussed above, namely: the Age Index (Z_1), the Additional Needs Index (N), the Input Price Index (Z_2), and the DFT Index (Z_3). Although N is endogenous, because it depends on historical mortality levels, we argue that the other three variables are uncorrelated with u and can be used as instruments. In practice, the funding rule is not exact but Z_1 , Z_2 and Z_3 are the main determinants of E ; as a result, we refer to these variables as “funding rule instruments” to reflect their relative importance in the determination of E . We also consider the inclusion of other other instruments, such as variables that are related to N (but not H), which tend to be of lesser importance in the determination of E ; these variables we have already labelled “secondary instruments”, and are similar to those used by Martin, Rice, and Smith (2008).

To give a flavour of the issues that this paper addresses, we estimate the model in Equation (6) using the following variables as controls: income deprivation among older people, education deprivation, and a constant term. Summary statistics for the data variables are presented in Table 3.

When the model is estimated using OLS, β is estimated as 0.090 with a robust standard error of 0.064. If taken at face value, this elasticity would imply increases in health expenditure levels have a positive, but statistically, insignificant effect on mortality. From a policy perspective, this positive elasticity is counter intuitive. When re-estimated by 2SLS, using only the three “funding rule instruments” for E , the estimate changes considerably, being -0.705 with a robust standard error of 0.245. In other words, increases in spending do have the expected negative effect on mortality, and the effect is significant, in spite of the increase in its standard error by a factor of 3.8.

This is an excellent example of where 2SLS works: the instruments can only plausibly work through the funding rule, and the upwards bias in OLS is ameliorated. To illustrate the contribution of some of the secondary instruments, we re-estimate once more, adding

the 7 secondary instruments (see Table 3 for full descriptions). Note that the R -squared from the first-stage regression with only the three funding rule instruments is 0.793, and this rises to 0.833 when the 7 secondary instruments are added.¹¹ Now the 2SLS estimate is -0.587 with a robust standard error of 0.132. The question is, which estimate should the policy maker use? Or put another way, does the change in the estimate reflect increased information due to the inclusion of the secondary instruments or does it reflect finite sample bias induced because the secondary instruments are effectively redundant?

Table 1: Summary statistics of variables in health expenditure example*

Variable	Mean	S.D.
<i>Dependent variable</i>		
Directly standardised mortality rate per 100,000: all causes ($\ln(H)$)	614.5	76.33
<i>Endogenous explanatory variable</i>		
Allocation per head ($\ln(E)$)	1,106	138.5
<i>Controls</i>		
Income deprivation among older people (proportion)	0.176	0.065
Education deprivation (proportion)	0.229	0.094
<i>Funding rule instruments</i>		
Age index (Z_1)	0.994	0.051
Input price index (Z_2)	1.036	0.159
Distance from target (DFT) index (Z_3)	1.005	0.081
<i>Secondary instruments^a</i>		
A: Inflow of persons all ages (rate per 1,000 persons)	0.845	0.140
A: Outflow of persons all ages (rate per 1,000 persons)	0.864	0.183
B: Proportion of people aged 16+ who have never married	0.314	0.077
B: Proportion of people in households that own their home	0.693	0.118
B: Proportion of houses failing ODPM ‘Decent Homes Standard’	0.346	0.053
C: Proportion of people aged 16-74 that have never worked	0.031	0.021
C: Proportion of people aged 16-74 that are long-term unemployed	0.011	0.005

* Notes: All variables are subsequently expressed in natural logarithms in the regressions.

^a We collect the secondary instruments into 3 groups later in the analysis, labelled A, B, and C.

In the next section, we examine both this issue and also compare the performance of

¹¹The instruments are clearly not “weak” in the usual sense of the label. The Cragg–Donald Wald F -statistic is 29.2; the corresponding Stock–Yogo weak identification test critical values are 13.91, 9.08, 6.46 respectively for 5%, 10%, 20% maximal IV relative bias.

GMM and IM estimation in a simulation calibrated to our health data. We then return to the above example in Section 5.

4 Simulation study

4.1 Design

In the simulation, we mimic the key properties of our empirical health example introduced in Section 3. However, for ease of exposition, we dispense with all the controls in the regression model apart from the constant ($c_t \equiv 1$), and so the model being estimated is written

$$y_t = \alpha w_t + \phi_0 + u_t, \quad t = 1, 2, \dots, T. \quad (7)$$

In other words, there are $p = 2$ regressors. Because we need to distinguish the stronger instruments from the less informative instruments, we write the reduced-form explicitly as

$$w_t = h_t' \pi_1 + z_{2,t}' \pi_2 + \pi_0 + v_t, \quad t = 1, 2, \dots, T. \quad (8)$$

h_t is (3×1) vector of stronger instruments, and $z_{2,t}$ is $(k \times 1)$ vector of less informative instruments, with the reduced-form parameter vectors π_1 and π_2 having 3 and k elements respectively. In other words, there are $q = k + 4$ instruments in total, and the model is over-identified by $k + 2$. k is the first important parameter in the simulations, because the number of less informative instruments has an ambiguous effect on the second order bias of $\hat{\alpha}$ identified by Proposition 2.

In the simulations, throughout we fix the following parameters as follows. First, $\alpha = -0.5$, so that the true elasticity of health outcomes with respect to health expenditure is negative; second, both constants ϕ_0 and π_0 are normalised to zero; third, $\pi_1 = 1_3$, so that the effect of the stronger instruments on health expenditures is normalised to unity; and fourth, $\pi_2 = a 1_k$, where 1_k is a $(k \times 1)$ vector of ones and a is a scalar. a is the second important choice parameter, because it captures the relative strength of the less informative

instruments compared with their stronger counterparts.

All $k + 3$ instruments are Normally distributed and are drawn independently of each other, each with a variance σ_z^2 . σ_z^2 is the third choice parameter in the simulation design. The reduced-form error v_t and the regression error u_t are drawn independently of the $k + 3$ instruments, but are jointly Normally distributed with the variance of v_t denoted σ_v^2 , the variance of u_t normalised to unity, and the covariance between v_t and u_t denoted ς . When ς is non-zero, w_t is endogenous. ς is the fourth choice parameter in the simulation design. As already explained, we restrict ς to being positive because OLS is upwards biased. Also, because the correlation between u and v is ς/σ_v , and is less than unity, ς is ultimately restricted to $0 \leq \varsigma < \sigma_v$. Finally, all of $u_t, v_t, h_t, z_{2,t}$ have zero mean, which implies that y_t and w_t also have zero mean (given $\phi_0 = \pi_0 = 0$).

Of the four parameters yet to be fixed, namely a, ς, σ_z^2 and k , we note that the number of less informative instruments k varies hereafter as 0, 4, 7, and 10. Although the other three parameters are allowed to vary, in what follows we report only what happens when $a = 1/\sqrt{10}$, $\sigma_z^2 = 1/4$, and $\sigma_v^2 = 3/2$.¹² Our choices are explained as follows. We choose $\sigma_v^2 = 3/2$ because this is what happens in the data. We then set the covariance to $\varsigma = 1$ to ensure a strong degree of endogeneity of the health expenditure variable w , with the correlation between u and v being $\varsigma/\sigma_v = 0.816$. Noting that

$$R_{w,h}^2 = \frac{3\sigma_z^2}{3\sigma_z^2 + \sigma_v^2},$$

and setting $R_{w,h}^2 = 1/3$ throughout, again because of the real data, this implies that $\sigma_z^2 = 1/4$ throughout. Next, note that

$$\sigma_w^2 = \sigma_z^2(3 + ka^2) + \sigma_v^2 \tag{9}$$

and

$$R_{w,[h,z_2]}^2 = \frac{\sigma_z^2(3 + ka^2)}{\sigma_z^2(3 + ka^2) + \sigma_v^2}. \tag{10}$$

¹²The wider set of results is available on request.

We now choose $ka^2 = 1$ so that the contribution of the less informative instruments moves $R_{w,h}^2$ from $1/3$ to $R_{w,[h,z_2]}^2 = 2/5$ when there are $k = 10$ less informative instruments in the reduced form for w . Hence $a = 1/\sqrt{10}$ throughout. To check that these are sensible choices, when $k = 7$ and $\sigma_w^2 = 2.425$, the bias in the OLS estimator, ς/σ_w^2 , is 0.412. In other words, a true α of -0.5 is estimated, on average, as -0.088 using OLS, which is roughly consistent with the real data described in Section 3.

Table 2 summarises the population values of the key parameters for $k = 0, 4, 7, 10$, together with the second order population biases given in Propositions 2 and 3 above. We do this for $T = 150$ and $T = 300$. The former is the sample size for the real data; the latter is to illustrate how the comparisons between the biases evolve as the sample size increases.

Table 2: Summary of simulation design as number of less informative instruments varies

	Number less informative instruments k			
	0	4	7	10
Variance of endogenous regressor σ_w^2 (Equation 9)	2.250	2.350	2.425	2.500
First stage $R_{w,[h,z_2]}^2$ (Equation 10)	0.333	0.362	0.381	0.400
$T = 150$				
Asy Std Error of IV estimator $\sqrt{V_\alpha/T}$ (Equation 12)	0.0943	0.0886	0.0849	0.0817
Second order bias IV estimators (Equation 4)	0.00889	0.03921	0.05766	0.07233
Second order bias IM estimators (Equation 5)	-0.00889	-0.00784	-0.00721	-0.00667
First order bias OLS ς/σ_w^2	0.444	0.426	0.412	0.400
Average F -statistic when estimating Equation (8)	151	35.0	24.1	19.2
$T = 300$				
Asy Std Error of IV estimator $\sqrt{V_\alpha/T}$ (Equation 12)	0.0667	0.0626	0.0600	0.0577
Second order bias IV estimators (Equation 4)	0.00444	0.01961	0.02883	0.03667
Second order bias IM estimators (Equation 5)	-0.00444	-0.00392	-0.00360	-0.00333
First order bias OLS ς/σ_w^2	0.444	0.425	0.412	0.400
Average F -statistic when estimating Equation (8)	301	69.0	47.2	37.4

* Data generation process given by Equations (7, 8). $p = 2$, $q = k + 4$, $\alpha = -0.5$, $\phi_0 = \pi_0 = 0$, $\pi_1 = 1_3$, $\pi_2 = a1_k$, and $\sigma_v^2 = 3/2$. In these simulations, $a = 1/\sqrt{10}$, $\sigma_z^2 = 1/4$, and $\varsigma = 1$.

The table shows that magnitude of the second order bias for the IV and IM estimators is the same when there are no less informative instruments, because $q - p - 1 = 1$. When $T = 150$, the bias is 0.00889. We now see what happens as more and more less informative instruments are added. The variance of the endogenous regressor σ_w^2 increases, and so the

first stage $R_{w,[h,z_2]}^2$ also increases. As the Propositions assert, the second order bias for the IM estimator falls, to 0.00667 for $k = 10$, whereas that for IV increases to 0.07233. For the latter, this is sizeable, as a true parameter of -0.5 will be estimated as -0.4277 on average; in Equation (4), the effect of q in the numerator is outweighing the increase in the fit in the denominator. When the sample size doubles to $T = 300$, the IV estimator is more precise, by a factor $\sqrt{2}$. Also, the second order biases all halve, whereas the first order bias of OLS remains the same.

We now examine the other properties of the IV estimators (2SLS and GMM) and IM estimators (ET and EL) assuming that the sample size is large whereas, in fact, it is a moderate $T = 150$ or $T = 300$. All estimations are performed using the Matlab[®] Optimization Toolbox. The EL and ET estimations utilize a GEL toolbox written by Kostas Kyriakoulis; this toolbox uses *fmincon* with the so-called *interior point* algorithm. This algorithm obtains EL (or ET) estimators by optimizing the EL objective with respect to the probabilities and θ subject to the constraints that the probabilities are non-negative, sum to one, and satisfy the moment condition. The number of replications is $N = 1000$.

4.2 Results

Table 3 summarises the properties of the estimators; that is, biases, coverage proportions, and rejection frequencies based on first order asymptotics (FOA).

The results for the 2SLS and GMM estimators are roughly the same throughout Table 3 because there is no heteroskedasticity in the simulation design. Given the analytical formulae already discussed, the biases reported in the first row of the table can be compared with Table 2: we see that the biases for all the estimators are close to their theoretical counterparts. (Recall that 2SLS/GMM are biased upwards whereas ET/EL are biased downwards.) The biases reported in the second row use the median rather than mean over $N = 1000$ replications; this is because the true sampling distributions might be non-symmetric. Now the biases for 2SLS/GMM get worse, whereas EL/ET get considerably better (roughly 5 times smaller).

Table 3: Coverage and rejection proportions assuming first order asymptotics, $T = 150$

	2SLS				GMM			
	0	4	7	10	0	4	7	10
Bias ^a	0.0091	0.0374	0.0517	0.0661	0.0091	0.0372	0.0520	0.0666
Bias (median) ^b	0.0175	0.0417	0.0592	0.0694	0.0176	0.0414	0.0587	0.0710
Bias mean corrected est ^c					0.0015	0.0093	0.0174	0.0311
Coverage prop, 95% nom, t -stat ^d	0.939	0.893	0.838	0.774	0.929	0.862	0.808	0.732
Coverage prop, 99% nom, t -stat	0.976	0.950	0.937	0.899	0.972	0.940	0.906	0.850
Rejection prop, 5% nom, J -stat ^e	0.052	0.065	0.066	0.079	0.053	0.046	0.045	0.047
Rejection prop, 1% nom, J -stat	0.009	0.013	0.017	0.019	0.008	0.005	0.007	0.009

	EL				ET			
	0	4	7	10	0	4	7	10
Bias	-0.0096	-0.0086	-0.0098	-0.0092	-0.0094	-0.0090	-0.0100	-0.0093
Bias (median)	-0.0019	-0.0021	-0.0007	-0.0024	-0.0018	-0.0022	0.0001	-0.0030
Bias mean corrected est	-0.0004	-0.0012	-0.0034	-0.0037	-0.0003	-0.0016	-0.0036	-0.0038
Coverage prop, 95% nom, t -stat	0.944	0.928	0.907	0.889	0.942	0.925	0.900	0.882
Coverage prop, 99% nom, t -stat	0.980	0.972	0.968	0.960	0.978	0.970	0.961	0.951
Rejection prop, 5% nom, LR ^f	0.059	0.078	0.120	0.178	0.063	0.113	0.173	0.262
Rejection prop, 5% nom, LM	0.056	0.081	0.137	0.208	0.056	0.062	0.075	0.079
Rejection prop, 5% nom, W	0.056	0.081	0.137	0.208	0.073	0.149	0.232	0.350
Rejection prop, 1% nom, LR	0.013	0.025	0.041	0.064	0.018	0.038	0.072	0.115
Rejection prop, 1% nom, LM	0.012	0.029	0.045	0.087	0.010	0.010	0.019	0.018
Rejection prop, 1% nom, W	0.012	0.029	0.045	0.087	0.023	0.064	0.121	0.215

^a Bias is $N^{-1} \sum_r \hat{\alpha}_r - \alpha$, where $\hat{\alpha}_r$ is the estimate of α on the r -th replication.

^b As [a], but using median rather than sample mean.

^c Estimate of $\hat{\alpha}_r$ is bias-corrected using suggestion of Newey and Smith (2004).

^d Proportion of replications where $H_0 : \alpha = -0.5$ not rejected using t -statistic $\sqrt{T}(\hat{\alpha} + 0.5)/\sqrt{\hat{V}_\alpha}$.

^e Proportion of replications where usual GMM overidentifying restrictions test is rejected.

^f Same as [e], but for LR, LM, and Wald analogues of overidentifying restrictions test for ET/EL.

Table 4: Coverage and rejection proportions assuming first order asymptotics, $T = 300^*$

	2SLS				GMM			
	0	4	7	10	0	4	7	10
Bias ^a	0.0038	0.0182	0.0269	0.0349	0.0040	0.0182	0.0271	0.0353
Bias (median) ^b	0.0086	0.0201	0.0282	0.0359	0.0089	0.0207	0.0280	0.0347
Bias mean corrected est ^c					-0.0005	0.0011	0.0066	0.0114
Coverage prop, 95% nom, t -stat ^d	0.933	0.907	0.879	0.856	0.930	0.891	0.860	0.823
Coverage prop, 99% nom, t -stat	0.980	0.965	0.954	0.935	0.977	0.956	0.939	0.922
Rejection prop, 5% nom, J -stat ^e	0.054	0.053	0.058	0.062	0.047	0.052	0.056	0.056
Rejection prop, 1% nom, J -stat	0.013	0.014	0.010	0.012	0.010	0.013	0.007	0.006

	EL				ET			
	0	4	7	10	0	4	7	10
Bias	-0.0051	-0.0052	-0.0047	-0.0035	-0.0052	-0.0053	-0.0049	-0.0036
Bias (median)	-0.0001	-0.0031	-0.0031	-0.0027	-0.0005	-0.0032	-0.0038	-0.0033
Bias mean corrected est	-0.0005	-0.0013	-0.0013	-0.0004	-0.0006	-0.0015	-0.0014	-0.0005
Coverage prop, 95% nom, t -stat	0.937	0.928	0.917	0.910	0.935	0.926	0.919	0.911
Coverage prop, 99% nom, t -stat	0.980	0.979	0.976	0.971	0.980	0.980	0.972	0.970
Rejection prop, 5% nom, LR ^f	0.056	0.068	0.085	0.102	0.059	0.082	0.103	0.139
Rejection prop, 5% nom, LM	0.055	0.064	0.084	0.104	0.050	0.060	0.063	0.077
Rejection prop, 5% nom, W	0.055	0.064	0.084	0.104	0.062	0.095	0.132	0.175
Rejection prop, 1% nom, LR	0.012	0.016	0.019	0.025	0.014	0.025	0.030	0.052
Rejection prop, 1% nom, LM	0.011	0.016	0.019	0.028	0.011	0.013	0.014	0.017
Rejection prop, 1% nom, W	0.011	0.016	0.019	0.028	0.014	0.031	0.045	0.078

* See tablenotes to Table 3.

For the GMM/EL/ET estimators, we implement a second-order bias-correction using a suggestion of Newey and Smith (2004). For ET/EL, this reduces the bias by a factor of 2 to 3. However, we note the correction seems sensitive to k with the remaining biases are 3 times worse for $k = 7, 10$ compared with $k = 4$. Importantly, our simulations confirm that EL/ET exhibits relatively little bias, even when there are $k = 10$ less informative instruments. Throughout, the biases for ET are very similar to EL. On the other hand, it is clear that the 2SLS/GMM estimators are biased, and the bias gets worse as more less informative instruments are added. However, the bias correction does make the biases smaller, and the “drift” with k is less pronounced. However, the bias corrected GMM estimator always exhibits more bias than bias corrected EL/ET and more bias than the uncorrected EL/ET for $k = 4, 7, 10$.

We now examine whether we obtain correct inference when testing the null hypothesis that $\alpha = -0.5$. The table shows that the coverage proportions are all too small for $k = 0$ (for example, the ET coverage is 0.944 instead of 0.95 and is 0.980 instead of 0.99) and these get worse as k increases. There is no difference between ET and EL. This deterioration is much worse for the 2SLS/GMM estimators.

Finally, we consider the performance of the standard model specification test statistics within the GMM and IM framework. For GMM, this is the overidentifying restrictions test statistic:

$$J = Tg_T(\hat{\theta}_T)'W_Tg_T(\hat{\theta}_T)$$

where $g_T(\theta) = T^{-1} \sum_{t=1}^T z_t u_t(\theta)$ and $W_T = \{T^{-1} \sum_{t=1}^T u_t^2(\hat{\theta}_T) z_t z_t'\}^{-1}$; the 2SLS version uses $W_T = \{\hat{\sigma}^2 T^{-1} \sum_{t=1}^T z_t z_t'\}^{-1}$. For the IM, we consider three model specification tests:

the LR, Wald and LM given respectively by¹³

$$\begin{aligned}
 LR &= 2 \sum_{t=1}^T [\rho(\hat{\lambda}' z_t u_t(\hat{\theta}_{IM})) - \rho_0] \\
 Wald &= T g_T(\hat{\theta}_{IM})' \hat{\Omega}^{-1} g_T(\hat{\theta}_{IM}) \\
 LM &= T \hat{\lambda}' \hat{\Omega} \hat{\lambda},
 \end{aligned}$$

where $\rho(a)$ equals $\ln(1-a)$ for EL and $-e^a$ for ET, $\rho_0 = \rho(0)$, $\hat{\lambda}$ is the Lagrange Multiplier associated with constraint that the moment conditions are satisfied in the sample, $\hat{\Omega} = T^{-1} \sum_{t=1}^T u_t^2(\hat{\theta}_{IM}) z_t z_t'$. Under $H_0 : E[z_t u_t(\theta_0)] = 0$ all four statistics converge to a χ_{q-p}^2 distribution. From Table 3, it can be seen the GMM overidentifying restrictions test, the so-called J -statistic, has the correct rejection proportions, and they are marginally worse for 2SLS. For the EL/ET estimators, when there are no less informative instruments, the rejection frequencies are also correct. However, when k increases, for the ET/EL estimators both the LR and Wald over-reject (for example, for ET and $k = 10$, the null hypothesis is rejected 35.0% of the time when using the Wald statistic when it should be 5%). Of the three IM-based statistics, only the LM test for the ET estimator exhibits a rejection frequency close to the nominal size.

Table 4 reports what happens when we double the sample size. As expected, all the biases halve. Also, for obvious reasons, the rejection proportions for EL/ET improve considerably.

To conclude, while EL and ET provide accurate point estimators, inferences based on the first order asymptotic distribution of the estimator and overidentification restriction tests are unreliable in these samples when less informative instruments are included. In the next sub-section, we explore whether a bootstrap can correct this problem.

¹³See Smith (2011)[Section 4].

4.3 Bootstrap

In this sub-section, we outline Brown and Newey's (2002) bootstrap procedure for constructing confidence intervals for the parameter estimators and performing the various versions of the overidentifying restrictions tests.¹⁴ For purposes of presentation, we let the random vector containing the data variables be $d = [y, w, z']'$, and d_t denote the t^{th} observation in the sample on d , that is $d_t = [y_t, w_t, z_t']'$. The statistics of interest are:

t-statistics:

$$\tau_{(\cdot)}(\hat{\theta}_j, \beta_{0,j}, \hat{V}_{jj}) = \left| \frac{\hat{\theta}_j^{(\cdot)} - \theta_{0,j}}{\sqrt{\hat{V}_{jj}}} \right|$$

for $(\cdot) = GMM, EL, ET$ with $\hat{\theta}_j$ and \hat{V}_{jj} denoting the appropriate estimator and its estimated first order asymptotic variance based on estimation method indicated by (\cdot) .

Overidentifying restrictions tests: $O_s^{(\cdot)}(d_1, d_2, \dots, d_T)$ where $(\cdot) = GMM, EL, ET$, and for $(\cdot) = GMM$ then $s = 1$ denotes the usual GMM overidentifying restrictions test, for $(\cdot) = EL$ or ET then $s = 1$ denotes the Wald statistic for the overidentifying restrictions, $s = 2$ denotes the LR statistic and $s = 3$ denotes the LM statistic.

Brown and Newey (2002) propose a version of the bootstrap based on GEL estimation of this model. To describe their procedure, we introduce the following definition.

- Let $\pi_t = P(d = d_t)$ and $\hat{\pi}_t$ denote the GEL estimator of π based on the sample.

For the purposes of the bootstrap, we treat d as discrete random vector with sample space $\mathcal{D}_T = \{d_t; t = 1, 2, \dots, T\}$ and probability distribution function $P_T(d = d_t) = \hat{\pi}_t$.¹⁵ The bootstrap samples are then created by sampling from replacement from this distribution

¹⁴Brown and Newey (2002) propose this procedure for GMM-based inference only but it can also be used for EL and ET because it is an example of the "b-bootstrap" methods analyzed in Hall and Presnell (1999). It has become customary in the econometrics literature to ascribe this resampling based on EL/ET to Brown and Newey, for example see Kitamura (2007), and so we continue this practice here.

¹⁵We ignore the possibility here that $d_t = d_s$ for some $t \neq s$ as this does not occur in our simulations or the health data.

for d . Let B be the total number of bootstrap samples generated, and index the bootstrap sample by b ; so we have $b = 1, 2, \dots, B$. Then, for each step of the bootstrap b , we proceed as follows.

1. On the b^{th} step of the bootstrap, draw a sample T observations $d_t^{(b)}$ with replacement from $P_T(d = d_t) = \hat{\pi}_t$.
2. Based on $\{d_t^{(b)}; t = 1, 2, \dots, T\}$, calculate:
 - the GEL and GMM estimators, denoted here by $\hat{\theta}_{GMM}^{(b)}$, $\hat{\theta}_{EL}^{(b)}$ and $\hat{\theta}_{ET}^{(b)}$.
 - the test statistics:
 - $\tau_{(\cdot)}^{(b)}(\hat{\theta}_j^{(b)}, \hat{\theta}_j, \hat{V}_{jj}^{(b)})$ where $\hat{\theta}_j^{(b)}$ and $\hat{V}_{jj}^{(b)}$ denote the appropriate estimator and its estimated first order asymptotic variance based on estimation method indicated by (\cdot)
 - $O_s^{(\cdot),b}(d_1^{(b)}, d_2^{(b)}, \dots, d_T^{(b)})$ where $(\cdot) = GMM, EL, ET$ and s is defined as above.

As a result of applying the bootstrap, this procedure generates a sampling distribution for each statistic of interest. It uses these distributions to provide bootstrap-based confidence intervals for the parameters and bootstrap-based p -values for the overidentifying restrictions tests as follows:

- *bootstrap-based confidence interval for $\beta_{0,j}$* : Let $\tau_b = \tau_{(\cdot)}^{(b)}(\hat{\theta}_j^{(b)}, \hat{\theta}_j, \hat{V}_{jj}^{(b)})$, that is the value of $\tau_{(\cdot)}^{(b)}(\hat{\theta}_j^{(b)}, \hat{\theta}_j, \hat{V}_{jj}^{(b)})$ based on the b^{th} bootstrap sample¹⁶. From the bootstrap, the procedure generates the following sampling distribution for $\tau_{(\cdot)}(\hat{\theta}_j, \beta_{0,j}, \hat{V}_{jj})$, $\{\tau_b\}_{b=1}^B$. Let q_α^B be the $100(1 - \alpha)^{th}$ quantile of $\{\tau_b\}_{b=1}^B$: the $100(1 - \alpha)\%$ bootstrap-based symmetric confidence interval for $\beta_{0,j}$ is:

$$\hat{\theta}_j \pm q_\alpha^B \sqrt{\hat{V}_{jj}}. \quad (11)$$

¹⁶Note this depends on j but this suppressed to simplify the notation

- *bootstrap-based p-values for the overidentifying restrictions tests:* To illustrate, consider the overidentifying restrictions test based on GMM, denoted $O_1^{GMM}(d_1, d_2, \dots, d_T)$ above. Put $O_b = O_1^{GMM}(d_1^{(b)}, d_2^{(b)}, \dots, d_T^{(b)})$, that is the value of the GMM overidentifying restrictions test based on the b^{th} bootstrap sample - again, for simplicity of notation, we suppress dependence (this time) on (\cdot) and s . From the bootstrap, the procedure generates the following sampling distribution for $O_1^{GMM}(d_1, d_2, \dots, d_T)$, $\{O_b\}_{b=1}^B$. Redefine q_α^B to be the $100(1 - \alpha)^{th}$ quantile of $\{O_b\}_{b=1}^B$: then the bootstrap version of the decision rule for this test is as follows: reject $H_0 : E[z_t u_t(\theta_0)] = 0$ if $O_1^{GMM}(d_1, d_2, \dots, d_T) \geq q_\alpha^B$, that is test statistic from original data is compared to the appropriate quantile obtained from the bootstrapped sampling distribution.

Table 5: Coverage and rejection proportions using the bootstrap, $T = 150$

	EL				ET			
	0	4	7	10	0	4	7	10
Coverage prop, 95% nom, t -stat	0.951	0.954	0.945	0.942	0.951	0.956	0.939	0.938
Coverage prop, 99% nom, t -stat	0.984	0.988	0.985	0.983	0.987	0.986	0.985	0.983
Rejection prop, 5% nom, LR ^f	0.049	0.042	0.045	0.039	0.052	0.047	0.047	0.045
Rejection prop, 5% nom, LM	0.052	0.052	0.045	0.043	0.049	0.034	0.029	0.019
Rejection prop, 5% nom, W	0.052	0.052	0.045	0.043	0.055	0.053	0.051	0.047
Rejection prop, 1% nom, LR	0.008	0.005	0.007	0.006	0.010	0.009	0.007	0.007
Rejection prop, 1% nom, LM	0.008	0.009	0.009	0.006	0.005	0.005	0.003	0.004
Rejection prop, 1% nom, W	0.008	0.009	0.009	0.006	0.011	0.011	0.011	0.006

* See tablenotes to Table 3.

Tables 5 and 6 report the outcomes, and shows that all the coverage and rejection probabilities are consistent with the nominal size.

Table 6: Coverage and rejection proportions using the bootstrap, $T = 300$

	EL				ET			
	0	4	7	10	0	4	7	10
Coverage prop, 95% nom, t -stat	0.942	0.943	0.928	0.937	0.942	0.938	0.934	0.932
Coverage prop, 99% nom, t -stat	0.985	0.987	0.985	0.985	0.985	0.988	0.987	0.985
Rejection prop, 5% nom, LR ^f	0.050	0.048	0.054	0.053	0.052	0.049	0.056	0.056
Rejection prop, 5% nom, LM	0.055	0.051	0.051	0.055	0.043	0.042	0.049	0.041
Rejection prop, 5% nom, W	0.055	0.051	0.051	0.055	0.051	0.047	0.056	0.058
Rejection prop, 1% nom, LR	0.009	0.011	0.010	0.006	0.009	0.012	0.010	0.007
Rejection prop, 1% nom, LM	0.011	0.010	0.010	0.006	0.010	0.007	0.007	0.003
Rejection prop, 1% nom, W	0.011	0.010	0.010	0.006	0.009	0.013	0.010	0.007

* See tablenotes to Table 3.

5 The health expenditure example continued

We continue with the empirical example we introduced in Section 3 above. Table 7 reports results from 2SLS and GMM estimation of the model using various choices of instrument.¹⁷ To recap Section 3, in contrast to the OLS estimate of 0.090 (0.064), the 2SLS estimate with the funding rule instruments is -0.705 (0.245) (see the column labelled “Base” and row labelled “2SLS”). We now report what happens when we add up to seven secondary instruments, and re-estimate the models using GMM, EL and ET.

In the rest of the row labelled “2SLS”, the secondary instruments are added in groups, so that the final column has 7 such instruments. Now the estimate is -0.587 (0.132). Recall that the first stage R -squared is 0.793 for the Base model and rises to 0.833 when the 7 extra instruments are included. In the second row, all the models are re-estimated using GMM. All the estimates are smaller in absolute value, and have larger standard errors, except for the Base model.

The issue we have addressed in this paper is the fact that the 2SLS/GMM estimates are sensitive to the number of secondary instruments. In particular, we note that the estimated elasticity tends to become smaller in absolute value as more instruments are included. By

¹⁷We have applied standard tests available in the literature to confirm the relevance and validity of all choices of instruments considered here. Details are omitted for brevity.

Table 7: 2SLS, GMM, EL and ET estimates of β and standard errors*

Estimator	Base	A	B	C	AB	AC	BC	ABC
2SLS	-0.705 (0.245)	-0.656 (0.148)	-0.705 (0.158)	-0.642 (0.148)	-0.608 (0.135)	-0.625 (0.142)	-0.641 (0.145)	-0.587 (0.132)
GMM	-0.627 (0.200)	-0.593 (0.192)	-0.574 (0.175)	-0.518 (0.172)	-0.527 (0.162)	-0.536 (0.171)	-0.507 (0.160)	-0.511 (0.157)
EL	-0.743 (0.218)	-0.787 (0.233)	-0.683 (0.192)	-0.658 (0.191)	-0.753 (0.204)	-0.694 (0.200)	-0.642 (0.179)	-0.734 (0.196)
ET	-0.747 (0.218)	-0.786 (0.231)	-0.681 (0.190)	-0.665 (0.192)	-0.729 (0.197)	-0.709 (0.201)	-0.642 (0.178)	-0.710 (0.189)
<i>Bias corrected</i>								
GMM	-0.659	-0.684	-0.646	-0.589	-0.638	-0.652	-0.608	-0.654
EL	-0.711	-0.756	-0.655	-0.633	-0.725	-0.670	-0.619	-0.708
ET ^a	-0.731	-0.770	-0.663	-0.653	-0.716	-0.698	-0.629	-0.698
<i>Bootstrap-based p-values for the overidentifying restrictions tests^b</i>								
EL, LR	0.81	0.74	0.47	0.75	0.69	0.65	0.41	0.57
EL, LM	0.80	0.73	0.45	0.74	0.68	0.63	0.39	0.56
EL, Wald	0.80	0.73	0.46	0.74	0.68	0.64	0.39	0.56
ET, LR	0.80	0.71	0.47	0.74	0.65	0.63	0.41	0.53
ET, LM	0.83	0.77	0.55	0.77	0.72	0.68	0.49	0.61
ET, Wald	0.79	0.70	0.44	0.72	0.64	0.62	0.39	0.52
<i>Bootstrapped-based confidence interval^c</i>								
EL (upper)	-0.216	-0.184	-0.186	-0.210	-0.239	-0.178	-0.188	-0.165
EL (lower)	-1.270	-1.390	-1.180	-1.105	-1.268	-2.211	-1.096	-1.303
ET (upper)	-0.160	-0.186	-0.088	-0.205	-0.193	-0.163	-0.119	-0.120
ET (lower)	-1.334	-1.387	-1.275	-1.125	-1.265	-1.254	-1.165	-1.299

* Notes: for definitions see Table 3. “Base” specification is 3 funding rule instruments only. “A” adds 2 migration variables; “B” adds 3 further socioeconomic variables; “C” adds 2 further labour market variables; so that ... “ABC” adds all 7 variables.

^a ET based on $B \approx 950$ because ET did not always converge.

^b See Section 4.3 for full details. (The numbers of bootstrapped samples that were discarded are 41, 30, 51, 21, 38, 29, 43, and 54 resp.)

^c See Equation (11).

contrast, the EL/ET estimates exhibit far less sensitivity to the choice of instrument than their 2SLS/GMM counterparts. In particular, it is interesting to compare the estimates of the Base specification (funding-rule instruments only) with the “ABC” specification (all the instruments). For GMM, the estimates for Base are -0.627 and with “ABC” are -0.511 ; where as for EL they are -0.743 and -0.734 respectively, and for ET, -0.747 and -0.710 respectively. As is apparent, the EL and ET estimates are close and different from those obtained via 2SLS/GMM.

Given the insights from first and second order asymptotic theory described above, we believe that the EL/ET estimates are the more reliable. Our simulations also suggest that the Newey and Smith (2004) second order bias correction reduces the bias for all values of k ; see the second panel of the table. In Table 7, the implied biases match the patterns predicted by the simulations. The biased-corrected EL estimates are approximately 0.03 closer to zero than the uncorrected estimate and for ET this is about 0.015, ie the estimates move in the right direction. There is no systematic variation across the columns. On the other hand, again as predicted, the GMM estimates move in the other direction, move further than EL/ET, are get bigger as more of the secondary instruments are added. The net effect is that the bias-corrected GMM estimates do not vary with the number of secondary instruments, as one would hope.

However, our simulations also show that the “usual” inference techniques based on first order asymptotic theory are unreliable and this problem can be corrected using the bootstrap. Therefore, we apply Brown and Newey’s (2002) procedure, described in Section 4.3, to our example. In the third panel, the p -values for the overidentifying restrictions tests all pass comfortably, and, in the fourth panel, we report the corresponding bootstrapped-based confidence intervals.

Of the second order biased corrected EL/ET estimates, which should we choose? Whilst the EL estimates range between -0.619 and -0.756 and the ET estimates range between -0.629 and -0.770 , because this paper is concerned with the relevance of secondary instruments, we focus on the ABC specifications, namely -0.708 for EL and -0.698 for ET.

These estimates are roughly in the middle of the 8 possibilities.

Thus we conclude that elasticity of mortality with respect to health expenditure is roughly -0.71. The equivalent GMM estimate is roughly -0.51, which we believe is biased because of the inclusion of the secondary instruments. In terms of policy implications, the differences between the GMM and EL estimators can be demonstrated as follows: the GMM estimates imply that increasing NHS expenditure by 10% leads to a 5.1% reduction in deaths which translates to a cost per death averted of approximately £350,000, but the EL estimates imply that a 10% increase in NHS expenditure would lead to 7.3% fewer deaths which translates to a cost per death averted of £250,000.¹⁸ However, the bootstrap confidence intervals suggest that there is considerable uncertainty about the estimate, with, for example, the EL estimate having a confidence interval of (-0.152, -1.264). The corresponding cost per death calculations turn out to be £1,120,000 and £140,000 respectively.

6 Concluding remarks

In his 1999 paper with Breusch, Qian and Wyhowski in the *Journal of Econometrics*, Peter Schmidt introduced the concept of “redundant” moment conditions. Such conditions arise when estimation is based on moment conditions that is valid and can be divided into two sub-sets: one that identifies the parameters and another that provides no further information. Their framework highlights an important concept in the moment-based estimation literature namely, that not all valid moment conditions need be informative about the parameters of interest.

In this paper, we demonstrate the empirical relevance of the concept in the context of the impact of government health expenditure on health outcomes in England because this is where exactly this type of structure is present. From the funding rule, there are

¹⁸These figures are obtained as follows. Using the summary statistics reported in Table 3 above, an increase in NHS expenditure by 10% would cost £107 per capita. The death probability is 0.006 and a 5.1% reduction in the death probability (from GMM) is a change of 0.000306. The cost per death averted is then $\text{£}107/(0.000306) = \text{£}350,000$ approximately. With the coefficient estimated from the EL (-0.71) the change in death probability is 0.000426, giving a cost per death averted of £250,000 approximately. In general, the cost per death averted is $\text{£}180,000/\beta$.

a set of 3 instruments that strongly identify the parameters and an additional group of seven secondary instruments capturing socio-economic characteristics that are potentially redundant, or nearly so, given the funding rule instruments. In estimating the elasticity of mortality with respect to health expenditure using English data for 152 PCTs in 2005–06, the 2SLS estimate falls from -0.705 (0.245) with only the funding rule instruments are used to -0.587 (0.132) when the additional seven instruments are also included. This raises the obvious question of which figure the policy maker should use.

Using second order asymptotic analysis backed up by a simulation study calibrated to these data, we perform a comparative study of the finite performance of inference procedures based on Generalized Method of Moment (GMM) and info-metric (IM) estimators. The results indicate that the properties of GMM procedures deteriorate as the number of less informative moment conditions increases; in contrast the IM methods provide reliable point estimators but the performance of associated inference techniques based on first order asymptotic theory, such as confidence intervals and overidentifying restriction tests, deteriorates as the number of less informative moment conditions increases. These results suggest that IM estimates combined with the Brown and Newey (2002) bootstrap provide reliable inferences.

When we return to the health example, we find that the IM point estimate implies a substantially lower cost per life saved than the GMM estimator. However, the bootstrapped confidence intervals suggest that there is considerable uncertainty about the estimate.

Appendix

Proof of Proposition 1: Using the arguments in Section 2, we have

$$\begin{aligned} V_\theta &= \sigma^2(M_{xz}M_{zz}^{-1}M_{zx})^{-1} = \sigma^2 A^{-1}, \text{ say,} \\ V_\theta^{(1)} &= \sigma^2(M_{xz_1}M_{z_1z_1}^{-1}M_{z_1x})^{-1} = \sigma^2\{A^{(1)}\}^{-1}, \text{ say.} \end{aligned}$$

Consider A . Using $x = [w, c']'$, we have (dropping the t for simplicity)

$$A = \begin{bmatrix} E[ww'] \\ E[cc'] \end{bmatrix} \{E[zz']\}^{-1} \begin{bmatrix} E[zw] & E[zc'] \end{bmatrix} = \begin{bmatrix} M_{wz}M_{zz}^{-1}M_{zw} & M_{wz}M_{zz}^{-1}M_{zc} \\ M_{cz}M_{zz}^{-1}M_{zw} & M_{cz}M_{zz}^{-1}M_{zc} \end{bmatrix}.$$

Using the partitioned inversion formula, it follows that

$$V_\alpha = \left\{ M_{wz}M_{zz}^{-1}M_{zw} - M_{wz}M_{zz}^{-1}M_{zc} (M_{cz}M_{zz}^{-1}M_{zc})^{-1} M_{cz}M_{zz}^{-1}M_{zw} \right\}^{-1}.$$

Since $zM_{zz}^{-1}M_{zc}$ is the (population) projection of c on z and $z = [c', h', z_2']'$, it follows that $c = zM_{zz}^{-1}M_{zc}$ and

$$V_\alpha = \sigma^2(M_{wz}M_{zz}^{-1}M_{zw} - M_{wc}M_{cc}^{-1}M_{cw})^{-1} = \frac{\sigma^2}{\sigma_w^2} \left(\frac{1}{R_{w,z}^2 - R_{w,c}^2} \right), \quad (12)$$

where σ_w^2 denotes the variance of w . Similarly, we have

$$V_\alpha^{(1)} = \frac{\sigma^2}{\sigma_w^2} \left(\frac{1}{R_{w,z_1}^2 - R_{w,c}^2} \right). \quad (13)$$

The result then follows from (12)–(13).

Proof of Propositions 2 and 3: From Newey and Smith (2004)[Theorem 4.5] and the discussion below on p.230, it follows that, under our assumptions, the second order biases

are as follows:

$$\begin{aligned} \text{bias}(\hat{\theta}_T) &= (q - p - 1)V_\theta E[x_t u_t] / \sigma^2 T \\ \text{bias}(\hat{\theta}_{IM}) &= -V_\theta E[x_t u_t] / T. \end{aligned}$$

The results follow from $E[x_t u_t] = [\sigma_{uv}, 0_{1 \times (p-1)}]$ where $0_{1 \times (p-1)}$ is the $1 \times (p-1)$ null vector, and (12).

References

- Appleby, J., and Boyle, S. (2000). ‘B Liar’s billions: where will he find the money for the NHS?’, *British Medical Journal*, March 25; 320(7238): 865–867.
- Breusch, T., Qian, H., Schmidt, P., and Wyhowski, D. (1999). ‘Redundancy of moment conditions’, *Journal of Econometrics*, 91: 89–111.
- Brown, B. W., and Newey, W. K. (2002). ‘Generalized Method of Moments, efficient bootstrapping, and improved inference’, *Journal of Business and Economic Statistics*, 20: 507–517.
- Chamberlain, G. (1987). ‘Asymptotic efficiency in estimation with conditional moment restrictions’, *Journal of Econometrics*, 34: 305–334.
- Cressie, N., and Read, T. R. C. (1984). ‘Multinomial goodness-of-fit tests’, *Journal of the Royal Statistical Society, series B*, 46: 440–464.
- DOH (2005). ‘Resource Allocation: Weighted Capitation Formula’, Fifth Edition.
- Golan, A. (2006). ‘Information and entropy econometrics - a review and synthesis’, *Foundations and Trends in Econometrics*, 2: 1–145.
- Hall, A. R. (2005). *Generalized Method of Moments*. Oxford University Press, Oxford, U.K.
- Hall, A. R., Inoue, A., Jana, K., and Shin, C. (2008). ‘Information in Generalized Method of Moments estimation and entropy based moment selection’, *Journal of Econometrics*, 138: 488–512.
- Hall, P., and Presnell, B. (1999). ‘Intentionally biased bootstrap methods’, *Journal of the Royal Statistical Society B*, 99: 143–158.
- Hansen, L. P. (1982). ‘Large sample properties of Generalized Method of Moments estimators’, *Econometrica*, 50: 1029–1054.

- (1985). ‘A method of calculating bounds on the asymptotic covariance matrices of generalized method of moments estimators’, *Journal of Econometrics*, 30: 203–238.
- Kitamura, Y. (2007). ‘Empirical Likelihood Methods in Econometrics: Theory and Practice’, in R. Blundell, W. K. Newey, and T. Personn (eds.), *Advances in Economics and Econometrics: Ninth World Congress of the Econometric Society*. Cambridge University Press, Cambridge.
- Kitamura, Y., and Stutzer, M. (1997). ‘An information-theoretic alternative to generalized method of moments estimation’, *Econometrica*, 65: 861–874.
- Martin, S., Rice, N., and Smith, P. C. (2008). ‘Does health care spending improve health outcomes? Evidence from English programme budgeting data’, *Journal of Health Economics*, 27: 826–842.
- Nagar, A. L. (1959). ‘The Bias and Moment Matrix of the General k-Class Estimators of the Parameters in Simultaneous Equations’, *Econometrica*, 27(4): pp. 575–595.
- Nelson, C. R., and Startz, R. (1990). ‘The distribution of the instrumental variables estimator and its t ratio when the instrument is a poor one’, *Journal of Business*, 63: S125–S140.
- Newey, W. K. (1993). ‘Efficient estimation of models with conditional moment restrictions’, in G. S. Maddala, C. R. Rao, and H. D. Vinod (eds.), *Handbook of Statistics*, vol. 11, pp. 419–454. Elsevier Science Publishers, Amsterdam, The Netherlands.
- Newey, W. K., and Smith, R. J. (2004). ‘Higher order properties of GMM and generalized empirical likelihood estimators’, *Econometrica*, 72: 219–255.
- ONS (2006). ‘Public Service Productivity: Health’, Office of National Statistics, Economic Trends 628 March 2006.
- Owen, A. (2001). *Empirical Likelihood*. Chapman Hall/CRC, New York, NY, USA.

- Qin, J., and Lawless, J. (1994). ‘Empirical likelihood and general estimating equations’, *Annals of Statistics*, 22: 300–325.
- Smith, R. (2011). ‘GEL Criteria for Moment Condition Models’, *Econometric Theory*, 27: 1192–1235.
- Smith, R. J. (1997). ‘Alternative semi-parametric likelihood approaches to generalized method of moments estimation’, *Economics Journal*, 107: 503–519.
- Staiger, D., and Stock, J. H. (1997). ‘Instrumental variables regression with weak instruments’, *Econometrica*, 65: 557–586.
- Stock, J. H., and Wright, J. H. (2000). ‘GMM with weak identification’, *Econometrica*, 68: 1055–1096.
- Stock, J. H., Wright, J. H., and Yogo, M. (2002). ‘A survey of weak instruments and weak identification in generalized method of moments’, *Journal of Business and Economic Statistics*, 20: 518–529.