

On the Brownian First-Passage Time Over a One-Sided Stochastic Boundary

G. PESKIR and A. N. SHIRYAEV

Let $B = (B_t)_{t \geq 0}$ be standard Brownian motion started at 0 under P , let $S_t = \max_{0 \leq r \leq t} B_r$ be the maximum process associated with B , and let $g : \mathbf{R}_+ \rightarrow \mathbf{R}$ be a (strictly) monotone continuous function satisfying $g(s) < s$ for all $s \geq 0$. Let τ be the first-passage-time of B over $t \mapsto g(S_t)$:

$$\tau = \inf \{ t > 0 \mid B_t \leq g(S_t) \} .$$

Let G be the function defined by:

$$G(y) = \exp \left(- \int_0^{g^{-1}(y)} \frac{ds}{s - g(s)} \right)$$

for $y \in \mathbf{R}$ in the range of g . Then, if g is increasing, we have:

$$\lim_{t \rightarrow \infty} \sqrt{t} P\{\tau \geq t\} = \sqrt{\frac{2}{\pi}} \left(-g(0) - \int_{g(0)}^{g(\infty)} G(y) dy \right)$$

and this number is finite. Similarly, if g is decreasing, we have:

$$\lim_{t \rightarrow \infty} \sqrt{t} P\{\tau \geq t\} = \sqrt{\frac{2}{\pi}} \left(-g(0) + \int_{g(\infty)}^{g(0)} G(y) dy \right)$$

and this number may be infinite. These results may be viewed as a *stochastic boundary* extension of some known results on the first-passage-time over deterministic boundaries. The method of proof relies on the classical Tauberian theorem and certain extensions of the Novikov-Kazamaki criteria for exponential martingales.

1. Introduction

In this paper we present the solution to the following problem. Let $B = (B_t)_{t \geq 0}$ be standard Brownian motion started at 0 under P , let $S_t = \max_{0 \leq r \leq t} B_r$ be the maximum process associated with B , and let $g : \mathbf{R}_+ \rightarrow \mathbf{R}$ be a (strictly) monotone continuous function satisfying $g(s) < s$ for all $s \geq 0$. Let τ be the first-passage-time of B over the stochastic boundary $t \mapsto g(S_t)$:

$$(1.1) \quad \tau = \inf \{ t > 0 \mid B_t \leq g(S_t) \}$$

(see Fig. 1 below). The problem we address is to compute the limit:

$$(1.2) \quad \lim_{t \rightarrow \infty} \sqrt{t} P\{\tau \geq t\}$$

and to express it in terms of function g .

AMS 1980 subject classifications. Primary 60G40, 60J65. Secondary 60G44, 60J25, 60J60.

Key words and phrases: Brownian motion, the first-passage-time, stochastic boundary, Novikov-Kazamaki criteria, Tauberian theorem, Girsanov measure change, local martingale, diffusion process. (Second edition) © goran@imf.au.dk




Figure 1: A schematic drawing of the Brownian motion B_t and the target process $g(S_t)$ with a decreasing function $s \mapsto g(s)$.

If $g(s) \equiv g(0) < 0$, it is well-known that:

$$(1.3) \quad \lim_{t \rightarrow \infty} \sqrt{t} P\{\tau \geq t\} = \sqrt{2/\pi} (-g(0)).$$

A more general problem was recently treated by Novikov [7] for deterministic one-sided boundaries of the form $t \mapsto g(t)$. The main novelty in our approach is that we allow the boundary to be *stochastic*. It turns out that such a reformulation of the problem enables one to compute the limit in (1.2) explicitly. The solution is presented in Theorem 2.1 below. The proof is divided into several parts of independent interest. For simplicity we work with Brownian motion but the methods used in principle could be extended to treat similar problems for general one-dimensional time-homogeneous diffusions.

Stopping times of the form (1.1) have been investigated in the past. Instead of going into a historical exposition on this extensive subject, we shall rather refer the reader to the paper of Azéma and Yor [1] where such stopping times are used to obtain a simple solution to the Skorokhod problem. In [1] one also finds the Laplace transform of τ , which (at least in principle) could lead to the explicit expression in (1.2). Such an approach, however, may have the drawback of computational complexity. In recent years stopping times of the form (1.1) gained a new attention by appearing as solutions to several optimal stopping problems [2], [5], [6], [9], [10].

The initial work on the problem above was motivated by the paper of Novikov [7]. From his work we learned about the useful role of the classical Tauberian theorem in such a context. At the same time we knew of the manuscript [3] where similar results were obtained for general continuous local martingales. While these general results could be used to handle the case above when g is increasing, they are not directly applicable to the more difficult case when g is decreasing. In this case we needed to use a refinement of the classical criteria for exponential martingales due to Novikov and Kazamaki (see [8] p.317-319). This refinement is in fact a new way of thinking of the exponential martingales which in essence relies on a simple Girsanov measure change.

2. The results and proofs

In this section we present the main result and proof. Before we state our main result, we want to make a few comments. Throughout we study the stopping time τ from (1.1). Given $p > 0$, below we shall characterize when $E(\tau^p)$ is finite (Corollary 2.2), and we shall compute $E(\tau)$ (Corollary 2.3), both in terms of g . It is well-known that in the context of one-sided boundaries the exponent $p = 1/2$ plays a special role. To relate the problem about the limit in (1.2) with the integrability properties of τ , the reader should note the following two facts which are easily verified by using standard arguments:

$$(2.1) \quad \limsup_{t \rightarrow \infty} \left(\sqrt{t} P\{\tau \geq t\} \right) < \infty \Rightarrow E(\tau^{1/2-\varepsilon}) < \infty \quad (\forall \varepsilon > 0)$$

$$(2.2) \quad E(\tau^{1/2}) < \infty \Rightarrow \lim_{t \rightarrow \infty} \left(\sqrt{t} P\{\tau \geq t\} \right) = 0 .$$

The converse implications are not fulfilled in general. Our main result may be now stated as follows.

Theorem 2.1

Let $B = (B_t)_{t \geq 0}$ be standard Brownian motion started at 0 under P , let $S_t = \max_{0 \leq r \leq t} B_r$ be the maximum process associated with B , and let $g : \mathbf{R}_+ \rightarrow \mathbf{R}$ be a strictly monotone continuous function satisfying $g(s) < s$ for all $s \geq 0$.

Let τ be the first-passage-time of B over $t \mapsto g(S_t)$:

$$(2.3) \quad \tau = \inf \{ t > 0 \mid B_t \leq g(S_t) \}$$

and let G be the function defined by:

$$(2.4) \quad G(y) = \exp \left(- \int_0^{g^{-1}(y)} \frac{ds}{s - g(s)} \right)$$

for $y \in \mathbf{R}$ in the range of g .

If g is increasing, then we have:

$$(2.5) \quad \lim_{t \rightarrow \infty} \sqrt{t} P\{\tau \geq t\} = \sqrt{\frac{2}{\pi}} \left(-g(0) - \int_{g(0)}^{g(\infty)} G(y) dy \right)$$

and this number is finite.

If g is decreasing, then we have:

$$(2.6) \quad \lim_{t \rightarrow \infty} \sqrt{t} P\{\tau \geq t\} = \sqrt{\frac{2}{\pi}} \left(-g(0) + \int_{g(\infty)}^{g(0)} G(y) dy \right)$$

and this number may be infinite.

Proof. 1. In order to compute the limits in (2.5) and (2.6) we shall use a classical Tauberian (Abelian) theorem (see [4] p.442-448) which states that for any monotone $f : \mathbf{R}_+ \rightarrow \mathbf{R}_+$ we have:

$$(2.7) \quad \lim_{\lambda \rightarrow 0} \left(\sqrt{\lambda} \int_0^{\infty} e^{-\lambda t} f(t) dt \right) = L \iff \lim_{t \rightarrow \infty} \left(\sqrt{t} f(t) \right) = \frac{L}{\sqrt{\pi}}.$$

We shall use this result with $f(t) = P\{\tau \geq t\}$. Then we have:

$$(2.8) \quad \sqrt{\lambda} \int_0^{\infty} e^{-\lambda t} f(t) dt = E \left(\frac{1 - e^{-\lambda \tau}}{\sqrt{\lambda}} \right)$$

for all $\lambda > 0$. Hence we see that the problem reduces to compute the limit:

$$(2.9) \quad \lim_{\lambda \rightarrow 0} E \left(\frac{1 - e^{-\lambda \tau}}{\sqrt{\lambda}} \right).$$

2. To compute this limit, consider the local martingale:

$$(2.10) \quad M_t = -B_{t \wedge \tau}$$

and the exponential local martingale associated with it:

$$(2.11) \quad \mathcal{E}_\lambda(M)_t = \exp \left(\lambda M_t - \frac{\lambda^2}{2} \langle M \rangle_t \right)$$

where $\lambda > 0$. It will be clear from our proof below that we may, and in the sequel do assume, that $P\{\tau < \infty\} = 1$. Then clearly $M_\infty := \lim_{t \rightarrow \infty} M_t = -B_\tau$ P -a.s. and $\langle M \rangle_\infty := \lim_{t \rightarrow \infty} \langle M \rangle_t = \tau < \infty$ P -a.s.

Suppose that M_∞ is integrable, and that the following two conditions are satisfied:

$$(2.12) \quad E \left(\mathcal{E}_\lambda(M)_\infty \right) = 1 \quad \text{for all } 0 < \lambda < \lambda_*$$

$$(2.13) \quad E \left(M_\infty^+ \mathcal{E}_\lambda(M)_\infty \right) \rightarrow E(M_\infty^+) \quad \text{as } \lambda \rightarrow 0$$

with some $\lambda_* > 0$. Then we claim that:

$$(2.14) \quad \lim_{\lambda \rightarrow 0} E \left(\frac{1 - e^{-\lambda \tau}}{\sqrt{\lambda}} \right) = \sqrt{2} E(M_\infty).$$

To prove (2.14) note that by (2.12) we have:

$$(2.15) \quad \begin{aligned} \frac{1}{\frac{\lambda}{\sqrt{2}}} E \left(1 - \exp \left(-\frac{\lambda^2}{2} \langle M \rangle_\infty \right) \right) &= \frac{\sqrt{2}}{\lambda} E \left(\mathcal{E}_\lambda(M)_\infty - \exp \left(-\frac{\lambda^2}{2} \langle M \rangle_\infty \right) \right) \\ &= \sqrt{2} E \left(\left(\frac{\exp(\lambda M_\infty) - 1}{\lambda} \right) \exp \left(-\frac{\lambda^2}{2} \langle M \rangle_\infty \right) \right) \end{aligned}$$

for all $0 < \lambda < \lambda_*$. To proceed introduce the following notation:

$$(2.16) \quad R_\lambda = \left(\frac{\exp(\lambda M_\infty) - 1}{\lambda} \right) \exp \left(-\frac{\lambda^2}{2} \langle M \rangle_\infty \right).$$

Then clearly $R_\lambda \rightarrow M_\infty$ as $\lambda \rightarrow 0$, and since $|e^x - 1| \leq x^+ e^x + x^-$ for all $x \in \mathbf{R}$, we have:

$$(2.17) \quad |R_\lambda| \leq \left(M_\infty^+ \exp(\lambda M_\infty) + M_\infty^- \right) \exp\left(-\frac{\lambda^2}{2} \langle M \rangle_\infty\right) \leq M_\infty^+ \mathcal{E}_\lambda(M)_\infty + M_\infty^-$$

for all $\lambda > 0$. Now note that $0 \leq M_\infty^+ \mathcal{E}_\lambda(M)_\infty \rightarrow M_\infty^+$ as $\lambda \rightarrow 0$, so the condition (2.13) is equivalent to the uniform integrability of $M_\infty^+ \mathcal{E}_\lambda(M)_\infty$ for $\lambda \rightarrow 0$. Thus from (2.17) we see that R_λ is uniformly integrable for $\lambda \rightarrow 0$, so by letting $\lambda \rightarrow 0$ in (2.15) we get (2.14).

3. The rest of the proof consists of two separate parts. The first part is to derive the distribution law of M_∞ , which will lead to a closed form expression for $E(M_\infty)$ in (2.14). The second part is to verify that (2.12) and (2.13) are fulfilled whenever M_∞ is integrable. Combining these two facts we will obtain (2.5) and (2.6).

To find the distribution law of M_∞ , recall that $M_\infty = -B_\tau = -g(S_\tau)$, so it will be sufficient to derive the distribution law of S_τ . For this, we shall use the fact that the process:

$$(2.18) \quad H(S_t) - (S_t - B_t) H'(S_t)$$

is a local martingale whenever $H \in C^1$ (see [1]). This is easily verified by Itô formula and in essence relies upon the fact that $dS_t = 0$ outside $X_t = S_t$ (see Fig. 2 below). We shall apply this fact with $H(s) = \int_0^s h(r) dr$ where $h: \mathbf{R}_+ \rightarrow \mathbf{R}_+$ is any continuous function which is zero at zero and outside a compact set. Then the process in (2.18) is uniformly bounded and therefore it is a uniformly integrable martingale. Thus by the optional sampling theorem we get:

$$(2.19) \quad E\left(H(S_\tau)\right) = E\left((S_\tau - B_\tau) H'(S_\tau)\right).$$

Integrating by parts, the left-hand side in (2.19) equals:

$$(2.20) \quad \begin{aligned} E\left(H(S_\tau)\right) &= \int_0^\infty H(s) dF_{S_\tau}(s) = H(s)F_{S_\tau}(s) \Big|_0^\infty - \int_0^\infty F_{S_\tau}(s) dH(s) \\ &= H(\infty) - \int_0^\infty F_{S_\tau}(s) dH(s) = \int_0^\infty (1 - F_{S_\tau}(s)) h(s) ds. \end{aligned}$$

On the other hand, the right-hand side is equal to:

$$(2.21) \quad \begin{aligned} E\left((S_\tau - B_\tau) H'(S_\tau)\right) &= E\left((S_\tau - g(S_\tau)) H'(S_\tau)\right) \\ &= \int_0^\infty (s - g(s)) h(s) dF_{S_\tau}(s). \end{aligned}$$

Since this is true for all h of the specified type, from (2.20) and (2.21) we find that:

$$(2.22) \quad \frac{dF_{S_\tau}(s)}{ds} = \frac{1 - F_{S_\tau}(s)}{s - g(s)}$$

for $s > 0$. The equation (2.22) is easily solved, and since $F_{S_\tau}(0) = 0$, this leads to:




Figure 2: A schematic drawing of the diffusion (B_t, S_t) with a decreasing target function $s \mapsto g(s)$. The diffusion (B_t, S_t) can increase in the second component (infinitesimally little) only after hitting the diagonal $x = s$.

$$(2.23) \quad F_{S_\tau}(s) = 1 - \exp\left(-\int_0^s \frac{dt}{t-g(t)}\right)$$

for $s \geq 0$ (see p. 258-264 in [8] and note in particular Exc. VI.5.9). The distribution law of $B_\tau = g(S_\tau)$ is now easily obtained from (2.23).

4. Having the distribution law of B_τ , we can compute $E(M_\infty) = -E(B_\tau)$ in (2.14). For this assume first that g is increasing. Then by (2.23) the integration by parts gives:

$$(2.24) \quad \begin{aligned} E(M_\infty) &= -E(B_\tau) = -\left(g(0) + \int_{g(0)}^{g(\infty)} P\{B_\tau \geq y\} dy\right) \\ &= -\left(g(0) + \int_{g(0)}^{g(\infty)} P\{g(S_\tau) \geq y\} dy\right) \\ &= -\left(g(0) + \int_{g(0)}^{g(\infty)} P\{S_\tau \geq g^{-1}(y)\} dy\right) \\ &= -\left(g(0) + \int_{g(0)}^{g(\infty)} (1 - F_{S_\tau}(g^{-1}(y))) dy\right) \\ &= -\left(g(0) + \int_{g(0)}^{g(\infty)} \exp\left(-\int_0^{g^{-1}(y)} \frac{ds}{s-g(s)}\right) dy\right) \end{aligned}$$

which together with (2.14) and (2.7)+(2.8) leads to (2.5) whenever conditions (2.12) and (2.13) are satisfied. If g is decreasing, then similarly:

$$\begin{aligned}
(2.25) \quad E(M_\infty) &= -E(B_\tau) = -\left(g(0) - \int_{g(\infty)}^{g(0)} P\{B_\tau \leq y\} dy\right) \\
&= -\left(g(0) - \int_{g(\infty)}^{g(0)} P\{g(S_\tau) \leq y\} dy\right) \\
&= -\left(g(0) - \int_{g(\infty)}^{g(0)} P\{S_\tau \geq g^{-1}(y)\} dy\right) \\
&= -\left(g(0) - \int_{g(\infty)}^{g(0)} (1 - F_{S_\tau}(g^{-1}(y))) dy\right) \\
&= -\left(g(0) - \int_{g(\infty)}^{g(0)} \exp\left(-\int_0^{g^{-1}(y)} \frac{ds}{s-g(s)}\right) dy\right)
\end{aligned}$$

which together with (2.14) and (2.7)+(2.8) leads to (2.6), whenever M_∞ is integrable, and conditions (2.12) and (2.13) are satisfied.

5. It should be noted that when g is increasing (thus bounded from below) the random variable $M_\infty = -B_\tau$ is integrable, and we have $0 \leq E(M_\infty) < \infty$. This is easily seen by analyzing the final expression in (2.24). To see this in another way, note that $M = (M_t)_{t \geq 0}$ is uniformly bounded from above by a constant, so that $(M_t^+)_{t \geq 0}$ is uniformly integrable. Moreover, since M is a continuous local martingale, there exists a sequence of (bounded) stopping times $\{\tau_n \mid n \geq 1\}$ such that $\tau_n \uparrow \infty$ and $(M_{t \wedge \tau_n})_{t \geq 0}$ is a (bounded) martingale. By the martingale property and Fatou's lemma we may conclude $E(M_\infty^-) \leq \lim_{t, n \rightarrow \infty} E(M_{t \wedge \tau_n}^-) = \lim_{t, n \rightarrow \infty} E(M_{t \wedge \tau_n}^+) = E(M_\infty^+) < \infty$.

If g is decreasing, however, the random variable $M_\infty = -B_\tau$ may not be integrable. We do have though that $E(M_\infty) \geq 0$ since M_∞ is non-negative. A simple truncation argument (by applying the preceding observations to the functions $g_n(s) = g(s) \vee (-n)$ which are bounded from below and in the limit tend to g as $n \rightarrow \infty$) it is clear that the limit on the left-hand side in (2.6) will be equal $+\infty$ if $E(M_\infty) = +\infty$. Thus to complete the proof it is enough to verify that conditions (2.12) and (2.13) are satisfied under the assumption that M_∞ is integrable.

In either of the cases it should be observed that since $M_\infty = -B_\tau = -g(S_\tau)$, the integrability of M_∞ implies that $S_\tau < \infty$ P -a.s., and therefore we have $P\{\tau < \infty\} = 1$.

6. To show that conditions (2.12) and (2.13) are satisfied, assume first that g is increasing. Then $M = (M_t)_{t \geq 0}$ is uniformly bounded from above by a constant, and therefore the exponential local martingale $\mathcal{E}_\lambda(M) = (\mathcal{E}_\lambda(M)_t)_{t \geq 0}$ is uniformly integrable for all $\lambda > 0$. Thus it is a martingale, and (2.12) is clearly satisfied. Moreover, since M_∞ is also bounded from above by a constant, we see that $(M_\infty^+ \mathcal{E}_\lambda(M)_\infty)_{\lambda > 0}$ is uniformly integrable, so that (2.13) is clearly valid as well. This completes the proof of (2.5) when g is increasing.

Assume now that g is decreasing. Consider the exponential martingale:

$$(2.26) \quad Z_t = Z_t^\lambda = \exp\left(-\lambda B_t - \frac{\lambda^2}{2} t\right)$$

for $\lambda > 0$. Denote $\mathcal{F}_t = \sigma(B_s \mid s \leq t)$ for $t \geq 0$, and let $\mathcal{F}_\infty = \sigma(\cup_{t \geq 0} \mathcal{F}_t)$. Let $\tilde{P} = \tilde{P}^\lambda$ denote the probability measure on \mathcal{F}_∞ determined by:

$$(2.27) \quad d\tilde{P}_t = Z_t dP_t$$

where \tilde{P}_t and P_t denote the restrictions of \tilde{P} and P to \mathcal{F}_t respectively. Then, under $\tilde{P} = \tilde{P}^\lambda$ with $\lambda > 0$, the process $\tilde{B}_t = \tilde{B}_t^\lambda = B_t + \lambda t$ is a standard Brownian motion. Moreover, it is well-known that (in the usual sense):

$$(2.28) \quad \tilde{E}\left(V 1_{\{\tau < \infty\}}\right) = E\left(V 1_{\{\tau < \infty\}} Z_\tau\right)$$

whenever V is \mathcal{F}_τ -measurable. This with $V \equiv 1$ shows that:

$$(2.29) \quad E(Z_\tau) = 1 \Leftrightarrow \tilde{P}\{\tau < \infty\} = 1.$$

Now note that in the notation of (2.12) above we have $Z_\tau = Z_\tau^\lambda = \mathcal{E}_\lambda(M)_\infty$. Thus (2.12) is equivalent to the fact that $\tau < \infty$ under $\tilde{P} = \tilde{P}^\lambda$ for all $0 < \lambda < \lambda_*$. To verify this note that:

$$(2.30) \quad \tau = \inf \left\{ t > 0 \mid \tilde{B}_t - \lambda t \leq g\left(\max_{0 \leq r \leq t} (\tilde{B}_r - \lambda r)\right) \right\}.$$

Since $t \mapsto \max_{0 \leq r \leq t} (\tilde{B}_r - \lambda r)$ is bounded from above \tilde{P} -a.s., and moreover $\tilde{B}_t - \lambda t \rightarrow -\infty$ \tilde{P} -a.s., we clearly see from (2.30) that $\tau < \infty$ under $\tilde{P} = \tilde{P}^\lambda$ for all $\lambda > 0$. Thus the condition (2.12) is fulfilled for all $\lambda > 0$.

To show (2.13) note by (2.28) that:

$$(2.31) \quad E\left(M_\infty^+ \mathcal{E}_\lambda(M)_\infty\right) = -E(B_\tau Z_\tau) = -\tilde{E}(B_\tau).$$

Motivated by (2.30) introduce the stopping time:

$$(2.32) \quad \tau_\lambda = \inf \left\{ t > 0 \mid B_t - \lambda t \leq g\left(\max_{0 \leq r \leq t} (B_r - \lambda r)\right) \right\}$$

for $\lambda > 0$. Then clearly:

$$(2.33) \quad Law(\tau_\lambda \mid P) = Law(\tau \mid \tilde{P})$$

$$(2.34) \quad Law(B_\tau \mid \tilde{P}) = Law(B_{\tau_\lambda} - \lambda \tau_\lambda \mid P)$$

since $B_\tau = \tilde{B}_\tau - \lambda \tau$. From (2.31) by (2.34) we get:

$$(2.35) \quad E\left(M_\infty^+ \mathcal{E}_\lambda(M)_\infty\right) = -\tilde{E}(B_\tau) = -E(B_{\tau_\lambda} - \lambda \tau_\lambda).$$

Now note that $\tau_\lambda \rightarrow \tau$ and $B_{\tau_\lambda} - \lambda \tau_\lambda \rightarrow B_\tau$ (both pointwise) when $\lambda \rightarrow 0$. Moreover, the sequence τ_λ is (pointwise) increasing for $\lambda \downarrow 0$. To see this, take and $\lambda' < \lambda$, then (since g is decreasing) at $t = \tau_{\lambda'}$ we have:

$$(2.36) \quad B_t - \lambda t \leq B_t - \lambda' t = g\left(\max_{0 \leq r \leq t} (B_r - \lambda' r)\right) \leq g\left(\max_{0 \leq r \leq t} (B_r - \lambda r)\right)$$

so we must have $\tau_\lambda \leq \tau_{\lambda'}$. Similarly, the sequence $B_{\tau_\lambda} - \lambda \tau_\lambda$ is (pointwise) decreasing for $\lambda \downarrow 0$. To see this, recall that:

$$(2.37) \quad B_{\tau_\lambda} - \lambda\tau_\lambda = g\left(\max_{0 \leq r \leq \tau_\lambda} (B_r - \lambda r)\right) .$$

Hence, since τ_λ is increasing, the maximum on the right-hand side is increasing for $\lambda \downarrow 0$, and the claim follows from the fact that g is decreasing. It remains to apply the monotone convergence theorem in (2.35) to conclude:

$$(2.38) \quad \lim_{\lambda \rightarrow 0} E\left(M_\infty^+ \mathcal{E}_\lambda(M)_\infty\right) = \lim_{\lambda \rightarrow 0} E(-B_{\tau_\lambda} + \lambda\tau_\lambda) = E(-B_\tau) = E(M_\infty^+) .$$

This establishes (2.13) in the case g is decreasing, and the proof of the theorem is complete. \square

Remark 1. In the proof above we consider strictly monotone g for simplicity. The proof can be generalized to treat more general functions g at the expense of technical complexity. A similar remark may be directed to the underlying Brownian motion process. We feel that the proof could be extended to treat similar problems for general one-dimensional time-homogeneous diffusions (or even for certain discontinuous processes like Levy's processes for instance).

Remark 2. Note that taking $g_n(s) = g(0) + s/n$ in (2.5) and letting $n \rightarrow \infty$, in the limit we obtain (1.3). A similar approximation procedure can be applied in (2.6). There one should rather use $g_n(s) = g(0) - \sqrt{s}/n$, since the right-hand side in (2.6) may be infinite for "too small" g . To experience the right size for g , note that if $g(s) = g(0) - s^\gamma$ with $\gamma > 0$, then the right-hand side in (2.6) is finite if and only if $0 < \gamma < 1$. Moreover, if $g(s) = g(0) - \varepsilon s$ for some $\varepsilon > 0$, then the right-hand side in (2.6) is infinite no matter how small ε is. These facts are easily obtained by analyzing the integral in (2.6). To conclude we may summarize that the functions g for which the limit in (2.6) is finite are of the form $g(s) = O(s^\gamma)$ for some $0 < \gamma < 1$.

Corollary 2.2

Let $B = (B_t)_{t \geq 0}$ be standard Brownian motion started at 0 under P , let $S_t = \max_{0 \leq r \leq t} B_r$ be the maximum process associated with B , and let $g : \mathbf{R}_+ \rightarrow \mathbf{R}$ be a strictly monotone continuous function satisfying $g(s) < s$ for all $s \geq 0$.

Let τ be the first-passage-time of B over $t \mapsto g(S_t)$:

$$(2.39) \quad \tau = \inf \{ t > 0 \mid B_t \leq g(S_t) \}$$

and let $p > 0$ be given and fixed.

If g is increasing, then $E(\tau^p) < \infty$ if and only if:

$$(2.40) \quad \int_0^\infty y^{2p-1} \exp\left(-\int_0^y \frac{ds}{s-g(s)}\right) dy < \infty .$$

If g is decreasing, then $E(\tau^p) < \infty$ if and only if (2.40) holds and:

$$(2.41) \quad \int_{-g(0)}^{-g(\infty)} y^{2p-1} \exp\left(-\int_0^{g^{-1}(-y)} \frac{ds}{s-g(s)}\right) dy < \infty .$$

Proof. By the Burkholder-Davis-Gundy inequality we know that $E(\tau^p) < \infty$ if and only

if $E(\max_{0 \leq t \leq \tau} |B_t|^{2p}) < \infty$. If g is increasing, the latter condition is clearly equivalent to $E(S_\tau)^{2p} < \infty$, which is by (2.23) expressed as in (2.40). If g is decreasing, the latter condition is clearly equivalent to both $E(S_\tau)^{2p} < \infty$ and $E(-B_\tau)^{2p} = E(-g(S_\tau))^{2p} < \infty$, which is again by (2.23) expressed as in (2.40) and (2.41). This completes the proof. \square

We shall conclude our exposition by indicating how one can compute the expectation of τ from (1.1) in a simple way when the underlying Brownian motion process starts at any point.

Corollary 2.3

Suppose we are given a standard Brownian motion $B = (B_t)_{t \geq 0}$ started at some $x \geq 0$ under P_x . Let $S_t = \max_{0 \leq r \leq t} B_r$ be the maximum process associated with B , and let $g : \mathbf{R}_+ \rightarrow \mathbf{R}$ be a (strictly) increasing continuous function satisfying $g(s) < s$ for all $s \geq 0$. Let τ be the first-passage-time of B over $t \mapsto g(S_t)$:

$$(2.42) \quad \tau = \inf \{ t > 0 \mid B_t \leq g(S_t) \}.$$

Then the following formula is valid:

$$(2.43) \quad E_x(\tau) = \int_x^\infty \frac{(x-g(t))^2}{t-g(t)} \exp\left(-\int_x^t \frac{ds}{s-g(s)}\right) dt$$

whenever $E_x(\tau)$ is finite, i.e. the condition (2.40) holds.

Proof. By a standard argument it is enough to prove (2.43) when the integral on the right-hand side is finite. Then $(B_t - x)_{t \geq 0}$ is a standard Brownian motion started at 0 under P_x , and clearly:

$$(2.44) \quad E_x(\tau) = E_x |B_\tau - x|^2 = E_0 |B_\tau|^2 = E_0 |h(S_\tau)|^2$$

where $h(s) = g(s + x) - x$. The distribution law of S_τ is then given by (2.23) where g is to be replaced by h . This easily leads to the formula (2.43) and completes the proof. \square

REFERENCES

- [1] AZEMA, J. and YOR, M. (1979). Une solution simple au probleme de Skorokhod. *Sem. Probab. XII, Lecture Notes in Math.* 721, Springer (90-115) & (625-633).
- [2] DUBINS, L. E. SHEPP, L. A. and SHIRYAEV, A. N. (1993). Optimal stopping rules and maximal inequalities for Bessel processes. *Theory Probab. Appl.* 38 (226-261).
- [3] ELWORTHY, K. L. LI X. M. and YOR, M. (1997). On the tails of the supremum and the quadratic variation of strictly local martingales. *Sem. Probab. XXXI, Lecture Notes in Math.* 1655, Springer-Verlag, (120-133).
- [4] FELLER, W. (1971). *An introduction to probability theory and its applications.* John Wiley & Sons, New York (Second edition).
- [5] GRAVERSEN, S. E. and PESKIR, G. (1995). On Doob's maximal inequality for Brownian

- motion. *Research Report* No. 337, *Dept. Theoret. Statist. Aarhus* (13 pp). *Stochastic Process. Appl.* 69 (111-125) 1997.
- [6] GRAVERSEN, S. E. and PESKIR, G. (1996). Optimal stopping in the $L \log L$ -inequality of Hardy and Littlewood. *Research Report* No. 360, *Dept. Theoret. Statist. Aarhus* (12 pp). *Bull. London Math. Soc.* 30 (171-181) 1998.
- [7] NOVIKOV, A. A. (1996). Martingales, Tauberian theorem and strategies of gambling. *Theory Probab. Appl.* 41 (716-729).
- [8] REVUZ, D. and YOR, M. (1994). *Continuous Martingales and Brownian Motion*. Springer-Verlag.
- [9] SHEPP, L. A. and SHIRYAEV, A. N. (1993). The Russian option: Reduced regret. *Ann. Appl. Probab.* 3 (631-640).
- [10] SHIRYAEV, A. N. (1996). Minimax optimality of the method of cumulative sums (cusum) in the case of continuous time. *Uspekhi Mat. Nauk* Vol. 51, No. 4 (173-174) (Russian). *Russian Math. Surveys* Vol. 51, No. 4 (750-751) (English translation).

Goran Peskir
Department of Mathematical Sciences
University of Aarhus, Denmark
Ny Munkegade, DK-8000 Aarhus
home.imf.au.dk/goran
goran@imf.au.dk

Albert N. Shiryaev
Steklov Mathematical Institute
Gubkina str. 8
117966 Moscow
Russia
shiryaev@mi.ras.ru