

ON THE STRENGTH OF SOME TOPOLOGICAL LATTICES

MARCUS TRESSL

– *Dedicated to Murray Marshall* –

ABSTRACT. We study the model theoretic strength of various lattices that occur naturally in topology, like closed (semi-linear or semi-algebraic or convex) sets. The method is based on weak monadic second order logic and sharpens previous results by Grzegorzczuk. We also answers a question of Grzegorzczuk on the 'algebra of convex sets'.

CONTENTS

1. Introduction	1
2. The weak monadic structure of a first order structure	3
3. Finite unions of closed intervals	8
4. Defining closed and bounded intervals	10
5. Linear spaces and convex sets	12
6. Lattices of closed definable sets in the ordered context	16
7. Zariski closed sets and p-adic sets	20
8. Grzegorzczuk's paper	21
References	22

1. INTRODUCTION

A lattice in this note is a partially ordered set that has suprema and infima for finite nonempty subsets. We are concerned with lattices arising naturally in topology, e.g. as zero sets of certain algebraic structures of continuous functions on topological spaces. Examples are:

- Closed subsets of \mathbb{R}^n : zero sets of continuous functions $\mathbb{R}^n \rightarrow \mathbb{R}$.
- Zariski closed subsets of \mathbb{C}^n : zero sets of polynomial maps $\mathbb{C}^n \rightarrow \mathbb{C}^m$.
- Linear subspaces of K^n for a field K : zero sets of linear functions $K^n \rightarrow K^m$.
- Convex hulls of finite sets in K^n for some ordered field K : zero sets of convex continuous semi-linear functions $K^n \rightarrow K$.

Date: May 13, 2018.

2010 Mathematics Subject Classification. Primary: 03C64, Secondary: 03G10, 06B99.

Key words and phrases. Lattices, decidability, topology, model theory, interpretation.

We call lattices of such kind *topological lattices*; there is no formal definition given, but we use the terminology to explain ideas.¹ Given such a lattice L , we are interested in the strength of the first order structure (L, \leq) , i.e., we want to know what is interpretable in the partially ordered set (L, \leq) . An important sub-question is: When is (L, \leq) decidable? Many of such decidability questions have been addressed in A. Grzegorzczuk's paper on *Undecidability of some topological theories*, [Grzego1951]. We revisit parts of Grzegorzczuk's paper, improve some results as well as answer questions from [Grzego1951, end of §5]. I have included footnotes en route when there are strong contact points to [Grzego1951].

On the other hand, our method and our context is different from those in [Grzego1951]. We study topological lattices with the aid of weak monadic second order structures: These are first order structures where it is allowed to quantify over finite subsets of the universe (however we will use a first order implementation of this second order structure, see section 2).

Conceptually, our results say that a topological lattice is decidable, when it comes from a 1-dimensional topological space, or, when the space has few connected sets; otherwise it interprets $(\mathbb{N}, +, \cdot)$ (or its theory).

Here is an outline of the paper. Our main focus are topological lattices related to some ordered context, although we talk briefly about Zariski closed sets and the p -adics in section 7. For this introduction, think of a given topological space X , like \mathbb{R}^n or \mathbb{Q}^n and the lattice as a lattice of closed subsets of X , like closed semi-linear sets, or closed connected 1-dimensional semi-algebraic subsets of \mathbb{R}^2 , or the lattice of all closed subsets of X .

In many lattices L of closed sets one can find isomorphic (in a sense to be made precise in the text) copies of a unit interval I , like the unit interval in \mathbb{R} or in \mathbb{Q} . This is explained in section 4. The overall strategy is to analyse which structure is imposed by the ambient L on these intervals. It turns out that L imposes the weak monadic order structure of the total order on I . More precisely L defines the partially ordered set of finite unions of closed subintervals $[a, b]$ of I . In section 3 we will study this partially ordered set first and show that it is essentially the same as the ordered set I extended by quantification over finite sets (cf. 3.1). Using Rabin's landmark result on the decidability of the monadic second order theory of two successor functions (cf. [Rabin1969]) it follows that this extension is decidable (cf. 3.6).

In particular topological lattices are decidable, when the space itself is 1-dimensional (in an informal sense). If the space is not 1-dimensional then one can frequently find a first order definition of "equal size" of finite sets (these are present as elements in the lattice). This phenomenon suffices to interpret arithmetic in the lattice (cf. 3.2).

In section 6 then, the method leads to interpretations of $(\mathbb{N}, +, \cdot)$ in various topological lattices that are not based on a 1-dimensional space, see 6.3, 6.7. A particular case is given by lattices of convex (not necessarily closed) sets in some K^n , K an ordered field. Here we show, using the strategy above, that L indeed interprets the ordered field K expanded by integers. This answers a question of Grzegorzczuk at the end of [Grzego1951, §5].

¹In the literature, a topological lattice is a lattice equipped with a topology. Our lattices carry natural Vietoris-like topologies, but this does not play a role here.

Our standard model theoretic set up follows Hodges book [Hodges1993]. The main reference for o-minimal structures is [vdDries1998]; however the text does not strongly depend on o-minimality and one can just read “semi-linear” or “semi-algebraic” instead of “o-minimal”.

2. THE WEAK MONADIC STRUCTURE OF A FIRST ORDER STRUCTURE

A common feature of the topological lattices that we are concerned with is that they frequently interpret the (weak) monadic second order logic of a structure. We will encompass this logic itself in a first order structure and first look at some model theory of these structures.

2.1. Notation. Let $L = (L, \leq, \wedge, \vee, \perp)$ be a lattice with smallest element \perp . An **atom** of L is an element that is minimal among all elements different from \perp . We say that an element $x \in L$ is **L -connected** (or just connected if L is clear from the context) if there are no $y, z \in L$, with $y \wedge z = \perp \neq y, z$ and $y \vee z = x$.

For example, if L is the poset of linear subspaces of \mathbb{C}^n . Then L is a (bounded) lattice with $a \wedge b = a \cap b$, $a \vee b = a + b$ (sum of linear spaces), $\perp = \{0\}$. Then the atoms of L are the one dimensional subspaces and these are the only L -connected elements of L . Notice that L is not distributive.

We will frequently work with a sublattice of some powerset. In this case we write $\subseteq, \cap, \cup, \emptyset$ instead of $\leq, \wedge, \vee, \perp$.

2.2. Definition. Let \mathcal{L} be a language (by which we always mean a first order language). Let $\underline{\mathcal{L}}$ be the language that extends the language of lattices ($\subseteq, \cap, \cup, \emptyset$) with bottom element by adjoining an n -ary predicate symbol Z_φ for every \mathcal{L} -formula φ in n variables.

Let M be an \mathcal{L} -structure. The **monadic second order structure of M** is defined as the following first order structure $\text{MSO}(M)$ in the language $\underline{\mathcal{L}}$: $\text{MSO}(M)$ is the expansion of the lattice $\mathcal{P}(M)$ (the powerset of M) by the following sets: For each \mathcal{L} -formula φ in n variables the predicate Z_φ is interpreted in $\text{MSO}(M)$ as

$$\{(\{z_1\}, \dots, \{z_n\}) \mid (z_1, \dots, z_n) \in M, M \models \varphi(z_1, \dots, z_n)\} \subseteq \mathcal{P}(M)^n.$$

We identify an element $m \in M$ with the atom $\{m\}$ of $\mathcal{P}(M)$. In particular M is considered as a subset of $\mathcal{P}(M)$ and more generally any subset Z of M^n is identified with the subset $\{(\{z_1\}, \dots, \{z_n\}) \mid (z_1, \dots, z_n) \in Z\}$ of $\mathcal{P}(M)^n$.² Notice that then, elements of M are, a priori, not 0-definable in $\text{MSO}(M)$.

The **weak monadic second order structure of M** is defined to be the substructure $W(M)$ of $\text{MSO}(M)$ induced on the finite subsets of M . Explicitly, $W(M)$ is the first order structure expanding the lattice of finite subsets of M by the 0-definable (in M) subsets of M^n (where again we identify M with $\{\{m\} \mid m \in M\} \subseteq \mathcal{P}(M)$).

2.3. We mainly work with $W(M)$ and record a few basic properties and conventions.

- (i) When dealing with definable subsets of $W(M)$ we use a human readable format. This mainly concerns subsets of M^n when M^n is considered as a subset of $\mathcal{P}(M)^n$. For example if $M = (M, \cdot)$ is a semigroup, then we still

² Another set up could be to introduce a new sort for the powerset of M and the element relation between M and its powerset. We choose a single sorted set up, since this fits better with applications later.

write $x \cdot y$ for the partial function $W(M) \times W(M) \rightarrow W(M)$ defined on $M \times M \subseteq W(M) \times W(M)$.

- (ii) The structure M is 0-definable in $W(M)$ as follows immediately from the definition of $W(M)$.
- (iii) For a map $f : M \rightarrow N$ of \mathcal{L} -structures, let $W(f) : W(M) \rightarrow W(N)$ be the map that sends a finite subset of M to its image under f . If f is elementary then clearly $W(f)$ is an $\underline{\mathcal{L}}$ -embedding, but it is in general not elementary.³ On the other hand, if $W(f)$ is elementary then f is elementary by (ii).
- (iv) If \mathcal{N} is an elementary $\underline{\mathcal{L}}$ -substructure of $W(M)$, then the intersection N of \mathcal{N} with M (recall that M is identified with the set of atoms) is an elementary substructure of M and $\mathcal{N} = W(N)$: Let N be the set of atoms of \mathcal{N} . Each element of N is an atom of $W(M)$, hence $N \subseteq M$ (using the convention of identifying atoms with elements of M). Now by (ii), M (as a structure) is defined in $W(M)$. The definition, when applied to \mathcal{N} , creates an \mathcal{L} -structure on N . We consider N equipped with this structure. Now, $W(M)$ 'knows' that it is the monadic second order structure of an \mathcal{L} -structure and so does \mathcal{N} : If φ is an \mathcal{L} -formula in n variables, then using the definition from (ii), there is an $\underline{\mathcal{L}}$ -formula ψ in n -variables such that for all $\bar{a} \in M^n$ we have $M \models \varphi(\bar{a}) \iff W(M) \models \psi(\bar{a})$. By definition of N it follows for $\bar{a} \in N^n$ that $N \models \varphi(\bar{a}) \iff \mathcal{N} \models \psi(\bar{a})$. Since $\mathcal{N} \prec W(M)$, this shows that N is the induced structure of M on N and $\mathcal{N} = W(N)$ (and so $N \prec M$).
- (v) There are not many model theoretic properties that pass from \mathcal{L} -structures (or maps between them) to their weak monadic second order correspondents; the reason is that $W(M)$ has in general a much richer structure of definable sets than M , see for example 2.6. To deal with definability in $W(M)$ it is useful to think of a second order language; I comment on this now in order to help the reader parsing the text, without explicitly using it later on: Given a first order language \mathcal{L} , a second order formula of \mathcal{L} is built just like a first order formula, with the exception that we have a second kind of variables $X, Y, Z \dots$ and quantifiers $\forall X, \exists Y$ over these variables; further we may use atomic formulas of the form $X(v)$ expressing $v \in X$, for a second order variable X and a first order variable v . In a first order \mathcal{L} -structure M , the variables and their quantifiers in a second order formula are thought of as ranging over (special classes) of subsets of the universe. In $\text{MSO}(M)$ we can read second order quantification over subsets of M as first order quantification over the powerset of M . In $W(M)$ we can read second order quantification over finite subsets of M as first order quantification over $W(M)$. This means: we can explicitly translate second order formulas φ of \mathcal{L} into first order formulas $\tilde{\varphi}$ of $\underline{\mathcal{L}}$ such that φ is true in M at some tuple (where second order quantifiers and variables range over finite sets) iff $\tilde{\varphi}$ is true in $W(M)$ at that tuple.

Many undecidability results in this note rely on the undecidability of $W((\mathbb{N}, +))$.⁴ We address this first, and, for subsequent use, in a slightly more general setting. If $S = (S, \cdot)$ is a semi-group (i.e. \cdot is associative) then we write $X \cdot Y = \{x \cdot y \mid x \in X, y \in Y\}$ for $X, Y \subseteq S$ and $s^K = \{s^k \mid k \in K\}$ for $K \subseteq \mathbb{N}$. An element $s \in S$ is

³To see an example: If K is a field of characteristic 0, then, using results from [Bauval1985] one can interpret in $\underline{\mathcal{L}}$, only depending on $\text{Th}(K)$, the pair of fields (K, \mathbb{Q}) . But in general the theory of this pair changes when we replace K by an elementary extension.

⁴ $\mathbb{N} = \{1, 2, 3, \dots\}$.

called **torsion** if $s^{\mathbb{N}}$ is finite. Notice that $s^{\mathbb{N}}$ does not need to be cancellative. On the other hand, if s is not torsion, then $(s^{\mathbb{N}}, \cdot) \cong (\mathbb{N}, +)$.

2.4. Proposition. *If $S = (S, \cdot)$ is a semigroup, then the binary relation $t \in s^{\mathbb{N}}$ of S is 0-definable in $W(S)$, independently of S ; i.e., the defining formula is the same for all semigroups S .*

Proof. In this proof, “definable” means 0-definable in $W(S)$. We proceed in several steps.

- (a) Obviously, the map $W(S) \times W(S) \rightarrow W(S)$, $(X, Y) \mapsto X \cdot Y$ is definable. Further it is clear that the set of finite semigroups (by which we mean sub-semigroups of S) is definable. Consequently the torsion elements of S are definable by saying that $s \in X$ for some finite semigroup X .
- (b) The following property of (s, X) , $s \in S$, $X \in W(S)$ is definable:

$$(*) \quad s \text{ is not torsion and } \exists n \in \mathbb{N} : X = s^{\{1, \dots, n\}}.$$

Proof. It suffices to show that $(*)$ holds if and only if the following properties hold (as “torsion” and the other properties are 0-definable in $W(S)$):

$$s \in X, \text{ } s \text{ is not torsion and}$$

$$\exists x \in X \left(X \cap x \cdot X = \emptyset \text{ and } s \cdot (X \setminus \{x\}) \subseteq X \right).$$

If $(*)$ holds, then $x = s^n$ has the required properties as $s^{\mathbb{N}}$ is cancellative.

Conversely, suppose $s \in X$ is not torsion and there is some $x \in X$ with $X \cap x \cdot X = \emptyset$ and $s \cdot (X \setminus \{x\}) \subseteq X$. Let $n \in \mathbb{N}$ be maximal with $s^{\{1, \dots, n\}} \subseteq X$. Since $s^{n+1} \notin X$ and $s \cdot (X \setminus \{x\}) \subseteq X$ we must have $x = s^n$. Consequently, $X \cap s^n \cdot X = \emptyset$. Now take $y \in X$ and let $k \in \{0, \dots, n-1\}$ be maximal with $s^k \cdot y \in X$ (where $s^0 \cdot y$ stands for y). Since $s^n y \notin X$ we have $s^{k+1} y \notin X$ and therefore $s^k y = x$. Hence $s^k y = s^n$, which implies $y = s^{n-k} \in s^{\{1, \dots, n\}}$ by cancelation. \square

- (c) We can now define $t \in s^{\mathbb{N}}$ by saying
 - t is in the smallest finite semigroup containing s , or, there is some $X \in W(S)$ with $t \in X$ such that property $(*)$ holds for s and X .

\square

2.5. Corollary. *Let S be the semigroup $(\mathbb{N}, +)$. Then in $W(S)$ multiplication of natural numbers is 0-definable.*

Proof. By 2.4 we can define, without parameters, division of natural numbers by saying that $k|n$ if and only if n is in the (additive!) semigroup generated by k . It is routine to check that multiplication is 0-definable in $(\mathbb{N}, +, |)$. \square

2.6. For infinite abelian groups G with $p \cdot G = 0$, p prime, the structure $W(G)$ is also undecidable, but for a different reason: Let T be the common theory of all finite dimensional \mathbb{F}_p -vector spaces expanded by 5 subspaces. This is known to be undecidable by [Toffal1997, Example 2, p. 246]. Now notice that G is an \mathbb{F}_p -vector space and as G is infinite the set of finite dimensional \mathbb{F}_p -vector spaces is (up to isomorphism) a definable subset of $W(G)$. It is then not difficult to effectively construct for each sentence φ in the language of T a sentence $\tilde{\varphi}$ in the language of $W(G)$, such that

$$T \vdash \varphi \iff W(G) \models \tilde{\varphi}.$$

As T is undecidable, also $W(G)$ must be undecidable. We omit the details here as we do not use it later on. However it is worth mentioning because it has a nice application in our context:

If M is an infinite set (viewed as a first order structure in the empty language), then $W(W(M))$ is undecidable: The reason is that $W(M)$ expands an infinite abelian group of exponent 2, where addition is symmetric difference of finite sets. It should be said that the undecidability of T here is based on the insolvability of the word problem for the class of finite groups. I do not know if $W(W(M))$ interprets Peano arithmetic.

We see from 2.4 and 2.6 that $W(M)$ may become wild for fairly simple structures M . There is one notable exception:

2.7. Proposition. *Let T be an ω -categorical theory in a countable language.*

- (i) *If $M \prec N$ are models of T , then $W(M) \prec W(N)$.*
- (ii) *For every $M \models T$, M is stably embedded into $W(M)$. In fact, every $S \subseteq M^n$ that is definable in $W(M)$ with parameters $a_1, \dots, a_k \in W(M)$ is definable in M with parameters from $a_1 \cup \dots \cup a_k$.*

Proof. (i). First assume M is countable and N is strongly \aleph_1 -homogeneous. Take a countable elementary substructure \mathcal{N} of $W(N)$. By 2.3(iv), there is a countable $N_0 \prec N$ with $\mathcal{N} = W(N_0)$. Since $M, N_0 \prec N$ are countable they are isomorphic (by ω -categoricity) and there is an automorphism of N mapping M to N_0 . This automorphism then extends to an automorphism of $W(N)$ and this extension maps $W(M)$ to $W(N_0) = \mathcal{N} \prec W(N)$. It follows that $W(M) \prec W(N)$.

For the general case choose $N_1 \succ N$ strongly \aleph_1 -homogeneous and take a sentence φ in the language of \mathcal{L} with parameters from $W(M)$ with $W(M) \models \varphi$. Take a countable elementary substructure $\mathcal{M}_0 \prec W(M)$ containing the parameters of φ . By 2.3(iv), we know that $\mathcal{M}_0 = W(M_0)$ for some countable $M_0 \prec M$. By what we have shown above we get $W(N_1) \models \varphi$. But then $W(N) \models \varphi$, too: Otherwise, the same argument as for M would entail $W(N_1) \models \neg\varphi$.

(ii) Let $S \subseteq M^n$ be definable in $W(M)$ by an \mathcal{L} -formula $\varphi(\bar{x}, \bar{a})$, \bar{x} an n -tuple, $\bar{a} \in W(M)^k$. Let E be the union of the finite sets a_1, \dots, a_k . By (i) we may replace M by a countable elementary substructure containing E . Suppose S is not definable in M with parameters from E .

Claim. There are $\bar{b}, \bar{c} \in M^n$ such that $\bar{b} \in S$, $\bar{c} \notin S$ and $\text{tp}^M(\bar{b}/E) = \text{tp}^M(\bar{c}/E)$.

Proof of the claim. Since T is ω -categorical, $S_n^M(E)$ is finite by Ryll-Nardzewski ([Hodges1993, Theorem 7.3.1]), say $S_n^M(E) = \{q_1, \dots, q_m\}$ is of size m . For each $i \in \{1, \dots, m\}$ take an $\mathcal{L}(E)$ -formula $\psi_i(\bar{x})$ isolating q_i . Since we assume that S is not definable in M with parameters from E , the set S is not a finite union of sets defined by some $\psi_i(\bar{x})$. Hence for some i , both S and $M^n \setminus S$ hit $\psi_i[M^n]$. Take $\bar{b} \in S \cap \psi_i[M^n]$ and $\bar{c} \in \psi_i[M^n] \setminus S$. Then $\text{tp}^M(\bar{b}/E) = q_i = \text{tp}^M(\bar{c}/E)$, as required. \square

Now take $\bar{b}, \bar{c} \in M^n$ as in the claim. Then $\text{tp}^M(\bar{b}/E) = \text{tp}^M(\bar{c}/E)$. Since M is countable and T is ω -categorical, M is saturated and there is an E -automorphism σ of M with $\sigma(\bar{b}) = \bar{c}$ (cf. [Hodges1993, Cor. 7.3.3]). But then σ extends to an a_1, \dots, a_k -automorphism $\bar{\sigma}$ of $W(M)$ and this contradicts the assumption that S is defined in $W(M)$ by $\varphi(\bar{x}, \bar{a})$. \square

2.8. *Remark.* Let M be an ω -categorical structure in a relational language \mathcal{L} . If the universal theory T_{\forall} of T is finitely axiomatisable, then the age of M (i.e. the set of finite substructures of M , cf. [Hodges1993, p. 324]) is a definable set in $W(M)$: There is obviously an \mathcal{L} -formula that holds in $W(M)$ at just those $A \in W(M)$ for which T_{\forall} is true, when A is considered as a substructure of M . This applies, for example, to the countable atomless boolean algebra A (in the language $\{\leq\}$, say). Notice that $W(A)$ still is undecidable by 2.6.

The following consequence of 2.5 will be used several times later to check if a structure interprets Peano arithmetic, or even $(\mathbb{N}, +, \cdot)$.

2.9. **Proposition.** *Let M be an infinite set and let $E \subseteq W(M) \times W(M)$ be defined by*

$$E(a, b) \iff a \text{ and } b \text{ are disjoint and of the same size.}$$

Let $(W(M), E)$ be the expansion of $W(M)$ by E .

(i) $(W(M), E)$ defines (without parameters) the relation “ a and b have the same size” of $W(M)$. If we identify the equivalence classes with $\mathbb{N}_0 = \{0, 1, 2, \dots\}$, where $n \in \mathbb{N}_0$ stands for the size of a representative, then $(W(M), E)$ interprets addition of natural numbers on \mathbb{N}_0 without parameters.

(ii) Suppose N is an expansion of $(W(M), E)$ and $S(x, \bar{y})$ is a formula of N with the following properties:

(a) For each $\bar{b} \in W(M)^{\bar{y}}$ the set $\{\text{size}(a) \mid a \in W(M), N \models S(a, \bar{b})\}$ is finite.

(b) For all $k \in \mathbb{N}$ and all $n_1, \dots, n_k \in \mathbb{N}$ there is some $\bar{b} \in W(M)^{\bar{y}}$ such that

$$\{n_1, \dots, n_k\} = \{\text{size}(a) \mid a \in W(M), N \models S(a, \bar{b})\}.$$

Then N interprets $(\mathbb{N}, +, \cdot)$ without parameters. The interpretation only depends on S and not on M .

Notation: If N is an expansion of $(W(M), E)$ for which there is a formula $S(x, \bar{y})$ satisfying (a) and (b) then we say that N **defines finite sequences of integers**.

Proof. (i). Two finite subsets a, b of M have the same size if and only if $E(a \setminus b, b \setminus a)$. This condition is first order in $(W(M), E)$, hence we may define the relation “ a and b have the same size” of $W(M)$. We thus may replace E by this relation. We write $[a]$ for the equivalence class of a modulo E . Then the relation

$$A([a], [b], [c]) \iff \exists a', b', c' : E(a, a') \& E(b, b') \& E(c, c') \& a' \cap b' = \emptyset \& a' \cup b' = c'$$

defines in $(W(M), E)$ the graph of addition induced by $(\mathbb{N}_0, +)$ via the bijection

$$W(M)/E \longrightarrow \mathbb{N}_0; [a] \mapsto \text{size}(a).$$

(ii). This follows from (i) and 2.5, since (a) and (b) guarantee that N (or better, the expansion of $(W(M), E)$ by the predicate S) interprets $W(\mathbb{N}, +)$. \square

2.10. **Outline on how 2.9 will be applied.** An important lattice that appears in topologies related to orders is the lattice L_0 of finite unions of closed intervals of the form $[a, b]$ for $a, b \in T$, where T is a dense linear order. In 3.1 we will see that the poset (L_0, \subseteq) is bi-interpretable with a certain weak monadic second order structure $W(M)$ of some first order structure M with universe T . This will be applied later as follows: In a topological lattice L one can frequently interpret $W(M)$ through the lattice L_0 ; then if L is essentially not based on a 1-dimensional set (the precise formulation can be found in 6.3), one can define the expansion

$(W(M), E)$ of $W(M)$ as in 2.9. Condition (ii) of 2.9 for $(W(M), E)$ then comes for free from the presence of an order on T (see 3.2) and so by 2.9, (L, \subseteq) will interpret $(\mathbb{N}, +, \cdot)$ (or at least its theory when parameters are involved).

3. FINITE UNIONS OF CLOSED INTERVALS

First some notation. If $T = (T, \leq)$ is any totally ordered set, then consider the **betweenness relation** B defined by T : B is a ternary relation and $B(x, y, z)$ holds just if $x \leq y \leq z$ or $z \leq y \leq x$. Betweenness relations will show up axiomatically in 4.1 below. Currently there is no need to talk about this axiomatically. We write $[[x, y]] = [x, y] \cup [y, x]$, for the set of all $z \in T$ between x, y and similarly $((x, y))$ for the set of all $z \in T$ properly between x, y . Hence $z \in [[x, y]]$ just means $B(x, z, y)$ and this relation is 0-definable in the structure (T, B) .⁵

3.1. Proposition. *Let (T, \leq) be an infinite totally ordered set with betweenness relation B and let $L = (L, \subseteq)$ be the partially ordered set of finite unions of closed intervals of T of the form $[a, b]$ (which is a distributive lattice with bottom element). Let B be the betweenness relation of (T, \leq) and consider the first order structure (T, B) .*

- (i) *The poset L is interpretable in $W(T, B)$ without parameters.*
- (ii) *If T is densely ordered, then $W(T, B)$ is definable in the poset L without parameters and – using the interpretation from (i) – L and $W(T, B)$ are bi-interpretable.⁶*

The definition and the interpretation are independent of T .

Proof. We write $M = (T, B)$.

- (i). For finite sets $E, F, G \in W(M)$ consider the set

$$A_{E,F,G} := \bigcup \{ [[e, f]] \mid e \in E, f \in F, \\ [[e, f]] \cap E = \{e\}, [[e, f]] \cap F = \{f\}, [[e, f]] \cap G = \emptyset \}.$$

Clearly $A_{E,F,G} \in L$. Conversely, if $U \in L$, then there are elements

$$e_1 \leq f_1 < g_1 < e_2 \leq f_2 < g_2 < \dots < g_{n-1} < e_n \leq f_n$$

in T with $U = [e_1, f_1] \cup \dots \cup [e_n, f_n]$. Then $U = A_{E,F,G}$ with $E = \{e_1, \dots, e_n\}$, $F = \{f_1, \dots, f_n\}$ and $G = \{g_1, \dots, g_{n-1}\}$. We can then interpret L in $W(M)$ without parameters as follows: The universe is $W(M)^3$. Routine checking shows that the binary relation $A_{E,F,G} \subseteq A_{E',F',G'}$ of $W(M)^3$ is 0-definable in $W(M)$. Hence also the equivalence relation $(E, F, G) \sim (E', F', G') \iff A_{E,F,G} = A_{E',F',G'}$ of $W(M)^3$ is 0-definable in $W(M)$. Thus L is interpreted as $W(M)^3 / \sim$ together with the partial order induced by $A_{E,F,G} \subseteq A_{E',F',G'}$.

- (ii). Assume now that T is densely ordered. Then we can define that a set $C \in L$ is definably connected⁷: We say that it is not the disjoint join of two nonempty sets in L . Consequently we can also talk about definably connected components of

⁵Obviously (T, \geq) has the same betweenness relation B as (T, \leq) . Routine checking shows that these are the only orders with betweenness relation B . Conversely, for a first order definition of \leq in (T, B) one generally needs to name two elements: think of \mathbb{Q} with its betweenness relation and consider reflections $2q - x$ at $q \in \mathbb{Q}$: They preserve the betweenness relation and fix q .

⁶Hence after naming two singletons of T (seen as elements of L), $W(T, \leq)$ is definable in the poset L .

⁷Definably connected means definably connected in the 0-minimal structure (T, \leq)

$A \in L$. This shows that, as a set, $W(M)$ is 0-definable in L as those $A \in L$ for which every definably connected component is an atom of L .

Having identified the finite subsets of T as a 0-definable subset of L it suffices to define the betweenness relation on the atoms of L that is naturally given by T : Given atoms $\{a\}, \{b\}, \{c\}$ of L we say that $\{b\}$ is between $\{a\}$ and $\{c\}$ if $\{b\}$ is included in the smallest definably connected set A with $\{a\}, \{c\} \subseteq A$.

The bi-interpretability is now routine checking and left to the reader. Both methods (i) and (ii) are independent of T . \square

3.2. Proposition. *Let (T, \leq) be an infinite totally ordered set and let B be the betweenness relation defined by (T, \leq) . Let M be the structure (T, B) .*

Let $E \subseteq W(M) \times W(M)$ be defined by

$$E(a, b) \iff a \text{ and } b \text{ are disjoint and of the same size.}$$

- (i) *The structure $(W(M), E)$ defines without parameters and independently of (T, \leq) , finite sequences of integers in the sense of 2.9.*
- (ii) *The structure $(W(M), E)$ interprets $(\mathbb{N}, +, \cdot)$ without parameters and independently of (T, \leq) . More precisely, it interprets the natural definition of addition and multiplication on the equivalence classes of T modulo the relation that identifies sets of equal size.*

Since (T, B) is 0-definable in (T, \leq) items (i) and (ii) are also true for $M = (T, \leq)$.

Proof. Item (ii) follows from 2.9(ii) once we have shown (i). In order to see (i) we may assume by using 2.9(i) that E satisfies

$$E(a, b) \iff a \text{ and } b \text{ are of the same size.}$$

for all $a, b \in W(M)$.

We call $x, y \in T$ a jump of $b \in W(M)$ if $x, y \in b$, $x \neq y$ and $((x, y)) \cap b = \emptyset$. For $a, b, c \in W(M)$ we define a relation

$$S(a, b, c) \iff \exists x, y \in b \left(((x, y)) \cap b = \emptyset \ \& \ E(((x, y)) \cap c, a) \right).$$

Hence $S(a, b, c)$ holds just if there is a jump x, y of b such that the size of $((x, y)) \cap c$ is the size of a . Since the relation $z \in ((x, y))$ is definable in $M = (T, B)$ by $B(x, z, y) \ \& \ z \neq x, y$, it is clear that S is 0-definable in $(W(M), E)$. Further, for all $b, c \in M$, the set of E -equivalence classes of $a \in W(M)$ with $S(a, b, c)$ is finite: It consist of all the E -classes of $a \in W(M)$ such that the size of a is the size of $((x, y)) \cap c$ for some jump x, y of b . Hence 2.9(ii)(a) holds.

On the other hand, also 2.9(ii)(b) holds: If $n_1, \dots, n_k \in \mathbb{N}$, then we may obviously choose finite subsets b, c of T such that the cardinalities of the sets $((x, y)) \cap c$ are n_1, \dots, n_k , when (x, y) runs through the jumps of b . Hence $S(u, b, c)$ defines a set of representatives of the elements of $\{n_1, \dots, n_k\}$. \square

We turn to decidability of the lattice of finite unions of closed intervals.

3.3. Theorem. *Let $S2$ be the binary tree $2^{<\omega}$ together with the two successor functions $\sigma \mapsto \sigma \hat{\ }1$ and $\sigma \mapsto \sigma \hat{\ }0$. Then any expansion of $\text{MSO}(S2)$ by naming finitely many elements from $W(S2)$ is decidable.*

Proof. This is the main result in [Rabin1969], see [Rabin1969, Theorem 1.1] and [Rabin1969, Corollary 1.9] \square

3.4. *Remark.* The following binary relations on $S2$ are 0-definable in $W(S2)$:

- (i) $\sigma \leq \tau$ defined as ' τ extends σ '. (One can express that σ is in the smallest finite subset of $S2$ containing τ , that is closed under immediate predecessors.)
- (ii) $\sigma \preceq \tau$ defined as ' σ is to the right of τ ' in the natural horizontal order of the binary tree $2^{<\omega}$. Explicitly, this means $\sigma = \tau$, or, $\sigma^{\wedge}1 \leq \tau$, or, $\tau^{\wedge}0 \leq \sigma$, or σ and τ are incomparable for \leq and if γ is the infimum of σ and τ for \leq , then $\gamma^{\wedge}0 \leq \sigma$.
- (iii) It is clear that the poset $(S2, \preceq)$ is a countable dense linear order without endpoints. Hence $(S2, \preceq) \cong (\mathbb{Q}, \leq)$ and therefore we may consider $W(S2)$ as an expansion of $W(\mathbb{Q}, \leq)$.

3.5. **Corollary.** *If T is a densely linearly ordered set, then any expansion of $W(T)$ by naming finitely many elements is decidable.*⁸

Proof. Since T is ω -categorical we may assume that T is countable, using 2.7(i). Then T is isomorphic to an interval of (\mathbb{Q}, \leq) with endpoints in \mathbb{Q} . Since all these intervals are parametrically definable in (\mathbb{Q}, \leq) we may assume that $T = (\mathbb{Q}, \leq)$. By 3.4(iii) it is therefore enough to show that any expansion of $W(S2)$ by naming finitely many elements is decidable. But this follows from 3.3, since $W(S2)$ is 0-definable in $\text{MSO}(S2)$: A subset S of $2^{<\omega}$ is infinite if and only if there is a nonempty subset Y of the down set generated by S for \leq such that for all $\sigma \in Y$ either $\sigma^{\wedge}1 \in Y$ or $\sigma^{\wedge}0 \in Y$. \square

3.6. **Corollary.** *Let T be a dense linear order. Any expansion of the lattice of finite unions of closed intervals of the form $[a, b]$ by finitely many parameters, is decidable.*

Proof. By 3.5 and 3.1. \square

3.7. *Remark.* The lattice $(\mathbb{Q} \times \mathbb{Q}, \leq)$ with component wise order is obviously definable in the totally ordered set (\mathbb{Q}, \leq) . However, the weak monadic second order structure of the lattice $(\mathbb{Q} \times \mathbb{Q}, \leq)$ is not interpretable in the weak monadic second order structure of (\mathbb{Q}, \leq) ; in other words, we can not talk about finite sets of pairs of rational numbers in the first order structure $(W(\mathbb{Q}, \leq))$: We know from 3.5 that $W(\mathbb{Q}, \leq)$ is decidable; however $W(\mathbb{Q} \times \mathbb{Q}, \leq)$ interprets the theory of $(\mathbb{N}, +, \cdot)$, see 6.7 (and 6.6).

4. DEFINING CLOSED AND BOUNDED INTERVALS

In order to apply the results from the previous section we need a method to define a totally ordered set T in a topological lattice. The lattice will then frequently define finite unions of closed intervals of T .

4.1. **Definition.** Let X be a set and B be a ternary relation on X . We say that B is a **bounded betweenness relation** if there are $a, b \in X$ such that

- (a) $B(a, x, y)$ and $B(x, y, b)$ define the same total order \leq on X with smallest element a and largest element b .
- (b) B is the betweenness relation of \leq .

⁸This is generally attributed to Läuchli, see [Laeuch1968], but I was unable to find a precise reference. So below is a proof relying on Rabin's work.

We write $[[x, y]] = \{z \in X \mid B(x, z, y)\}$, $((x, y)) = [[x, y]] \setminus \{x, y\}$, and $[[x, y)) = [[x, y]] \setminus \{y\}$. Then B is called **dense** if $((x, y)) \neq \emptyset$ for all $x \neq y$.

Clearly bounded (dense) betweenness relations are first order axiomatisable in the language that has a ternary relation symbol.

4.2. Definition of $I(x)$. Let $I(x)$ be the formula in the language of posets saying the following:

- I1. x is connected, not an atom.
- I2. The ternary relation of the set of atoms contained in x , defined as “ u is in the smallest connected element $\leq x$ containing v, w ”, is a bounded and dense betweenness relation.⁹
- I3. For all atoms $u \leq x$ and every $y \leq x$ with $u \not\leq y$ one of the following hold true:
 - (a) There are atoms $v, w \leq x$ such that $u \in ((v, w))$ and $((v, w))$ contains no atom $\leq y$.
 - (b) u is one of the boundary points for one/both of the two total orders given by the betweenness relation in I2, and there exists an atom $v \leq x$ such that $u \neq v$ and $[[u, v))$ contains no atom $\leq y$.

In both (a) and (b) the double bracketed expression refer to the betweenness relation given by I2.

The idea here is that for an element a in a lattice L (and assuming that L is atomic), $I(a)$ says that the set of atoms of L underneath a should be orderable as a dense linear order with endpoints, in such a way that the elements of L are (or, mimic) a basis of closed sets of the order topology. We use the formulation with betweenness relations as this avoids the use of parameters in applications.

4.3. Nota Bene. For $a \in L$ the formula $I(x)$ is true in L at a if and only if it is true in the lattice $\{b \in L \mid b \leq a\}$ at a .

4.4. Proposition. Let X be a Hausdorff space that has a countable basis and let L be a lattice of closed subsets of X with the following properties:

- (a) If $A \in L$ with $L \models I(A)$, then all closed subsets of A are in L
- (b) If $A \subseteq X$ is homeomorphic to $[0, 1]_{\mathbb{R}}$ then all closed subsets of A are in L .

(The largest such lattice is the set of all closed subsets of X , the smallest such lattice is the set of all closed subsets of X that are contained in a finite union of homeomorphic copies of $[0, 1]_{\mathbb{R}}$.)

Then $I(x)$ defines in L the set of all subsets of X that are homeomorphic to the closed unit interval of \mathbb{R} .

Proof. By 4.3 and (b), $I(x)$ is satisfied for subsets of X that are homeomorphic to the closed unit interval of \mathbb{R} . Conversely take $T \in L$ with $L \models I(T)$. Since T also has a countable basis we may use 4.3 again and assume $X = T$. Now condition (a) says that L is the lattice of all closed subsets of X , in particular L -connected is just connected in the usual sense of topology. We pick one of the two linear orders \sqsubseteq on X defined by the betweenness relation B asserted to exist by the formula $I(x)$. Since \sqsubseteq has B as its betweenness relation we know for all $a, b \in X$ that the smallest L -connected set from L containing $\{a, b\}$, is the interval $[a, b]_{\sqsubseteq}$. This shows that $[a, b]_{\sqsubseteq} \in L$ is closed in X . Therefore the order topology of \sqsubseteq is contained in the

⁹This implicitly implies that for all atoms $v, w \leq x$ there is a smallest connected $b \leq x$ with $v, w \leq b$.

original topology of X . Now property I3 for X precisely says that the converse is true as well. Hence the order topology of \sqsubseteq is the original topology of X . Since X has a countable basis it also has an infinite countable dense subset Y (X is infinite since the betweenness relation B is dense).

Altogether we see that the topology of X is given by the order topology of a dense and bounded order that has a countable dense subset and that is connected. By [Bing1960] this characterizes the homeomorphism type of the closed unit interval of \mathbb{R} . \square

4.5. *Example.* Without condition I3 of 4.2, the formula $I(x)$ would not define homeomorphic copies of $[0, 1]_{\mathbb{R}}$: For example let $S = \{\frac{1}{n} \mid n \in \mathbb{N}\}$ and let

$$\tau = \{O \setminus A \mid O \subseteq [0, 1]_{\mathbb{R}} \text{ open and } A \subseteq S\}.$$

Then τ is *Smirnov's Deleted Sequence Topology* on $[0, 1]_{\mathbb{R}}$ see [SteSee1995, no. 64, p. 86]. Let X be $[0, 1]_{\mathbb{R}}$ with this topology. Then X still has a countable basis, e.g. take all $O \setminus Y$, where O runs through a countable basis of $[0, 1]_{\mathbb{R}}$ in the order topology and $Y \subseteq S$ is cofinite. Routine checking shows that all intervals of $[0, 1]_{\mathbb{R}}$ are τ -connected and in the lattice of closed subsets of X , the element X satisfies I1 and I2. The order \sqsubseteq in the proof of 4.4 is the natural order of $[0, 1]_{\mathbb{R}}$ or its inverse, but clearly X is not homeomorphic to the unit interval.

4.6. **Proposition.** *Let R be an o -minimal expansion of a dense linear order and let $X \subseteq R^n$ be definable with parameters. Let L be a lattice of parametrically definable subsets of X that are closed in X and suppose L contains all such sets of dimension ≤ 1 . Then $I(x)$ defines in L the set of all subsets of X that are definably homeomorphic to an interval of the form $[a, b]$ of R , $a < b$.*

Proof. Using 4.3, it is clear that $I(x)$ is satisfied for all $A \in L$ that are definably homeomorphic to some $[a, b] \subseteq R$.

Conversely, by the cell decomposition theorem [vdDries1998, chapter 3, 2.11, p. 52] we see that every $A \in L$ at which $I(x)$ is true must be definably connected and of dimension 1 (if $\dim A \geq 2$, then there are points $x, y \in A$ such that there is no smallest L -connected $B \in L$ containing x, y). Then A is a finite union of definably homeomorphic copies of intervals of R and routine checking shows that A must be definably homeomorphic to some $[a, b] \subseteq R$. \square

5. LINEAR SPACES AND CONVEX SETS

In [Grzego1951, §5] the “algebra of convexity” is shown to be undecidable. A prototype of such an algebra is the Boolean algebra of subsets of \mathbb{R}^n , $n \geq 2$, together with the convex hull operator. Generalizations of this type of closure operators have shown to be undecidable, e.g. see [Davis2013] and [Dornhe1998]. However lattices of convex sets themselves are much less studied from a logic perspective. At the end of [Grzego1951, §5] the question is raised whether the lattice of closed convex subsets of \mathbb{R}^n , $n \geq 2$ is undecidable. We confirm this in a strong sense, see 5.2.

5.1. Reminder on incidence geometry.

For any field k , the lattice U of sub-vector spaces of k^3 interprets the field k after naming three parameters. In fact (U, \subseteq) and $(k, +, \cdot)$ are bi-interpretable with definable parameters (cf. [Hodges1993, Remark 5, section 5.3, p. 215]). This is done via incidence geometry. We recall it briefly. Fix a 2-dimensional subspace H of k^3 (hence a maximal element of $U \setminus \{k^3\}$). Let P be the set of all 1-dimensional


subspaces of k^3 (these are the minimal elements of $U \setminus \{(0)\}$) that are not contained in H and let L be the set of all 2-dimensional subspaces of k^3 , except H . Then the relation $p \subseteq l$ between $p \in P$ and $l \in L$ is isomorphic to the affine incidence geometry defined by k between points of k^2 and affine 1-dimensional subspaces of k^2 . The isomorphism between the incidence geometries is given as follows: We may assume that $H = k \times k \times \{0\}$. Then the map

$$\tau : k^2 \longrightarrow P; (a, b) \mapsto (a, b, 1) \cdot k \subseteq k^2$$


is a bijection and the induced map between affine 1-dimensional subspaces of k^2 and L that maps A to $\tau(A)$ is a bijection as well. This also shows that the partially ordered sets $(P \cup L, \subseteq)$ and (\mathbb{A}, \subseteq) are isomorphic, where \mathbb{A} is the set of all proper, nonempty affine subspaces of k^2 .

Having the incidence geometries identified we may define the field k in (\mathbb{A}, \subseteq) after naming two points $0 \neq 1$ of \mathbb{A} as parameters (in order to get the field in (U, \subseteq) we need a third parameter for the definition of (\mathbb{A}, \subseteq) in (U, \subseteq)): As a set it will be the line through $0, 1$ in the affine geometry, or a 2-dimensional subspace in the poset of sub-vector spaces of k^3 :

Addition of A and C on a line with fixed point 0 is defined as follows: Choose a point B not on the line and then use twice the fact that opposite lines in a parallelogram are of the same length (this is independent of B). Of course in an arbitrary field this has to be phrased appropriately, but over an ordered field it is exactly this. The pictures show how to construct $A + C$ out of $0, A$ and C step by step:



Multiplication of A and C on a line with fixed points 0 and 1 is done by applying the Intercept Theorem twice. Choose a point B not on the line. The pictures show how to construct $A \cdot C$ out of $0, 1, A, C$ step by step.



Hence the field is defined in (\mathbb{A}, \subseteq) once we have named two distinct points $0 \neq 1$.

The method of defining "intervals" is particularly easy for convex sets and identifies the strongest topological lattices: these are lattices of convex sets in higher dimensions. For example the lattice of (semi-linear) convex subsets of \mathbb{R}^n , $n \geq 2$, interprets $(\mathbb{R}, \mathbb{N}, +, \cdot)$, the expansion of the real field by the set of natural numbers. In detail:

5.2. Theorem. *Let K be an ordered field and let M be any expansion of the ordered K -vector space K . Let L be the poset of all parametrically definable and convex subsets of M^n and let $\bar{L} = \{C \in L \mid C \text{ is closed}\}$. Then the map $L \rightarrow \bar{L}$ that sends C to its closure is 0-definable in (L, \subseteq) ; in particular \bar{L} is 0-definable in (L, \subseteq) . The formulas defining these objects are independent of M and n .*

Further, in the poset (\bar{L}, \subseteq) the following sets are 0-definable:

- (i) *The set of all segments $[a, b]$ (= the convex hull of $\{a, b\}$ in M^n).*
- (ii) *The set of affine subspaces of M^n as well as the set of affine subspaces of M^n of a fixed dimension d .*
- (iii) *The set $\{C \in \bar{L} \mid C \text{ is bounded}\}$.*
- (iv) *If $n \geq 2$, then (\bar{L}, \subseteq) defines the ordered field K on any 1-dimensional subspace ℓ after naming two atoms of ℓ (ℓ itself is then definable over these atoms).*

Now assume $n \geq 2$ and every definable, bounded subset of M that is discrete in M , is finite.¹⁰ Then

- (v) *The family*

$$(\{\ell\} \times F \mid \ell \text{ is a line and } F \text{ is a finite set of atoms } \leq \ell)$$

is 0-definable in \bar{L} .

- (vi) *The expansion of the ordered field K by naming \mathbb{Q} as a subset, is interpretable in the poset (\bar{L}, \subseteq) .*

All formulas in (i)-(vi), defining the various objects can be chosen to be independent of M and n .

Proof. Firstly, observe that any $A \in L$ is uniquely determined by the atoms $\leq A$. We will talk about points of A rather than atoms $\leq A$ when we formulate elementary properties within the poset (L, \subseteq) .

Claim 1. The set of all segments $[a, b]$ is 0-definable in L by the formula $I^*(x)$, which says that there are atoms $A, B \leq x$ such that x is the supremum of $\{A, B\}$ in L . This is clear.

The formula $I^*(x)$ also defines the segments in $\bar{L} = (\bar{L}, \subseteq)$, which shows (i).

Claim 2. The map $L \rightarrow \bar{L}$ that sends C to its closure is 0-definable in L .

Proof. It is routine to check that indeed for every $C \in L$, the closure of C is in \bar{L} (just copy the proof for $M = \mathbb{R}$). Let $C \in L$. Then the convexity of C implies that $C \in \bar{L}$ if and only if for every proper (i.e., not a singleton) segment S and all $a \in S$ with $S \setminus \{a\} \subseteq C$ we have $a \in C$. This shows that \bar{L} is 0-definable in L . But then the map $L \rightarrow \bar{L}$ that sends C to its closure is also 0-definable in L : For $C \in L$ its closure \bar{C} is the smallest $D \in \bar{L}$ with $C \subseteq D$. \square

Claim 3. The set of all lines (i.e. 1-dimensional affine subspaces of M^n) is 0-definable in \bar{L} : A set $A \in \bar{L}$ is contained in a line if and only if for all $a, b, c \in A$

¹⁰This for example is true if M is o-minimal, or, if $K = \mathbb{R}$.

there is a segment containing a, b, c . Then a set $A \in \bar{L}$ is a line if and only if it is maximal among elements of \bar{L} that are contained in a line.

(ii). A set $A \in \bar{L}$ is an affine subspace if it is not empty and for all proper segments S and every line ℓ with $S \subseteq A, \ell$ we have $\ell \subseteq A$. A set $A \in \bar{L}$ is an affine subspace of dimension d if there is a non refinable chain of affine subspaces $A_0 \subsetneq \dots \subsetneq A_d = A$.

We will from now on assume that $n \geq 2$; item (iii) holds for $n = 1$ by (i).

(iv). By (ii) the plane, affine incidence geometry of K is definable in \bar{L} after naming a 2-dimensional subspace U . Hence $(K, +, \cdot)$ is definable in \bar{L} after naming U , a line $\ell \subseteq U$ and two points $0, 1 \in \ell$ (see 5.1). However, this field structure is independent of the chosen U , so we can wrap U in an existential quantifier (by (ii), the set of two-dimensional subspaces is 0-definable in \bar{L}). Furthermore, the order of the field is definable by using the betweenness relation on the segments of ℓ and declaring $0 < 1$. This shows (iv).

Claim 4. For each affine hyperplane A of M^n and each line $\ell \not\subseteq A$, let $\pi_{A,\ell}$ be the projection: $M^n \rightarrow \ell$ along A . If we consider $\pi_{A,\ell}$ as a map $\text{atoms}(\bar{L}) \rightarrow \{\text{atoms} \leq \ell\}$, then the family of (graphs of) all the $\pi_{A,\ell}$ is 0-definable in \bar{L} (by a formula that is independent of n and M).

Proof. The affine hyperplanes are precisely the maximal elements in the poset of proper affine subspaces. For $a \in M^n$ and $b \in \ell$ we then have $\pi_{A,\ell}(a) = b$ if and only if $a \in \ell$ and $b = a$, or, the unique affine hyperplane containing a and disjoint from A , intersects ℓ in b . This gives a 0-definable definition of the family of all the $\pi_{A,\ell}$. \square

(iii). A convex set C is bounded if and only if for every projection $\pi_{A,\ell}$ as in claim 4 there is a segment contained in ℓ that contains $\pi_{A,\ell}(C)$. (Of course it suffices to test with the n coordinate axes, but in our formulation the definition is independent of n .)

For the rest of the proof we now also assume that every definable bounded subset of M that is discrete in M , is finite.

Claim 5. The set of all $C \in \bar{L}$ that are convex hulls of finitely many points is 0-definable in \bar{L} .

Proof. If $F \subseteq K^n$ is finite, then the convex hull C of F in K^n is the convex hull of the extremal points of C ; this is proved as for \mathbb{R}^n . The extremal points of C are the points $a \in C$ with the property that for all segments $[c, b] \subseteq C$ with $a \in [c, b]$ we have $a = b$ or $a = c$. Further, the extremal points of C are contained in F . Hence we can express the property of $C \in \bar{L}$ being the convex hull of finitely many points as follows:

C is the smallest element in \bar{L} containing the set F of extremal points (or atoms) of C and any projection of F with a projection function $\pi_{A,\ell}$ as in claim 4 is bounded and discrete in ℓ .

Using segments we can express “ $\pi_{A,\ell}(F)$ is discrete in ℓ ” as a first order property of C, A, ℓ in (\bar{L}, \subseteq) . Notice that F here is seen a definable set of atoms (defined using the parameter $C \in \bar{L}$); F is not an element of L . Further, by the assumption made right before claim 5, we know that then F must be finite. \square

(v). Consider the following property of $A, \ell, C, D \in \bar{L}$:

(*) A is an affine hyperplane, $\ell \not\subseteq A$ is a line, C is the convex hull of finitely many points and there is an extremal point E of C with $D = \pi_{A,\ell}(E)$

Using claims 4 and 5, this can be expressed by an \leq -formula $\varphi(u, v, w, x)$. Consequently, given a line ℓ the sets defined by the formulas $\varphi(A, \ell, C, x)$, where A, C vary in \bar{L} is the set of finite subsets (of atoms) of ℓ .

(vi). By (iv) and (v), the weak monadic second order structure $W(K, +, \cdot)$ is interpretable in \bar{L} . By [Bauval1985] this implies that the polynomial ring in one variable over K is interpretable in \bar{L} . In this ring the rational numbers are definable (cf. [Robins1951, Section 2]). \square

6. LATTICES OF CLOSED DEFINABLE SETS IN THE ORDERED CONTEXT

We start with a combinatorial tool needed here and in 7.1.

6.1. Lemma. *Let K be an infinite field, $m \in \mathbb{N}$ and let S be a set of functions $K \rightarrow K$ containing all scalar multiplication functions, such that for all mutually distinct $\alpha_1, \dots, \alpha_m \in K$ and all $\beta_1, \dots, \beta_m \in K$, there is some $f \in S$ with $f(\alpha_i) = \beta_i$ ($1 \leq i \leq m$).*

Fix $n \geq 2$ and let V be the set of all functions $(t_1, \dots, t_n) : K^n \rightarrow K^n$ where the t_j are terms in x_1, \dots, x_n built from addition and the functions from S .

Take $a_1, b_1, \dots, a_m, b_m \in K^n$ with $a_i \neq a_j$ and $b_i \neq b_j$ for all $i \neq j \in \{1, \dots, m\}$. Then there is a bijective $F \in V$ whose compositional inverse is in V such that $F(a_i) = b_i$ for all $i \in \{1, \dots, m\}$.¹¹

Proof. Since $a_i \neq a_j$ for all $i \neq j$ and K is infinite, some projection on a one dimensional subspace of K^n must be injective on $\{a_1, \dots, a_m\}$ (choose a line that does not contain any $a_i - a_j$, $i \neq j$; then the projection to a linear complement of this line is injective on $\{a_1, \dots, a_m\}$ and then use induction). Hence there is a K -vector space isomorphism $l : K^n \rightarrow K^n$ such that the first coordinates of the points $l(a_1), \dots, l(a_m)$ are mutually distinct

Since V is closed under composition and V contains all linear maps $K^n \rightarrow K^n$, we may replace a_i by $l(a_i)$ and assume that $a_i = (\alpha_i, c_i) \in K \times K^{n-1}$ with $\alpha_i \neq \alpha_j$ for all $i \neq j$. Similarly we may assume that $b_i = (d_i, \beta_i) \in K^{n-1} \times K$ with $\beta_i \neq \beta_j$ for all $i \neq j$.

We will now find a bijective map $G \in V$ with $G^{-1} \in V$ such that

$$G(a_i) = (\alpha_i, \bar{0}, \beta_i) \quad (1 \leq i \leq m),$$

where $\bar{0} \in K^{n-2}$. Similarly there is then also some $H \in V$ with $H^{-1} \in V$ such that $H(b_i) = (\alpha_i, \bar{0}, \beta_i)$ and so with $F = H^{-1} \circ G \in V$ we have a function as required for the lemma.

For the definition of G we use the assumption on S , which implies that there are $f_2, \dots, f_n \in S$ with

$$(f_2, \dots, f_n)(\alpha_i) = c_i - (\bar{0}, \beta_i)$$

for all $i \in \{1, \dots, m\}$. Then the function $G : K^n \rightarrow K^n$ defined by

$$G(x_1, \dots, x_n) = (x_1, x_2 - f_2(x_1), \dots, x_n - f_n(x_1))$$

¹¹We will apply this for $S = K[X]$ and (when K is ordered) for the set of all piecewise linear continuous functions $K \rightarrow K$. In the first case, all functions in V will be polynomial maps and in the second case all functions in V will be piecewise linear continuous functions $K^n \rightarrow K^n$.

is in V and $G(a_i) = G(\alpha_i, c_i) = (\alpha_i, c_i - (f_2, \dots, f_n)(\alpha_i)) = (\alpha_i, \bar{0}, \beta_i)$. The compositional inverse of G maps (x_1, \dots, x_n) to $(x_1, x_2 + f_2(x_1), x_n + f_n(x_1))$. Hence $G^{-1} \in V$. \square

6.2. Corollary. *Let K be an ordered field and let $n \geq 2$. If $p_1, q_1, \dots, p_k, q_k \in K^n$ are $2k$ distinct points then there are piecewise linear homeomorphisms $\sigma_1, \dots, \sigma_k : [0, 1]_K \rightarrow K^n$ onto their respective images with mutually disjoint images such that $\sigma_i(0) = p_i$ and $\sigma_i(1) = q_i$ for all i .*

Proof. By 6.1 applied to the set S of continuous piecewise linear maps $K \rightarrow K$ there is a piecewise linear homeomorphism $F : K^n \rightarrow K^n$ such that $F(p_i) = (2i - 1, 0) \in K \times K^{n-1}$ and $F(q_i) = (2i, 0) \in K \times K^{n-1}$. For these points we can just take σ_i to be the segment between p_i and q_i . Applying F^{-1} then gives the corollary. \square

For the rest of this section we fix an ordered field K and work with an expansion M of the ordered K -vector space K . We will always assume that M is o-minimal, or, $K = \mathbb{R}$ and M is the full structure on \mathbb{R} , meaning that all subsets of all \mathbb{R}^n are definable.

6.3. Theorem. *Let $n \geq 2$. Let L be a lattice of closed and definable subsets of M^n and suppose that L contains all closed subsets of M^n that are contained in a finite union of definably homeomorphic copies of $[0, 1]_K$; when M is the total expansion of \mathbb{R} we also assume that condition (a) of 4.4 is satisfied.¹² Then in L the binary relation*

$$E(A, B) \iff A, B \text{ are finite and of the same size}$$

is 0-definable in L . Further, (L, \subseteq) interprets $(\mathbb{N}, +, \cdot)$ after naming an element $A \in L$ with $L \models I(A)$ (see 4.2). The interpretation is independent of M and n .¹³

Proof. Firstly, by 4.4 (if M is the full structure on \mathbb{R}) and 4.6 (if M is o-minimal) we can define the property "A is finite": We say that there is some $C \in L$ with $L \models I(C)$ such that A is a subset of C that is discrete in C (for the o-minimal case recall that every definable and discrete set is finite). By 2.9 it then suffices to show that for disjoint $A, B \in L$ we can define that they are of the same size. We claim that this is the case if and only if there is some $C \in L$ such that

- (a) each L -connected component of C hits A in exactly one atom and for each atom $a \leq A$ there is a unique L -connected component of C containing a , and,
- (b) each L -connected component of C hits B in exactly one atom and for each atom $b \leq B$ there is a unique L -connected component of C containing b .

The condition implies that the map from the L -connected components of C to the atoms of A , $D \mapsto D \cap A$ is a bijection, and similarly with B . Conversely, if A and B are disjoint and of size k , say, then choose $\sigma_1, \dots, \sigma_k$ according to 6.2. Then the union C of the images of the σ_i satisfies (a) and (b) since L contains all closed subsets of C and L is contained in the lattice of all closed sets.

¹²The largest such lattice is the lattice of all definable subsets and the smallest such lattice is the set of all closed subsets that are contained in a finite union of definably homeomorphic copies of $[0, 1]_K$.

¹³In [Grzego1951, §3] the undecidability of the lattice of all closed subsets of \mathbb{R}^n , $n \geq 2$ is proved; but it is neither shown that the lattice interprets $(\mathbb{N}, +, \cdot)$ nor that the property "finite" is definable in the lattice. See section 8 for a discussion.

This shows that E is definable. It follows that for each $A \in L$ with $L \models I(A)$, the structure $(W(A, B), E|_{W(A, B)})$ is definable in (L, \subseteq) , where B is the betweenness relation of A asserted to exist by the formula $I(x)$. Now 3.2 implies that $(\mathbb{N}, +, \cdot)$ is interpretable in L . \square

6.4. Corollary. *Let $n \geq 2$ and assume M is o-minimal with the convention of this section in force. Let P be the set of all closed, definable, definably connected and bounded subsets of M^n of dimension ≤ 1 , partially ordered by inclusion.¹⁴ Then the poset P interprets, independently of M , the lattice L of all closed and bounded subsets of M^n of dimension ≤ 1 . Hence by 6.3, (P, \subseteq) interprets $(\mathbb{N}, +, \cdot)$.*

Proof. (L, \subseteq) is interpretable in (P, \subseteq) because every set A in L is the intersection of two sets in P : By o-minimality, there is a definable continuous map $\sigma : [0, 1]_K \rightarrow M^n$ that has A in its image. A suitable perturbation of σ will then cut out the complement of A in the image of σ ; the easy details are left to the reader.

Now we can interpret (L, \subseteq) in (P, \subseteq) as $P \times P$ modulo the equivalence relation $(A, B) \sim (C, D) \iff$ the set of atoms below A and B is the set of atoms below C and D . Similarly inclusion of L can be interpreted. \square

6.5. Remark. When the dimension of the ambient space is 1 or if the ambient space has no definably connected sets, the lattices in 6.3 generally behave better:

- (i) The lattice of closed subsets of \mathbb{Q}^n is decidable and also the closure algebra of \mathbb{Q}^n is decidable¹⁵. This is because \mathbb{Q}^n is homeomorphic to \mathbb{Q} and for \mathbb{Q} it is deduced by Rabin from 3.3 in [Rabin1969].
- (ii) In [Rabin1969], Rabin also shows that the lattice of closed subsets of \mathbb{R} is decidable, answering a question of [Grzego1951]. In contrast, Shelah has proved in [Shelah1975] (under the continuum hypothesis) that the closure algebra of \mathbb{R} is undecidable.
- (iii) If X is a Boolean space, then the lattice of closed subsets is decidable, see 7.2 below.

6.6. It follows from 6.3 that for every ordered field K , the lattice of closed and semi-linear subsets of K^2 interprets $(\mathbb{N}, +, \cdot)$. We can also interpret $(\mathbb{N}, +, \cdot)$ in finite unions of rectangles of various sorts.¹⁶ We focus on one case, which also addresses the theme of 3.7: If T is an infinite totally ordered set, then consider the lattice L of finite unions of closed infinite rectangles of the form $(-\infty, p] \times (-\infty, q] \subseteq T \times T$ with $p, q \in T$. The partially ordered set of such rectangles is obviously itself a lattice, which is isomorphic to the poset $T \times T$ with componentwise partial order. Further, the elements of L are in bijection with the nonempty, finite anti-chains of $T \times T$ and the order of L translates to the partial order $A \leq B$ of finite anti-chains of $T \times T$ given by $\forall a \in A \exists b \in B : a \leq b$. All these data are 0-definable in $W(T \times T, \leq)$ and so L is 0-definable in $W(T \times T, \leq)$.

¹⁴The interest in P comes from the observation that P is the set of images of continuous definable maps $[0, 1]_K \rightarrow M^n$. In a forthcoming paper we will show that the lattice ordered abelian group of these function is decidable, in contrast to the assertion of the corollary.

¹⁵The closure algebra of \mathbb{Q}^n is the expansion of the boolean algebra of all subsets of \mathbb{Q}^n by the map that sends a set to its closure.

¹⁶In [Grzego1951, §5] the closure operation on the powerset of \mathbb{R}^2 with values in such lattices is dealt with, but not the lattices themselves.

6.7. Proposition. *Let T be an infinite totally ordered set and let \leq be the componentwise partial order on $T \times T$. Let L be the poset of finite anti-chains of $(T \times T, \leq)$ under the partial order described in 6.6 (hence L indeed is the lattice described there). Then L interprets $(\mathbb{N}, +, \cdot)$ after naming two definable parameters.¹⁷ The interpretation is independent of T .*

Proof. In a nutshell, this is true because L is in bijection with the set

$$\{(U, V) \in W(T) \times W(T) \mid U \text{ and } V \text{ have the same size}\};$$

the map sends $A \in L$ to $(p_1(A), p_2(A))$, where p_i are the projections onto the coordinate axes. Hence L “is” the equivalence E of the structure $(W(T), E)$ of 3.2. In detail:

Firstly, the poset $(T \times T, \leq)$ is 0-definable in (L, \leq) as the sub-poset of join irreducible elements. We now define certain data in $(T \times T, \leq)$.

For $x, y \in T^2 = T \times T$ we write $[x, y] = \{z \in T^2 \mid x \leq z \leq y\}$. Let $p, q \in T^2$. We say p, q define a line segment if $p \neq q$ and the binary relation $x \leq y$ restricted to $[p, q]$ is a total order with least element p and largest element q .

Hence the binary relation “ p, q define a line segment” is 0-definable in (T^2, \leq) . Obviously, p, q define a line segment if and only if $p \leq q$, $p \neq q$ and $(p_1 = q_1 \text{ or } p_2 = q_2)$. A line segment is a set of the form $[p, q]$, where p, q define a line segment.

Suppose $p, q \in T^2$ define a line segment. We define

$$\ell(p, q) = \{r \in T^2 \mid [p, r] \cup [r, p] \cup [p, q] \text{ is a line segment}\},$$

which is the line through p and q . Hence the ternary relation “ $r \in \ell(p, q)$ ” is 0-definable in (T^2, \leq) . Further, we define a map $\pi_{p,q} : T^2 \rightarrow \ell(p, q)$ by

$$\pi_{p,q}(r) = \begin{cases} r, & \text{if } r \in \ell(p, q), \\ \text{the unique } s \in \ell(p, q) \text{ s.th. } r, s \text{ define a} \\ \text{line segment, or, } s, r \text{ define a line segment,} & \text{if } r \notin \ell(p, q). \end{cases}$$

Hence $\pi_{p,q}$ is the projection onto $\ell(p, q)$.

Let $o, p, q \in T^2$. We say o, p, q define a coordinate system if o, p and o, q define line segments and $q \notin \ell(o, p)$, $p \notin \ell(o, q)$. Hence the ternary relation “ o, p, q define a coordinate system” is 0-definable in (T^2, \leq) .

Now fix $o, p, q \in T^2$ defining a coordinate system. We write $\ell = \ell(o, p)$ and show that the structure $(W(\ell, \leq), E)$ from 3.2 is interpretable in (L, \leq) by using the bijection from the beginning of the proof.

Firstly, we now read the data above in the poset L . Hence for example, $\pi_{o,p}$ is a map from the join irreducible elements of L onto the set of join-irreducible elements that correspond to points of $\ell_{o,p}$. Further, we read elements of L as the set of join irreducible elements contained in it. In this set up then, $W(\ell, \leq)$ is interpretable in L by using the identification of $A, B \in L$ when $\pi_{o,p}(A) = \pi_{o,p}(B)$ (which is definable in L). Observe that projections are injective on anti-chains.

It remains to interpret “equal size” of elements of $W(\ell, \leq)$: Let $A, B \in L$. Then $\pi_{o,p}(A)$ and $\pi_{o,p}(B)$ have the same size if and only if there are $G, H_A, H_B \in L$ such that

$$\bullet \pi_{o,p}(H_A) = \pi_{o,p}(A),$$

¹⁷This means that the set of parameters needed for the interpretation is 0-definable.

- $\pi_{o,p}(H_B) = \pi_{o,p}(B)$,
- $\pi_{o,q}(H_A) = \pi_{o,q}(G)$,
- $\pi_{o,q}(H_B) = \pi_{o,q}(G)$.

This is first order expressible in (L, \leq) , using o, p, q as parameters.

Hence indeed $(W(\ell, \leq), E)$ is interpretable in L and 3.2 entails that (L, \leq) interprets $(\mathbb{N}, +, \cdot)$ after naming the two definable parameters o, p ; the appearance of q above can be wrapped into an existential quantifier. \square

7. ZARISKI CLOSED SETS AND P-ADIC SETS

7.1. Proposition. *Let $n \geq 2$, and let K be an infinite field. Let L be the lattice of Zariski closed subsets of K^n , let L_1 be the lattice of Zariski closed sets of Krull dimension ≤ 1 and let L_2 be the lattice generated by all absolutely irreducible Zariski closed sets of Krull dimension ≤ 1 .¹⁸ Then L defines L_1 and L_1 as well as L_2 interpret $(\mathbb{N}, +, \cdot)$. The definition and both interpretations are parameter free and independent of K and n .¹⁹*

Proof. By definition of the Krull dimension of Zariski closed subsets of K^n , a set $Z \in L$ is in L_1 if and only if there are no irreducible sets (i.e., join irreducible elements of L) $\emptyset \neq A \subsetneq B \subsetneq C \subseteq Z$. Hence L_1 is definable in the lattice L .

The interpretation of $(\mathbb{N}, +, \cdot)$ in L_1 and in L_2 is the same (just evaluated on different structures) and we write L_* to mean either of them.

Notice that the set of finite subsets of K^n is 0-definable in L_* as those sets whose irreducible components are atoms. We will show that the structure N induced by L_* on the set $W(K^n)$ defines “same size” as required for 2.9(i) and that N has a definable relation as in 2.9(ii). Then 2.9 will give us an interpretation as required.

(a). Let $p_1, \dots, p_k, q_1, \dots, q_k \in K^n$ be $2k$ points. Then there is a Zariski closed subset $V \subseteq K^n$ of dimension 1 such that

- each irreducible component of V is absolutely irreducible,
- for each irreducible component C of V there is some (necessarily unique) $i \in \{1, \dots, k\}$ with $C \cap \{p_1, \dots, p_k, q_1, \dots, q_k\} = \{p_i, q_i\}$, and,
- the resulting map from the set of irreducible components of V to $\{1, \dots, k\}$ is a bijection.

This property implies that the structure $(W(K^n), E)$ from 2.9(i) is 0-definable in (L_*, \subseteq) .

Proof. Firstly, if $E, F \subseteq K^n$ are finite and disjoint, then there is an absolutely irreducible curve $C \subseteq K^n$ containing E and disjoint from F . To see this we apply 6.1 with S being the polynomials over K in one variable. By 6.1 then, there is a polynomial isomorphism $f : K^n \rightarrow K^n$ (so all entries of f and f^{-1} are polynomials over K in n variables), such that $f(E \cup F) \subseteq K \times \{0\}^{n-1}$. So we may assume $E \cup F \subseteq K \times \{0\}^{n-1}$ all along. Write $E = \{(a_1, 0), \dots, (a_m, 0)\}$ with $a_i \neq a_j \in K$ for $i \neq j$. Then the curve defined by $X_2 = (X_1 - a_1) \cdot \dots \cdot (X_1 - a_m), X_3 = 0, \dots, X_n = 0$ has the required property.

¹⁸We are dealing here with the naive Zariski topology on K^n , which has zero sets in K^n of polynomials $\in K[X_1, \dots, X_n]$ as a basis of closed sets. An absolutely irreducible Zariski closed set then is a set whose Zariski closure in \overline{K}^n is still irreducible, \overline{K} = algebraic closure of K .

¹⁹The undecidability of the lattice L is also established in [Grzego1951, §3]. See section 8 for a discussion.

Using this fact for each $i \in \{1, \dots, k\}$ and writing $P = \{p_1, \dots, p_k\}$ and $Q = \{q_1, \dots, q_k\}$ we get an absolutely irreducible curve $C_i \subseteq K^n$ with $C_i \cap (P \cup Q) = \{p_i, q_i\}$. Then $V = C_1 \cup \dots \cup C_k$ has the required property. \square

(b) Now we define a family as required in 2.9(ii). Define a ternary relation S on L_* by

$$S(a, b, c) \iff a, c \text{ are finite and there is an irreducible component } x \text{ of } b \text{ with } a = c \cap x.$$

It is clear that S is 0-definable in L_* . For all $b, c \in L_*$, the set of cardinalities of $a \in L_*$ with $S(a, b, c)$ is finite: c has to be finite and the size of a is the size of $c \cap x$ for some irreducible component x of b . Hence 2.9(ii)(a) is satisfied for S .

On the other hand, if $T = \{n_1, \dots, n_k\} \subseteq \mathbb{N}$ is finite of size k then pick a set $b \in L_*$ with exactly k irreducible components, each infinite and choose a finite set a_i of size n_i in the i^{th} -irreducible component, disjoint from the other components. Then for $c = a_1 \cup \dots \cup a_k$, the set of cardinalities of $a \in L_*$ with $S(a, b, c)$ is T . \square

7.2. p -adic sets. In contrast to the ordered case (cf. 6.3), lattices of closed sets coming from a valuation topology of a valued field behave much better. This again follows from a result of Rabin in [Rabin1969]: He shows that any Boolean algebra with a second order quantifier over ideals is decidable. In topological terms, this means that the lattice of closed subsets of any Boolean space is decidable. Hence, for example, the lattice of closed subsets of \mathbb{Z}_p^n is decidable as this space is Boolean.

8. GRZEGORCZYK'S PAPER

We give a brief outline of [Grzego1951] and talk about the parts that have not been addressed in our note. This hopefully helps the reader studying the beautiful results in [Grzego1951] (which are a bit dipped in logic formalism).

The main theme of Grzegorzczuk's paper is the creation of a topological version of Robinson Arithmetic (see [Grzego1951, §1] and [TaMoRo1953, II.3, page 51]) and then to find models of that. This means, he states an axiom system for partially ordered sets (mainly intended to be applied to topological lattices) such that any lattice that is consistent with these axioms is undecidable. In fact, ordinary Robinson Arithmetic is interpreted in the topological set up and then Tarski's theorem on interpretations is invoked. As with ordinary Robinson Arithmetic, models of the topological axioms do not need to interpret Peano arithmetic, or the standard model $(\mathbb{N}_0, +, \cdot)$.

The models of this topological Robinson Arithmetic in [Grzego1951, §2, §4 and §5] are then "closure systems". This means that Boolean algebras are furnished with an operator that behaves in one sense or another like taking closures of arbitrary subsets of a topological space. The set up also encompasses closure operators like $S \mapsto \text{convex hull of } S \subseteq \mathbb{R}^n$, see [Grzego1951, §5]. (For a comparison: In our paper the weaker structures of the images of such operators are studied; recall from 6.5(ii) that the two points of view can differ quite substantially.) The set up in [Grzego1951, §2] is stated as a formal axiom system A1-A6, but this system is infinitary (see A4) and not first order. So the class of closure algebras addressed in §2 is not first order. A more down to earth way of reading the results in §2 is: Every closure algebra satisfying A1-A6 is a model of the topological Robinson arithmetic from §1.

In [Grzego1951, §3], topological lattices are studied in the sense of our paper. More precisely they are studied as a particular class of so-called *Brouwerian algebras*, also known as *(co-)Heyting algebras*. However they are still implicitly defined in terms of the infinitary language from §2 and again one might want to read the theorems in that section as “every lattice satisfying the properties stated at the beginning of §3 satisfies the topological Robinson arithmetic from §1”. This should be compared with our results in section 6. For example 6.3 is stronger for the lattices addressed there, because we can also define finite elements in the lattice and interpret $(\mathbb{N}, +, \cdot)$; on the other hand 6.3 is weaker because there are lattices outside the ordered context where [Grzego1951, §3] applies. For example Zariski closed sets, see 7.1.

In [Grzego1951, §4] the topic is “the algebra of bodies”. Boolean algebras with a binary predicate “ A, B are tangent” are studied. The theme here is closely related to what is called mereotopology (see [AiPrvB2007, Definition 2.5, p. 18]): One should think of the Boolean algebra as the algebra of regular *open* subsets of a topological space and then “being tangent” could be read as “the closures intersect”, or as “contact” in mereotopological terminology. Our paper is not addressing this interesting subject, instead we refer to the handbook of spatial logics [AiPrvB2007] (for example see p. 69 there).

We conclude with two questions.

- (1) Let P be the set of irreducible Zariski closed subsets of \mathbb{C}^2 . Is the partially ordered set (P, \subseteq) decidable?
Does (P, \subseteq) interpret the poset of affine subspaces of \mathbb{C}^2 (and therefore also $(\mathbb{C}, +, \cdot)$ by 5.1)? Notice that the affine subspaces are not 0-definable, even in the lattice of Zariski closed subsets of \mathbb{C}^2 since \mathbb{C}^2 is k -transitive for all k under the group of polynomial isomorphisms (use 6.1).
- (2) Let L be the lattice of closed subsets of \mathbb{R}^2 . Does L interpret the real field $(\mathbb{R}, +, \cdot)$? Note that by 6.3 together with standard coding tricks, we can interpret the field of real algebraic numbers in L . There is evidence that in the lattice of closed and semi-algebraic subsets of \mathbb{R}^2 the real field is not parametrically *definable*; this will be explained in a forthcoming paper.

Acknowledgement. I wish to thank the anonymous referee for carefully proof-reading the manuscript and for many valuable comments to improve the text.

REFERENCES

- [AiPrvB2007] Marco Aiello, Ian Pratt-Hartmann, and Johan van Benthem, editors. Handbook of spatial logics. Springer, Dordrecht, 2007. 22
- [Bauval1985] Anne Bauval. Polynomial rings and weak second-order logic. J. Symbolic Logic, 50(4):953–972 (1986), 1985. 4, 16
- [Bing1960] R. H. Bing. A Topological Characterization of a Line. Amer. Math. Monthly, 67(7):38–41, 1960. 12
- [Davis2013] Ernest Davis. The expressive power of first-order topological languages. J. Logic Comput., 23(5):1107–1141, 2013. 12
- [Dornhe1998] Christoph Dornheim. Undecidability of plane polygonal mereotopology. In Principles of Knowledge Representation and Reasoning: Proceedings of the 6th International Conference (KR-98, pages 342–353. Morgan Kaufman, 1998. 12
- [Grzego1951] Andrzej Grzegorzcyk. Undecidability of some topological theories. Fund. Math., 38:137–152, 1951. 2, 12, 17, 18, 20, 21, 22

- [Hodges1993] Wilfrid Hodges. Model theory, volume 42 of Encyclopedia of Mathematics and its Applications. Cambridge University Press, Cambridge, 1993. 3, 6, 7, 12
- [Laeuch1968] H. Läuchli. A decision procedure for the weak second order theory of linear order. In Contributions to Math. Logic (Colloquium, Hannover, 1966), pages 189–197. North-Holland, Amsterdam, 1968. 10
- [Rabin1969] Michael O. Rabin. Decidability of second-order theories and automata on infinite trees. Trans. Amer. Math. Soc., 141:1–35, 1969. 2, 9, 18, 21
- [Robins1951] Raphael M. Robinson. Undecidable rings. Trans. Amer. Math. Soc., 70:137–159, 1951. 16
- [Shelah1975] Saharon Shelah. The monadic theory of order. Ann. of Math. (2), 102(3):379–419, 1975. 18
- [SteSee1995] Lynn Arthur Steen and J. Arthur Seebach, Jr. Counterexamples in topology. Dover Publications, Inc., Mineola, NY, 1995. Reprint of the second (1978) edition. 12
- [TaMoRo1953] Alfred Tarski. Undecidable theories. Studies in Logic and the Foundations of Mathematics. North-Holland Publishing Company, Amsterdam, 1953. In collaboration with Andrzej Mostowski and Raphael M. Robinson. 21
- [Toffal1997] Carlo Toffalori. An undecidability theorem for lattices over group rings. Ann. Pure Appl. Logic, 88(2-3):241–262, 1997. Joint AILA-KGS Model Theory Meeting (Florence, 1995). 5
- [vdDries1998] Lou van den Dries. Tame topology and o-minimal structures, volume 248 of London Mathematical Society Lecture Note Series. Cambridge University Press, Cambridge, 1998. 3, 12

THE UNIVERSITY OF MANCHESTER, SCHOOL OF MATHEMATICS, OXFORD ROAD, MANCHESTER M13 9PL, UK

HOME PAGE: [HTTP://PERSONALPAGES.MANCHESTER.AC.UK/STAFF/MARCUS.TRESSL/INDEX.PHP](http://PERSONALPAGES.MANCHESTER.AC.UK/STAFF/MARCUS.TRESSL/INDEX.PHP)

Email address: marcus.tressl@manchester.ac.uk