

Censored α -stable processes

Andreas Kyprianou

Juan-Carlos Pardo*

Alex Watson

*CIMAT, Mexico

Prob-Lab seminar, Bath, 5/12/2011

Outline

- 1 Problem
- 2 Tools
 - pssMps and the Lamperti transform
 - α -stable processes
 - Lamperti-stable processes
- 3 The censored process
 - Construction
 - Lamperti transform
 - Wiener-Hopf factorisation
- 4 Results

Problem statement

Definition: α -stable

A Lévy process $(X, P_x)_{x \in \mathbb{R}}$ is called **α -stable** if it satisfies the **scaling property**

$$(cX_{c^{-\alpha}t})_{t \geq 0} \Big|_{P_x} \stackrel{d}{=} X \Big|_{P_{cx}}, \quad c > 0$$

The problem

Let

$$\tau_{-1}^1 = \inf\{t > 0 : X_t \in (-1, 1)\}$$

be the first hitting time of $(-1, 1)$.

What is $P_x(X_{\tau_{-1}^1} \in dy, \tau_{-1}^1 < \infty)$?

Problem statement

Definition: α -stable

A Lévy process $(X, P_x)_{x \in \mathbb{R}}$ is called α -stable if it satisfies the scaling property

$$(cX_{c^{-\alpha}t})_{t \geq 0} \Big|_{P_x} \stackrel{d}{=} X \Big|_{P_{cx}}, \quad c > 0$$

The problem

Let

$$\tau_{-1}^1 = \inf\{t > 0 : X_t \in (-1, 1)\}$$

be the first hitting time of $(-1, 1)$.

What is $P_x(X_{\tau_{-1}^1} \in dy, \tau_{-1}^1 < \infty)$?

Problem: history

- Blumenthal, Gettoor, Ray (1961): symmetric, d -dimensional
- Port (1967): one-sided jumps

Problem: history

- Blumenthal, Gettoor, Ray (1961): symmetric, d -dimensional
- Port (1967): one-sided jumps

Theorem (B-G-R)

Let $x > 1$. Then, when $\alpha \in (0, 1]$,

$$\begin{aligned} P_x(X_{\tau_{-1}^1} \in dy, \tau_{-1}^1 < \infty) / dy \\ = \frac{\sin(\pi\alpha/2)}{\pi} (x^2 - 1)^{\alpha/2} (1 - y^2)^{-\alpha/2} (x - y)^{-1} \end{aligned}$$

for $y \in (-1, 1)$.

Problem: history

- Blumenthal, Gettoor, Ray (1961): symmetric, d -dimensional
- Port (1967): one-sided jumps

Theorem (B-G-R)

Let $x > 1$. Then, when $\alpha \in (1, 2)$,

$$\begin{aligned}
 & P_x(X_{\tau_{-1}^1} \in dy)/dy \\
 &= \frac{\sin(\pi\alpha/2)}{\pi} (x^2 - 1)^{\alpha/2} (1 - y^2)^{-\alpha/2} (x - y)^{-1} \\
 &\quad - (\alpha - 1) \frac{\sin(\pi\alpha/2)}{\pi} (1 - y^2)^{-\alpha/2} \int_1^x (t^2 - 1)^{\alpha/2 - 1} dt,
 \end{aligned}$$

for $y \in (-1, 1)$.

Outline

- 1 Problem
- 2 Tools
 - pssMps and the Lamperti transform
 - α -stable processes
 - Lamperti-stable processes
- 3 The censored process
 - Construction
 - Lamperti transform
 - Wiener-Hopf factorisation
- 4 Results

Positive, self-similar Markov processes

α -pssMp

$[0, \infty)$ -valued Markov process,
equipped with initial measures P_x , $x > 0$,
with 0 an absorbing state,
satisfying the scaling property

$$(cX_{c^{-\alpha}t})_{t \geq 0} \Big|_{P_x} \stackrel{d}{=} X \Big|_{P_{cx}}, \quad x, c > 0$$

Positive, self-similar Markov processes

α -pssMp

$[0, \infty)$ -valued Markov process,
equipped with initial measures $P_x, x > 0$,
with 0 an absorbing state,
satisfying the scaling property

$$(cX_{c^{-\alpha}t})_{t \geq 0} \Big|_{P_x} \stackrel{d}{=} X \Big|_{P_{cx}}, \quad x, c > 0$$

Positive, self-similar Markov processes

α -pssMp

$[0, \infty)$ -valued Markov process,
 equipped with initial measures P_x , $x > 0$,
 with 0 an **absorbing state**,
 satisfying the scaling property

$$(cX_{c^{-\alpha}t})_{t \geq 0} \Big|_{P_x} \stackrel{d}{=} X \Big|_{P_{cx}}, \quad x, c > 0$$

Positive, self-similar Markov processes

α -pssMp

$[0, \infty)$ -valued Markov process,
equipped with initial measures P_x , $x > 0$,
with 0 an absorbing state,
satisfying the **scaling property**

$$(cX_{c^{-\alpha}t})_{t \geq 0} \Big|_{P_x} \stackrel{d}{=} X \Big|_{P_{cx}}, \quad x, c > 0$$

Lamperti transform

$(X, P_x)_{x>0}$ α -pssMp

\leftrightarrow

$(\xi, \mathbb{P}_y)_{y \in \mathbb{R}}$ killed Lévy

$$X_t = \exp(\xi_{S(t)}),$$

$$\xi_s = \log(X_{T(s)}),$$

S a random time-change

T a random time-change

Lamperti transform

$(X, P_x)_{x>0}$ α -pssMp

$$X_t = \exp(\xi_{S(t)}),$$

S a random time-change

\leftrightarrow

$(\xi, \mathbb{P}_y)_{y \in \mathbb{R}}$ killed Lévy

$$\xi_s = \log(X_{T(s)}),$$

T a random time-change

$\left. \begin{array}{l} X \text{ never hits zero} \\ X \text{ hits zero continuously} \\ X \text{ hits zero by a jump} \end{array} \right\} \leftrightarrow$

\leftrightarrow

$\left\{ \begin{array}{l} \xi \rightarrow \infty \text{ or } \xi \text{ oscillates} \\ \xi \rightarrow -\infty \\ \xi \text{ is killed} \end{array} \right.$

Stable processes

Definition 1

A Lévy process X is called α -stable if it satisfies the scaling property

$$(cX_{c^{-\alpha}t})_{t \geq 0} \Big|_{P_x} \stackrel{d}{=} X \Big|_{P_{cx}}, \quad c > 0.$$

Necessarily $\alpha \in (0, 2]$. [$\alpha = 2 \rightarrow$ BM, exclude this.]

The quantity $\rho = P_0(X_t \geq 0)$ is well defined.

Stable processes

Definition I

A Lévy process X is called α -stable if it satisfies the scaling property

$$(cX_{c^{-\alpha}t})_{t \geq 0} \Big|_{P_x} \stackrel{d}{=} X \Big|_{P_{cx}}, \quad c > 0.$$

Necessarily $\alpha \in (0, 2]$. [$\alpha = 2 \rightarrow$ BM, exclude this.]

The quantity $\rho = P_0(X_t \geq 0)$ is well defined.

Definition II

Let α, ρ be admissible parameters, X the Lévy process with Lévy density

$$c_+ x^{-(\alpha+1)} \mathbb{1}_{(x>0)} + c_- |x|^{-(\alpha+1)} \mathbb{1}_{(x<0)}, \quad x \in \mathbb{R},$$

no Gaussian part and a certain centre. Setting $\hat{\rho} := 1 - \rho$, we can put

$$c_+ = \frac{\Gamma(\alpha + 1)}{\Gamma(\alpha\rho)\Gamma(1 - \alpha\rho)}, \quad c_- = \frac{\Gamma(\alpha + 1)}{\Gamma(\alpha\hat{\rho})\Gamma(1 - \alpha\hat{\rho})}.$$

Stable processes

We make two assumptions:

- X does not have one-sided jumps,
- When $\alpha = 1$, X is symmetric.

A Lamperti-stable process

Few examples where the Lamperti transform is known.

A Lamperti-stable process

Few examples where the Lamperti transform is known.

Let X be a stable process, and define

$$X_t^* = X_t \mathbb{1}_{(t < \tau_0^-)}, \quad t \geq 0,$$

where

$$\tau_0^- = \inf\{t > 0 : X_t < 0\}.$$

A Lamperti-stable process

Few examples where the Lamperti transform is known.

Let X be a stable process, and define

$$X_t^* = X_t \mathbb{1}_{(t < \tau_0^-)}, \quad t \geq 0,$$

where

$$\tau_0^- = \inf\{t > 0 : X_t < 0\}.$$

Then X^* is a pssMp, with Lamperti transform ξ^* .

A Lamperti-stable process

Few examples where the Lamperti transform is known.
Let X be a stable process, and define

$$X_t^* = X_t \mathbb{1}_{(t < \tau_0^-)}, \quad t \geq 0,$$

where

$$\tau_0^- = \inf\{t > 0 : X_t < 0\}.$$

Then X^* is a pssMp, with Lamperti transform ξ^* .
 ξ^* has Lévy density

$$c_+ \frac{e^x}{(e^x - 1)^{\alpha+1}} \mathbb{1}_{(x>0)} + c_- \frac{e^x}{(1 - e^x)^{\alpha+1}} \mathbb{1}_{(x<0)},$$

and is killed at rate $c_-/\alpha = \frac{\Gamma(\alpha)}{\Gamma(\alpha\hat{\rho})\Gamma(1-\alpha\hat{\rho})}$.

Lamperti-stable subordinators

A **Lamperti-stable subordinator** with parameters (α, β) has Lévy measure

$$c \frac{e^{\beta x}}{(e^x - 1)^{\alpha+1}}, \quad x > 0.$$

Outline

- 1 Problem
- 2 Tools
 - pssMps and the Lamperti transform
 - α -stable processes
 - Lamperti-stable processes
- 3 The censored process
 - Construction
 - Lamperti transform
 - Wiener-Hopf factorisation
- 4 Results

Construction

- Start with X , the stable process.

Construction

- Start with X , the stable process.
- Let $A_t = \int_0^t \mathbb{1}_{(X_t > 0)} dt$.

Construction

- Start with X , the stable process.
- Let $A_t = \int_0^t \mathbb{1}_{(X_t > 0)} dt$.
- Let γ be the right-inverse of A , and put $\check{Y}_t := X_{\gamma(t)}$.

Construction

- Start with X , the stable process.
- Let $A_t = \int_0^t \mathbb{1}_{(X_t > 0)} dt$.
- Let γ be the right-inverse of A , and put $\check{Y}_t := X_{\gamma(t)}$.
- Finally, make zero an absorbing state: $Y_t = \check{Y}_t \mathbb{1}_{(t < T_0)}$.
This is the **censored stable process**.

The Lamperti transform and its structure

Censoring **preserves self-similarity**: Y is a pssMp.

The Lamperti transform and its structure

Censoring preserves self-similarity: Y is a pssMp.
Let ξ be the Lamperti transform of Y .

The Lamperti transform and its structure

Censoring preserves self-similarity: Y is a pssMp.
Let ξ be the Lamperti transform of Y .

Theorem

$\xi \stackrel{d}{=} \xi^L \oplus \xi^C$, with

- ξ^L equal in law to ξ^* with the killing removed,
- ξ^C a compound Poisson process with jump rate c_-/α .

The Lamperti transform and its structure

Censoring preserves self-similarity: Y is a pssMp.
Let ξ be the Lamperti transform of Y .

Theorem

$\xi \stackrel{d}{=} \xi^L \oplus \xi^C$, with

- ξ^L equal in law to ξ^* with the killing removed,
- ξ^C a compound Poisson process with jump rate c_-/α .

Proof.

By diagram.

The Lamperti transform and its structure

Censoring preserves self-similarity: Y is a pssMp.
Let ξ be the Lamperti transform of Y .

Theorem

$\xi \stackrel{d}{=} \xi^L \oplus \xi^C$, with

- ξ^L equal in law to ξ^* with the killing removed,
- ξ^C a compound Poisson process with jump rate c_-/α .

Proof.

By diagram.

Tricky element – show Δ independent of ξ^L .

Lamperti: $\Delta \leftrightarrow \frac{X_\sigma}{X_{\tau-}}$. By Markov property, reduces to showing

$P_x\left(\frac{X_\sigma}{X_{\tau-}} \in \cdot\right)$ does not depend on x .

Scaling property gives $X_\sigma|_{P_x} \stackrel{d}{=} xX_\sigma|_{P_1}$, etc. □

Wiener-Hopf factorisation

Recall: Wiener-Hopf factorisation

Let ξ be a Lévy process, $\mathbb{E}[e^{i\theta\xi_1}] = e^{-\Psi(\theta)}$.

Then there exist $\kappa, \hat{\kappa}$, such that:

$$\Psi(\theta) = \kappa(-i\theta)\hat{\kappa}(i\theta),$$

κ and $\hat{\kappa}$ Laplace exponents of increasing, possibly killed Lévy processes (subordinators) H and \hat{H} :

$$\mathbb{E}[e^{-\lambda H_1}] = e^{-\kappa(\lambda)}, \quad \mathbb{E}[e^{-\lambda \hat{H}_1}] = e^{-\hat{\kappa}(\lambda)}, \quad \lambda \geq 0.$$

Wiener-Hopf factorisation

Recall: Wiener-Hopf factorisation

Let ξ be a Lévy process, $\mathbb{E}[e^{i\theta\xi_1}] = e^{-\Psi(\theta)}$.

Then there exist $\kappa, \hat{\kappa}$, such that:

$$\Psi(\theta) = \kappa(-i\theta)\hat{\kappa}(i\theta),$$

κ and $\hat{\kappa}$ Laplace exponents of increasing, possibly killed Lévy processes (subordinators) H and \hat{H} :

$$\mathbb{E}[e^{-\lambda H_1}] = e^{-\kappa(\lambda)}, \quad \mathbb{E}[e^{-\lambda \hat{H}_1}] = e^{-\hat{\kappa}(\lambda)}, \quad \lambda \geq 0.$$

- unique

Wiener-Hopf factorisation

Recall: Wiener-Hopf factorisation

Let ξ be a Lévy process, $\mathbb{E}[e^{i\theta\xi_1}] = e^{-\Psi(\theta)}$.

Then there exist $\kappa, \hat{\kappa}$, such that:

$$\Psi(\theta) = \kappa(-i\theta)\hat{\kappa}(i\theta),$$

κ and $\hat{\kappa}$ Laplace exponents of increasing, possibly killed Lévy processes (subordinators) H and \hat{H} :

$$\mathbb{E}[e^{-\lambda H_1}] = e^{-\kappa(\lambda)}, \quad \mathbb{E}[e^{-\lambda \hat{H}_1}] = e^{-\hat{\kappa}(\lambda)}, \quad \lambda \geq 0.$$

- unique
- H and \hat{H} related to maxima and minima of ξ :
ascending and descending ladder processes.

Wiener-Hopf factorisation for ξ : $\alpha \in (0, 1]$

WHF for $\alpha \in (0, 1]$

$$\kappa(\lambda) = \frac{\Gamma(\alpha\rho + \lambda)}{\Gamma(\lambda)}, \quad \hat{\kappa}(\lambda) = \frac{\Gamma(1 - \alpha\rho + \lambda)}{\Gamma(1 - \alpha + \lambda)}, \quad \lambda \geq 0.$$

H : Lamperti-stable subordinator with parameters $(\alpha\rho, 1)$,

\hat{H} : Lamperti-stable subordinator with parameters $(\alpha\hat{\rho}, \alpha)$.

ξ a hypergeometric Lévy process
with parameters $(1, \alpha\rho, 1 - \alpha, \alpha\hat{\rho})$.

Lamperti-stable subordinators are nice! We can calculate:

- The Lévy measure of ξ ,
- The Lévy measures of H and \hat{H} ,
- The renewal measures, $\mathbb{E} \int_0^\infty \mathbb{1}_{(H_t \in \cdot)} dt$ and $\mathbb{E} \int_0^\infty \mathbb{1}_{(\hat{H}_t \in \cdot)} dt$.

Wiener-Hopf factorisation for ξ : $\alpha \in (1, 2)$ WHF for $\alpha \in (1, 2)$

$$\kappa(\lambda) = (\alpha - 1 + \lambda) \frac{\Gamma(\alpha\rho + \lambda)}{\Gamma(1 + \lambda)}, \quad \hat{\kappa}(\lambda) = \lambda \frac{\Gamma(1 - \alpha\rho + \lambda)}{\Gamma(2 - \alpha + \lambda)},$$

for $\lambda \geq 0$.

- $\kappa(\lambda) = \frac{\lambda}{\mathcal{T}_{\alpha-1}\psi(\lambda)}$, with ψ LSS($1 - \alpha\rho, \alpha\hat{\rho}$).
- $\hat{\kappa}(\lambda) = \frac{\lambda}{\phi(\lambda)}$, with ϕ LSS($1 - \alpha\hat{\rho}, \alpha\rho$).

Not as nice, but we can still calculate Lévy measures and renewal measures.

Outline

- 1 Problem
- 2 Tools
 - pssMps and the Lamperti transform
 - α -stable processes
 - Lamperti-stable processes
- 3 The censored process
 - Construction
 - Lamperti transform
 - Wiener-Hopf factorisation
- 4 Results

Results

Recall: the problem

Let X be a stable process and $x > 1$.

$$P_x(X_{\tau_{-1}^1} \in dy, \tau_{-1}^1 < \infty) = \text{what?}$$

Results

Recall: the problem

Let X be a stable process and $x > 1$.

$$P_x(X_{\tau_{-1}^1} \in dy, \tau_{-1}^1 < \infty) = \text{what?}$$

First shift up by 1:

$$P_{x+1}(X_{\tau_0^2} \in dy + 1, \tau_0^2 < \infty),$$

Results

Recall: the problem

Let X be a stable process and $x > 1$.

$$P_x(X_{\tau_{-1}^1} \in dy, \tau_{-1}^1 < \infty) = \text{what?}$$

First shift up by 1:

$$P_{x+1}(X_{\tau_0^2} \in dy + 1, \tau_0^2 < \infty),$$

then scale out x :

$$P_1\left(X_{\tau_0^{2/(x+1)}} \in \frac{dy+1}{x+1}, \tau_0^{2/(x+1)} < \infty\right)$$

Results

Recall: the problem

Let X be a stable process and $x > 1$.

$$P_x(X_{\tau_{-1}^1} \in dy, \tau_{-1}^1 < \infty) = \text{what?}$$

First shift up by 1:

$$P_{x+1}(X_{\tau_0^2} \in dy + 1, \tau_0^2 < \infty),$$

then scale out x :

$$P_1\left(X_{\tau_0^{2/(x+1)}} \in \frac{dy+1}{x+1}, \tau_0^{2/(x+1)} < \infty\right)$$

So, we just need to look at

$$P_1(X_{\tau_0^b} \in dz, \tau_0^b < \infty), \quad 0 < b < 1.$$

Results

We are looking for

$$P_1(X_{\tau_0^b} \in dz, \tau_0^b < \infty) = P_1(Y_{\tau_0^b} \in dz, \tau_0^b < \infty).$$

Results

We are looking for

$$P_1(X_{\tau_0^b} \in dz, \tau_0^b < \infty) = P_1(Y_{\tau_0^b} \in dz, \tau_0^b < \infty).$$

Recall: Lamperti transform

$$Y_t = \exp(\xi_{S(t)}), \quad \text{and} \quad \xi_s = \log Y_{T(s)},$$

where S, T are random, mutually inverse time-changes.

Key fact: if $a = \log b$, then $(0, b)$ corresponds to $(-\infty, a)$ and τ_0^b corresponds to $S_a^- = \inf\{s > 0 : \xi_s < a\}$. Then,

$$Y_{\tau_0^b} = \exp(\xi_{S_a^-}).$$

Results

We are looking for

$$\mathbb{P}_1(X_{\tau_0^b} \in dz, \tau_0^b < \infty) = \mathbb{P}_1(Y_{\tau_0^b} \in dz, \tau_0^b < \infty).$$

Recall: Lamperti transform

$$Y_t = \exp(\xi_{S(t)}), \quad \text{and} \quad \xi_s = \log Y_{T(s)},$$

where S, T are random, mutually inverse time-changes.

Key fact: if $a = \log b$, then $(0, b)$ corresponds to $(-\infty, a)$ and τ_0^b corresponds to $S_a^- = \inf\{s > 0 : \xi_s < a\}$. Then,

$$Y_{\tau_0^b} = \exp(\xi_{S_a^-}).$$

So now we are looking for $\mathbb{P}(\xi_{S_a^-} \in dw, S_a^- < \infty)$, for $a < 0$.

Results

Now we are looking for $\mathbb{P}(\xi_{S_a^-} \in dw, S_a^- < \infty)$, for $a < 0$.

Method for $\alpha \in (0, 1]$

Use the ladder process:

$$\begin{aligned}\mathbb{P}(\xi_{S_a^-} \in dw, S_a^- < \infty) &= \mathbb{P}(\underline{\xi}_{S_a^-} \in dw, S_a^- < \infty) \\ &= \mathbb{P}(-\hat{H}_{S_{-a}^+} \in dw) \\ &= \int_{[0, -a]} \hat{U}(dz) \Pi_{\hat{H}}(-dw - z),\end{aligned}$$

recalling that $-\hat{H}$ is a time-change of the running minimum $\underline{\xi}$.

Results

Now we are looking for $\mathbb{P}(\xi_{S_a^-} \in dw, S_a^- < \infty)$, for $a < 0$.

Method for $\alpha \in (1, 2)$

Use the Pecherskii-Rogozin identity:

$$\int_0^\infty \int \exp(qa - \beta(a - \xi_{S_a^-})) d\mathbb{P} da = \frac{\hat{\kappa}(q) - \hat{\kappa}(\beta)}{(q - \beta)\hat{\kappa}(q)},$$

for $a < 0, q, \beta > 0$.

The theorem

Theorem

Let $x > 1$. Then, when $\alpha \in (0, 1]$,

$$\begin{aligned} & P_x(X_{\tau_{-1}^1} \in dy, \tau_{-1}^1 < \infty)/dy \\ &= \frac{\sin(\pi\alpha\hat{\rho})}{\pi} (x+1)^{\alpha\rho} (x-1)^{\alpha\hat{\rho}} (1+y)^{-\alpha\rho} (1-y)^{-\alpha\hat{\rho}} (x-y)^{-1}, \end{aligned}$$

for $y \in (-1, 1)$.

The theorem

Theorem

Let $x > 1$. Then, when $\alpha \in (1, 2)$,

$$\begin{aligned} & P_x(X_{\tau-1}^1 \in dy)/dy \\ &= \frac{\sin(\pi\alpha\hat{\rho})}{\pi} (x+1)^{\alpha\rho} (x-1)^{\alpha\hat{\rho}} (1+y)^{-\alpha\rho} (1-y)^{-\alpha\hat{\rho}} (x-y)^{-1} \\ &\quad - (\alpha-1) \frac{\sin(\pi\alpha\hat{\rho})}{\pi} (1+y)^{-\alpha\rho} (1-y)^{-\alpha\hat{\rho}} \\ &\quad \quad \quad \times \int_1^x (t-1)^{\alpha\hat{\rho}-1} (t+1)^{\alpha\rho-1} dt, \end{aligned}$$

for $y \in (-1, 1)$.

Thank you!