
Galois Automorphisms and Classical Groups

A. A. Schaeffer Fry and Jay Taylor

Abstract. In a previous work, the second-named author gave a complete description of the action of automorphisms on the ordinary irreducible characters of the finite symplectic groups. We generalise this in two directions. Firstly, using work of the first-named author, we give a complete description of the action of the Galois group on irreducible characters. Secondly, we extend both descriptions to cover the case of special orthogonal groups. As a consequence, we obtain explicit descriptions for the character fields of symplectic and special orthogonal groups.

1. Introduction

1.1. Given a finite group G , one has natural actions of its automorphism group $\text{Aut}(G)$ and the absolute Galois group $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ on the set of its ordinary irreducible characters $\text{Irr}(G)$. Many questions in character theory concern these actions. For instance, determining the *character field* $\mathbb{Q}(\chi) = \mathbb{Q}(\chi(g) \mid g \in G)$ of $\chi \in \text{Irr}(G)$ is equivalent to understanding the stabiliser of χ in $\text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$. It is the purpose of this article to study, in some detail, these actions in the case of finite symplectic and special orthogonal groups. Our results sharpen, in these cases, the general statements on character fields obtained by Geck [Gec03] and Tiep–Zaleski [TZ04] and extend previous results in type A [Tur01; SFV19].

The need for such precise information regarding these actions has become increasingly more relevant owing to recent developments regarding the McKay–Navarro Conjecture, sometimes referred to as the Galois–McKay Conjecture. Specifically, this conjecture has just been reduced to a problem about quasi-simple groups by Navarro–Späth–Vallejo [NSV20]. The resulting problem involves understanding explicitly the actions of automorphisms and Galois automorphisms on irreducible characters.

Checking these conditions seems to be extremely complicated, with it already being a challenge in the case of $\text{SL}_2(q)$. In the verification of the McKay conjecture for the prime 2 (recently completed by Malle–Späth [MS16]), special consideration was needed for the case of symplectic groups, see [Mal08]. It stands to reason that the same will be true for the McKay–Navarro conjecture.

To give our first results, let us fix a pair (\mathbf{G}^*, F^*) dual to (\mathbf{G}, F) , where \mathbf{G} is a connected reductive algebraic group and F is a Frobenius morphism on \mathbf{G} yielding an \mathbb{F}_q -rational structure. Given any semisimple element $s \in \mathbf{G}^{*F^*}$, we have a corresponding (rational)

2010 Mathematics Subject Classification. Primary 20C30; Secondary 20C15.

Keywords and phrases. Classical groups, character fields, Galois automorphisms, Harish-Chandra theory, generalised Gelfand–Graev characters, unipotent elements.

Lusztig series $\mathcal{E}(\mathbf{G}^F, s) \subseteq \text{Irr}(\mathbf{G}^F)$. We use $\text{Inn}(\mathbf{G}, F)$ to denote the group of inner diagonal automorphisms, defined precisely in 2.2 below, and $\text{Inn}(\mathbf{G}, F)_\chi$ for the stabiliser of $\chi \in \text{Irr}(\mathbf{G}^F)$ under the natural action.

Theorem A. *Assume $\mathbf{G} = \text{SO}(V)$ is a special orthogonal group defined over \mathbb{F}_q with q odd. Let $F_p, \gamma : \mathbf{G} \rightarrow \mathbf{G}$ be field and graph automorphisms, respectively, with γ the identity when $\dim(V)$ is odd. Assume $s \in \mathbf{G}^{*F^*}$ is a quasi-isolated semisimple element, i.e., $s^2 = 1$, and that s is not contained in the centre of \mathbf{G}^{*F^*} . Then for any $\chi \in \mathcal{E}(\mathbf{G}^F, s)$, the following hold:*

- (i) $\mathbf{Q}(\chi) = \mathbf{Q}$,
- (ii) $\chi^{F_p} = \chi$,
- (iii) $\chi^\gamma = \chi$ if and only if $\text{Inn}(\mathbf{G}, F)_\chi = \text{Inn}(\mathbf{G}^F)$.

1.2. The case when s is central is very well known. In that case (i) and (ii) also hold but (iii) does not. When $\dim(V)$ is odd or q is even, the group $\text{SO}(V)$ has a trivial, hence connected, centre. In these cases, the statements in Theorem A are all easy consequences of the unicity of multiplicities in Deligne–Lusztig virtual characters, see [DM90, Prop. 6.3] or [GM20, Thm. 4.4.23]. However, we include the odd-dimensional case in our analysis as it is little extra effort and some results used on the way to prove Theorem A may be of independent interest.

Our next main result gives the analogue of (i) of Theorem A in the case of symplectic groups. The character fields themselves are a little more complicated in this case, but the behaviour is still uniform across a Lusztig series.

Theorem B. *Assume $\mathbf{G} = \text{Sp}(V)$ is a symplectic group defined over \mathbb{F}_q with q odd. Assume $s \in \mathbf{G}^*$ is quasi-isolated and $\chi \in \mathcal{E}(\mathbf{G}^F, s)$. Then $\mathbf{Q}(\chi) \subseteq \mathbf{Q}(\sqrt{\omega p})$, where $\omega = (-1)^{\frac{p-1}{2}}$ is the unique element of $\{\pm 1\}$ for which $p \equiv \omega \pmod{4}$. Moreover, the following hold:*

- (i) if q is a square or $s = 1$, then $\mathbf{Q}(\chi) = \mathbf{Q}$;
- (ii) if q is not a square and $s \neq 1$, then $\mathbf{Q}(\chi) = \mathbf{Q}(\sqrt{\omega p})$. In particular, if $\sigma \in \text{Gal}(\overline{\mathbf{Q}}/\mathbf{Q})$, then $\chi^\sigma = \chi$ if and only if $\sqrt{\omega p}^\sigma = \sqrt{\omega p}$.

1.3. On the way to Theorem B, we give further details for the action of the Galois group on the characters in this case. Moreover, in Sections 4 and 14, we describe the action of the specific Galois automorphisms relevant for the McKay–Navarro conjecture, which will be useful in studying the conjecture for finite reductive groups. Note that, whilst seemingly only covering special cases, Theorems A and B do allow one to completely determine the field of values for all irreducible characters of \mathbf{G}^F when $\mathbf{G} = \text{SO}(V)$ or $\text{Sp}(V)$. Specifically, we obtain the following:

Theorem C. *Assume $\mathbf{G} = \text{Sp}(V)$ or $\text{SO}(V)$ is defined over \mathbb{F}_q with q odd and let $s \in \mathbf{G}^{*F^*}$ be a semisimple element. Assume s has order $d > 0$ and let $\zeta_d \in \overline{\mathbf{Q}}^\times$ be a primitive d th root of unity. We let $\mathbf{Q}_s = \overline{\mathbf{Q}}^{G_s} \subseteq \mathbf{Q}(\zeta_d)$ be the subfield fixed by the subgroup $G_s \leq G$ stabilising the series $\mathcal{E}(\mathbf{G}^F, s)$, which can be described in terms of conjugacy in \mathbf{G}^{*F^*} , as in Remark 16.8.*

- (i) If $\mathbf{G} = \mathrm{SO}(V)$, then any $\chi \in \mathcal{E}(\mathbf{G}^F, s)$ has character field $\mathbb{Q}(\chi) = \mathbb{Q}_s$.
- (ii) If $\mathbf{G} = \mathrm{Sp}(V)$, then any $\chi \in \mathcal{E}(\mathbf{G}^F, s)$ has character field $\mathbb{Q}(\chi)$, where:
- $\mathbb{Q}(\chi) = \mathbb{Q}_s$ if q is a square or -1 is not an eigenvalue of s ;
 - $\mathbb{Q}_s \subsetneq \mathbb{Q}(\chi) = \mathbb{Q}_s(\sqrt{\omega p})$ otherwise, where $\omega = (-1)^{(p-1)/2}$.

1.4. We note that, assuming q is sufficiently large, it is theoretically possible to compute the character values of the characters of symplectic groups using a result of Waldspurger [Wal04]. However, no such statement is known for the special orthogonal groups, as the results in [Wal04] concern the (disconnected) orthogonal group. Our arguments, which are based on Kawanaka's theory of Generalised Gelfand–Graev Characters (GGGCs) and the Howlett–Lehrer parametrisation of Harish-Chandra series, require no restriction on q and do not require the precise determination of character values.

We now wish to describe two consequences of our description of the action of $\mathrm{Aut}(\mathbf{G}^F)$ on $\mathrm{Irr}(\mathbf{G}^F)$ when $\mathbf{G} = \mathrm{SO}(V)$. For this, let us recall that the automorphism group $\mathrm{Aut}(\mathbf{G}^F)$ has a subgroup $\mathrm{Aso}(\mathbf{G}, F) \leq \mathrm{Aut}(\mathbf{G}^F)$ generated by inner, diagonal, field, and graph automorphisms. This group is defined precisely in 2.2. There is a complement $\Gamma(\mathbf{G}, F) \leq \mathrm{Aso}(\mathbf{G}, F)$ to the subgroup $\mathrm{Inn}(\mathbf{G}, F) \leq \mathrm{Aso}(\mathbf{G}, F)$ such that $\Gamma(\mathbf{G}, F)$ consists of field and graph automorphisms. We choose this explicitly in Lemma 7.9. Recalling the notion of a Jordan decomposition from [DM20, §11.5], we will show the following.

Theorem D. *Assume $\mathbf{G} = \mathrm{SO}(V)$ is a special orthogonal group defined over \mathbb{F}_q with q odd and $\dim(V)$ even. Moreover, let $\gamma : \mathbf{G} \rightarrow \mathbf{G}$ and $\gamma^* : \mathbf{G}^* \rightarrow \mathbf{G}^*$ be dual graph automorphisms.*

- (i) *If $s \in \mathbf{G}^{*\mathbf{F}^*}$ is γ^* -fixed, then there exists a Jordan decomposition $J_s^{\mathbf{G}} : \mathcal{E}(\mathbf{G}^F, s) \rightarrow \mathcal{E}(\mathbf{C}_{\mathbf{G}^*}(s)^{\mathbf{F}^*}, 1)$ satisfying $J_s^{\mathbf{G}}(\chi^\gamma)^{\gamma^*} = J_s^{\mathbf{G}}(\chi)$ for all $\chi \in \mathcal{E}(\mathbf{G}^F, s)$.*
- (ii) *Each $\chi \in \mathrm{Irr}(\mathbf{G}^F)$ satisfies the condition $\mathrm{Aso}(\mathbf{G}, F)_\chi = \mathrm{Inn}(\mathbf{G}, F)_\chi \rtimes \Gamma(\mathbf{G}, F)_\chi$.*

1.5. The statement in (i) of Theorem D was claimed in the proof of a result of Aubert–Michel–Rouquier [AMR96, Prop. 1.7]. Their result states that the orthogonal group $\mathrm{GO}(V)$ also has a Jordan decomposition when $\dim(V)$ is even. The existence of such a Jordan decomposition has recently been used by Vinroot in his work on totally orthogonal groups [Vin20], who noticed the arguments in [AMR96] are insufficient. We thank him for bringing this issue to our attention.

The statement in (ii) of Theorem D was shown to hold in the case $\mathbf{G} = \mathrm{Sp}(V)$ in [Tay18] and independently, by a completely different argument, in [CS17]. Such a statement is of interest owing to Späth's criterion for the Inductive McKay Conditions [Spä12]. We also remark that, using the known actions of $\mathrm{Aut}(\mathbf{G}^F)$ on irreducible characters, Li [Li19, Thm. 5.9] has shown the existence of an equivariant Jordan decomposition when $\mathbf{G} = \mathrm{Sp}(V)$. Similar arguments can be applied to our results here to establish such a statement when $\mathbf{G} = \mathrm{SO}(V)$. However, we have decided not to pursue this.

1.6. Given Theorem C, one might expect that when $\mathbf{G} = \mathrm{SO}(V)$, there is a Jordan decomposition that is equivariant with respect to Galois automorphisms, as in the case of connected centre established in [SV19]. Of course, such a statement cannot hold in general, as it will not hold when $\mathbf{G} = \mathrm{Sp}(V)$. The case of $\mathrm{SL}_2(q) \cong \mathrm{Sp}_2(q)$ already provides a counterexample, as pointed out in [SV19, §5]. Given our work here, the only issue remaining to prove such a statement for $\mathrm{SO}(V)$ is to determine the rationality of unipotent characters of the (disconnected) orthogonal group $\mathrm{GO}(V)$.

With regards to this, we obtain an important partial result, see Theorem 10.8. Namely, we prove that cuspidal unipotent characters of $\mathrm{GO}(V)$ are rational valued. Extending the results of [SF19] to disconnected groups would easily yield the complete description. We intend to consider this in future work.

1.7. We end with a comment on the automorphism group of \mathbf{G}^F when $\mathbf{G} = \mathrm{SO}(V)$ and $\dim(V)$ is even. Under certain conditions on q , we have $Z(\mathbf{G}^F) \leq \mathrm{O}^{p'}(\mathbf{G}^F)$. When this happens, the group \mathbf{G}^F admits an automorphism that is not contained in the group $\mathrm{Aso}(\mathbf{G}, F)$, see Proposition 2.3. The action of this automorphism is closely related to the study of central characters. To give a complete description for the action of $\mathrm{Aut}(\mathbf{G}^F)$ on $\mathrm{Irr}(\mathbf{G}^F)$, we must also describe the action of this automorphism. This is dealt with in Section 15.

Structure of the Paper

1.8. In Sections 2 to 4, we give some generalities on automorphisms and Galois automorphisms, in addition to recalling a few results from the literature to be used in further sections. In this paper, we use Kawanaka's GGGCs to translate the automorphism and Galois actions to questions on unipotent conjugacy classes. To understand the Galois actions, we study the permutation on conjugacy classes induced by power maps in Sections 5 and 9. This allows us to conclude, for instance, that all GGGCs of special orthogonal groups are rational valued.

To understand the actions of $\mathrm{Aut}(\mathbf{G}^F)$ and $\mathrm{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ on irreducible characters of $\mathrm{SO}(V)$, it is necessary to understand such actions on extensions of these characters to $\mathrm{GO}(V)$. In Section 6, we define extensions of GGGCs for disconnected groups and study the relevant automorphism and Galois actions on these characters. To show (i) of Theorem D, we need to describe the action of automorphisms on unipotent characters of certain disconnected groups, which we do in Section 8.

We study the case of quasi-isolated Lusztig series in Sections 10 to 13. This is broken up by studying individual Harish-Chandra series. We first treat the case of cuspidal characters of special orthogonal groups in Section 10, which extends results in [Tay18]. For another approach to the action of $\mathrm{Aut}(\mathbf{G}^F)$ on cuspidal characters see [Mal17]. Hecke algebra techniques and the results of [SF19] are then used in Sections 12 and 13 to study the members of a Harish-Chandra series, first in the special orthogonal group case and then in the symplectic group case. The extra automorphism of $\mathrm{SO}(V)$ mentioned in 1.7 is studied in Section 15. Finally, our reductions to the quasi-isolated case, as well as the proof of Theorems C and D, are contained in Section 16.

Notation

1.9. For a set X and elements $x, y \in X$, we denote by $\delta_{x,y} \in \{0, 1\}$ the usual Kronecker delta. Assume G is a group. If G acts on a set X , then we denote by $G_x \leq G$ the stabiliser of $x \in X$. We denote by $\text{Aut}(G)$ the automorphism group of G as an abstract group, ignoring any additional structure on G . For any $g \in G$, we will use $\iota_g : G \rightarrow G$ to denote the inner automorphism defined by $\iota_g(x) = {}^g x = x^{g^{-1}} = gxg^{-1}$ and $\text{Inn}(G) \leq \text{Aut}(G)$ to denote the inner automorphism group.

Assume S is an arbitrary set. If $\phi : G \rightarrow H$ is a group isomorphism, then for any function $f : G \rightarrow S$, respectively $f : H \rightarrow S$, we let ${}^\phi f = f \circ \phi^{-1} : H \rightarrow S$, respectively $f^\phi = f \circ \phi : G \rightarrow S$. This defines left and right actions of $\text{Aut}(G)$ on functions $f : G \rightarrow S$. If $g \in G$, then we also let ${}^g f = \iota_g f$ and $f^g = f^{\iota_g}$. Throughout, we will usually use right actions, although we will use the left action when it is notationally convenient.

We fix a prime p and denote by $\mathbb{F} = \overline{\mathbb{F}}_p$ an algebraic closure of the finite field \mathbb{F}_p . Any algebraic group \mathbf{G} , usually denoted in bold, is assumed to be affine and defined over \mathbb{F} . We denote by \mathbf{G}° the connected component. For any $x \in \mathbf{G}$, we denote by $A_{\mathbf{G}}(x)$ the component group $C_{\mathbf{G}}(x)/C_{\mathbf{G}}^\circ(x)$ of the centraliser. Note that if $x \in \mathbf{G}^\circ$, then $A_{\mathbf{G}^\circ}(x) \leq A_{\mathbf{G}}(x)$ is a subgroup, since $C_{\mathbf{G}^\circ}^\circ(x) = C_{\mathbf{G}^\circ}^\circ(x)$.

We fix a prime $\ell \neq p$ and let $\overline{\mathbb{Q}}_\ell$ be an algebraic closure of the ℓ -adic numbers. We identify $\overline{\mathbb{Q}}$ with a subfield of $\overline{\mathbb{Q}}_\ell$. If G is a finite group, then $\text{Irr}(G)$ is the set of $\overline{\mathbb{Q}}_\ell$ -irreducible characters of G . We denote by $1_G \in \text{Irr}(G)$ the trivial character. If G is cyclic, we denote by $\varepsilon_G \in \text{Irr}(G)$ the unique character with kernel $\{x^2 \mid x \in G\}$. For each $\chi \in \text{Irr}(G)$, we have the central character $\omega_\chi : Z(G) \rightarrow \overline{\mathbb{Q}}_\ell$ defined by $\omega_\chi(z) = \chi(z)/\chi(1)$. For a subgroup $H \leq G$ and characters $\chi \in \text{Irr}(G)$ and $\varphi \in \text{Irr}(H)$, we will write $\text{Res}_H^G(\chi)$ and $\text{Ind}_H^G(\varphi)$ for the corresponding restricted and induced characters, respectively.

Throughout, we let $\mathcal{G} := \text{Gal}(\overline{\mathbb{Q}}/\mathbb{Q})$ denote the absolute Galois group of \mathbb{Q} . Note that if G is a finite group and $\chi \in \text{Irr}(G)$, then $\chi(g) \in \overline{\mathbb{Q}}$. For any $\xi \in \overline{\mathbb{Q}}$ and $\sigma \in \mathcal{G}$, we let ξ^σ be the image of ξ under σ . We have an action of \mathcal{G} on $\text{Irr}(G)$ given by $\chi^\sigma(g) = \chi(g)^\sigma$. For each character χ of G , we fix a $\overline{\mathbb{Q}}_\ell G$ -module M_χ affording χ . We then write $\text{End}_G(\chi)$ to denote the endomorphism algebra $\text{End}_{\overline{\mathbb{Q}}_\ell G}(M_\chi)$.

2. Automorphisms

Lemma 2.1. *Assume G is a group and $A \leq \text{Aut}(G)$ is a subgroup. Let $N \triangleleft G$ be an A -invariant normal subgroup. We denote by $K \leq A$ the kernel of the natural restriction map $A \rightarrow \text{Aut}(N)$ and let $Z = N \cap Z(G)$.*

- (i) *If $\alpha \in \text{Hom}(G/N, Z)$ then we have an automorphism $\tau_\alpha \in \text{Aut}(G)$ given by $\tau_\alpha(g) = g \cdot \alpha(gN)$. Moreover, $\phi \tau_\alpha \phi^{-1} = \tau_{\phi \alpha \phi^{-1}}$ for any $\phi \in A$.*
- (ii) *If $C_G(N) = Z(G)$ then $K \leq \{\tau_\alpha \mid \alpha \in \text{Hom}(G/N, Z)\}$ and $K \cap \text{Inn}(G) = \{1\}$.*
- (iii) *If G is finite, $\chi \in \text{Irr}(G)$, and $\alpha \in \text{Hom}(G/N, Z)$, then $\chi^{\tau_\alpha} = (\omega_\chi \circ \alpha)\chi$, where τ_α is as in (i) and $\omega_\chi \circ \alpha \in \text{Irr}(G)$ is a linear character.*

Proof. (i). Clearly τ_α is a homomorphism, and if $\tau_\alpha(g) = 1$, then $g \cdot \alpha(gN) = 1$ so $g \in Z \leq N$ which implies $\alpha(gN) = 1$, hence $g = 1$. If $g \in G$, then $z = \alpha(gN) \in Z \leq N$ and we have $\tau_\alpha(gz^{-1}) = gz^{-1}\alpha(gz^{-1}N) = gz^{-1}\alpha(gN) = g$ so the map is surjective, hence an automorphism. The last statement is clear.

(ii). Assume $\varphi \in K$. For any $g \in G$ and $x \in N$ we have ${}^g x = \varphi({}^g x) = \varphi({}^g)x$ so $g^{-1}\varphi(g) \in C_G(N) = Z(G)$. If $\alpha : G \rightarrow Z(G)$ is the map given by $\alpha(g) = g^{-1}\varphi(g)$ then α is a homomorphism factoring through G/N and $\varphi = \tau_\alpha$. If $\varphi = \iota_h$ for some $h \in G$ then we must have $h \in C_G(N) = Z(G)$ so φ is trivial.

(iii). This follows from the definitions. ■

2.2. Assume \mathbf{G} is an affine algebraic group over \mathbb{F} (not necessarily connected). A bijective homomorphism of algebraic groups $\varphi : \mathbf{G} \rightarrow \mathbf{G}$ will be called an *asogeny*, and we denote by $\text{Aso}(\mathbf{G}) \leq \text{Aut}(\mathbf{G})$ the *submonoid* consisting of all asogenies. For any Frobenius root $F : \mathbf{G} \rightarrow \mathbf{G}$, in the sense of [DM20, Def. 4.2.4], we have $F \in \text{Aso}(\mathbf{G})$. Restricting to \mathbf{G}^F , we obtain a natural monoid homomorphism $C_{\text{Aso}(\mathbf{G})}(F) \rightarrow \text{Aut}(\mathbf{G}^F)$. The subgroup of $\text{Aut}(\mathbf{G}^F)$ generated by the image of this map will be denoted by $\text{Aso}(\mathbf{G}, F)$. If $\mathbf{H} \leq \mathbf{G}$ is an F -stable subgroup then we let $\text{Inn}(\mathbf{H}, F) \leq \text{Aso}(\mathbf{G}, F)$ be the image of $C_{\text{Inn}(\mathbf{H})}(F)$, which is a subgroup containing $\text{Inn}(\mathbf{H}^F)$. We define $\text{Out}(\mathbf{G}, F)$ to be the quotient $\text{Aso}(\mathbf{G}, F)/\text{Inn}(\mathbf{G}^F)$.

Recall that if G is a finite group, then $\text{Op}'(G) \triangleleft G$ is the unique minimal normal subgroup whose quotient $G/\text{Op}'(G)$ is a p' -group. Equivalently, this is the subgroup generated by all p -elements of G . Using the notation of Lemma 2.1, we define a subgroup

$$K(G) = \{\tau_\alpha \mid \alpha \in \text{Hom}(G/\text{Op}'(G), Z(G) \cap \text{Op}'(G))\} \leq \text{Aut}(G)$$

of the automorphism group of G . The following is well known to finite group theorists, but we include some details in the proof as it is difficult to extract this exact statement from the literature.

Proposition 2.3. *Assume \mathbf{G} is a quasisimple algebraic group with $F : \mathbf{G} \rightarrow \mathbf{G}$ a Frobenius root. Then the map $C_{\text{Aso}(\mathbf{G})}(F) \rightarrow \text{Aso}(\mathbf{G}, F)$ is surjective and $\text{Aut}(\mathbf{G}^F) = K(\mathbf{G}^F) \rtimes \text{Aso}(\mathbf{G}, F)$.*

Proof. The subgroup $\text{Aut}_1(\mathbf{G}) \leq \text{Aut}(\mathbf{G})$ generated by $\text{Aso}(\mathbf{G})$ is described in [GLS98, Thm. 1.15.7]. Picking a set of Steinberg generators for \mathbf{G} , we have a splitting $\text{Aut}_1(\mathbf{G}) = \text{Inn}(\mathbf{G}) \rtimes \Gamma(\mathbf{G})$, where $\Gamma(\mathbf{G})$ is generated by field and graph automorphisms. In fact $\Gamma(\mathbf{G})$ is generated, as a group, by graph automorphisms whose inverse is also an asogeny and an element ψ , which is either a standard Frobenius F_p or $\psi^2 = F_p$ is such. In any case, one sees that the submonoid of $\text{Aut}(\mathbf{G})$ generated by $\text{Aso}(\mathbf{G})$ and $F_p^{-1} \in \text{Aut}(\mathbf{G})$ is the subgroup $\text{Aut}_1(\mathbf{G})$.

Up to conjugacy, we can assume $F \in \Gamma(\mathbf{G})$, so that $C_{\text{Aso}(\mathbf{G})}(F) = C_{\text{Inn}(\mathbf{G})}(F) \rtimes C_{\Gamma(\mathbf{G})}(F)$. We have $F_p \in C_{\Gamma(\mathbf{G})}(F)$ and so $\text{Aso}(\mathbf{G}, F)$ is generated by the image of $C_{\text{Aso}(\mathbf{G})}(F)$ and the image of F_p^{-1} in $\text{Aut}(\mathbf{G}^F)$. However, $\langle F_p \rangle \leq \text{Aut}(\mathbf{G}^F)$ is finite, so $C_{\text{Aso}(\mathbf{G})}(F)$ contains a preimage of the image of F_p^{-1} . This implies $C_{\text{Aso}(\mathbf{G})}(F) \rightarrow \text{Aso}(\mathbf{G}, F)$ is surjective.

By the proof of [Bon06, Lem. 6.1], there exist two unipotent elements $u, v \in \mathbf{G}^F$ such that $C_{\mathbf{G}}(u) \cap C_{\mathbf{G}}(v) = Z(\mathbf{G})$. These are clearly contained in $\text{Op}'(\mathbf{G}^F)$, so we have

$C_{\mathbf{G}^F}(\mathrm{OP}'(\mathbf{G}^F)) \leq Z(\mathbf{G}) \cap \mathbf{G}^F = Z(\mathbf{G})^F$. Hence $C_{\mathbf{G}^F}(\mathrm{OP}'(\mathbf{G}^F)) = Z(\mathbf{G}^F) = Z(\mathbf{G})^F$. According to [GLS98, 2.5.14(g)] the restriction map $C_{\mathrm{Aso}(\mathbf{G})}(\mathbf{F}) \rightarrow \mathrm{Aut}(\mathrm{OP}'(\mathbf{G}^F))$ is surjective. Therefore, so is the map $\mathrm{Aso}(\mathbf{G}, \mathbf{F}) \rightarrow \mathrm{Aut}(\mathrm{OP}'(\mathbf{G}^F))$. It follows from (ii) of Lemma 2.1 that $\mathrm{Aut}(\mathbf{G}^F) = \mathbf{K}(\mathbf{G}^F) \mathrm{Aso}(\mathbf{G}, \mathbf{F})$.

We now need only show the intersection is trivial. For this, let $\pi : \mathbf{G} \rightarrow \mathbf{G}_{\mathrm{ad}}$ be an adjoint quotient of \mathbf{G} and let $F : \mathbf{G}_{\mathrm{ad}} \rightarrow \mathbf{G}_{\mathrm{ad}}$ be the Frobenius root such that $F \circ \pi = \pi \circ F$. By [GLS98, 1.15.6, 2.5.14], restriction through π defines injective homomorphisms $C_{\mathrm{Aso}(\mathbf{G})}(\mathbf{F}) \rightarrow C_{\mathrm{Aso}(\mathbf{G}_{\mathrm{ad}})}(\mathbf{F})$ and $\mathrm{Aut}(\mathrm{OP}'(\mathbf{G}^F)) \rightarrow \mathrm{Aut}(\mathrm{OP}'(\mathbf{G}_{\mathrm{ad}}^F))$. Suppose $\gamma \in C_{\mathrm{Aso}(\mathbf{G})}(\mathbf{F})$ is the identity on $\mathrm{OP}'(\mathbf{G}^F)$. Then $\gamma \circ \pi$ is the identity on $\mathrm{OP}'(\mathbf{G}_{\mathrm{ad}}^F)$ so $\gamma \circ \pi \in \langle F \rangle$ by [GLS98, 2.5.7]. Therefore $\gamma \in \langle F \rangle$ so γ is the identity on \mathbf{G}^F . ■

After (iii) of Lemma 2.1, we see that to understand the action of $\mathbf{K}(\mathbf{G}^F)$ on irreducible characters, it is crucial to have a good understanding of central and linear characters. For convenience, we recall a few results from the literature regarding this. We let $\mathcal{C}(\mathbf{G}, \mathbf{F})$ and $\mathcal{S}(\mathbf{G}^*, \mathbf{F}^*)$ be as in [Tay18].

Proposition 2.4. *Assume \mathbf{G} is a connected reductive algebraic group with Frobenius root $F : \mathbf{G} \rightarrow \mathbf{G}$. Let $(\mathbf{G}^*, \mathbf{F}^*)$ be dual to (\mathbf{G}, \mathbf{F}) .*

- (i) *For any semisimple element $s \in \mathbf{G}^{*\mathbf{F}^*}$, there exists a unique character $\omega_s \in \mathrm{Irr}(Z(\mathbf{G})^F)$ such that $\omega_s = \mathrm{Res}_{Z(\mathbf{G})^F}^{\mathbf{G}^F}(\theta)$ for any pair $(\mathbf{T}, \theta) \in \mathcal{C}(\mathbf{G}, \mathbf{F})$ corresponding to $(\mathbf{T}^*, s) \in \mathcal{S}(\mathbf{G}^*, \mathbf{F}^*)$ under duality. Moreover, $\omega_\chi = \omega_s$ for all $\chi \in \mathcal{E}(\mathbf{G}^F, s)$.*
- (ii) *There is a group isomorphism $Z(\mathbf{G}^*)^{\mathbf{F}^*} \rightarrow \mathrm{Irr}(\mathbf{G}^F / \mathrm{OP}'(\mathbf{G}^F))$, denoted by $z \mapsto \hat{z}$, such that $-\otimes \hat{z} : \mathcal{E}(\mathbf{G}^F, s) \rightarrow \mathcal{E}(\mathbf{G}^F, sz)$ is a bijection.*
- (iii) *We have a well-defined group isomorphism $\mathbf{G}^{*\mathbf{F}^*} / \mathrm{OP}'(\mathbf{G}^{*\mathbf{F}^*}) \rightarrow \mathrm{Irr}(Z(\mathbf{G})^F)$ given by $s\mathrm{OP}'(\mathbf{G}^{*\mathbf{F}^*}) \mapsto \omega_s$, where $s \in \mathbf{G}^{*\mathbf{F}^*}$ is a semisimple element representing the coset.*

Proof. (i) is [Bon06, 11.1(d)] and (ii) is [DM20, 11.4.12, 11.4.14]. Dualising the arguments in [DM20] gives (iii), see also [NT13, Lem. 4.4]. ■

Corollary 2.5. *Assume $s \in \mathbf{G}^{*\mathbf{F}^*}$ is a semisimple element and $\sigma = \tau_\alpha \in \mathbf{K}(\mathbf{G}^F)$. Then there exists a central element $z \in Z(\mathbf{G}^*)^{\mathbf{F}^*}$ such that $\omega_s \circ \alpha = \hat{z}$ and $\mathcal{E}(\mathbf{G}^F, s)^\sigma = \mathcal{E}(\mathbf{G}^F, sz)$.*

Proof. After (iii) of Lemma 2.1, this follows from the definition of Lusztig series and the property of Deligne–Lusztig virtual characters in [Bon06, 11.1(d)]. ■

Remark 2.6. As is well known, the elements of $\mathrm{Aso}(\mathbf{G}, \mathbf{F})$ permute rational and geometric Lusztig series, see [NTT08, Cor. 2.4] and [Tay18, Prop. 7.2]. However, this is not necessarily the case for elements of $\mathbf{K}(\mathbf{G}^F)$. It follows from Corollary 2.5 that the elements of $\mathbf{K}(\mathbf{G}^F)$ will permute rational Lusztig series. However, they do not necessarily permute geometric Lusztig series. This is due to the fact that, in a geometric Lusztig series, characters can have both trivial and non-trivial central character.

3. Galois Automorphisms

3.1. Let \mathbf{G} be a connected reductive algebraic group and $F : \mathbf{G} \rightarrow \mathbf{G}$ a Frobenius endomorphism endowing \mathbf{G} with an \mathbb{F}_q -rational structure. Let $\mathbf{T} \leq \mathbf{B}$ be, respectively, an F -stable maximal torus and Borel subgroup of \mathbf{G} . We denote by $\mathbf{W} = N_{\mathbf{G}}(\mathbf{T})/\mathbf{T}$ the corresponding Weyl group with simple reflections $S \subseteq \mathbf{W}$ determined by \mathbf{B} . Let Φ be the root system of \mathbf{G} with respect to \mathbf{T} and $\Delta \subseteq \Phi^+$ the set of simple and positive roots determined by \mathbf{B} .

For each subset $J \subseteq \Delta$ of simple roots, we have a corresponding standard parabolic and Levi subgroup $\mathbf{L}_J \leq \mathbf{P}_J \leq \mathbf{G}$. We denote by $\text{Cusp}(\mathbf{G}, F)$ the set of *cuspidal pairs* (\mathbf{L}_J, λ) with $J \subseteq \Delta$ an F -invariant subset of simple roots and $\lambda \in \text{Irr}(\mathbf{L}_J^F)$ a cuspidal character. To each such pair $(\mathbf{L}, \lambda) \in \text{Cusp}(\mathbf{G}, F)$, we have a corresponding Harish-Chandra series $\mathcal{E}(\mathbf{G}^F, \mathbf{L}, \lambda)$. By definition, this is the set of irreducible constituents of the corresponding Harish-Chandra induced character $\mathbf{R}_{\mathbf{L}}^{\mathbf{G}}(\lambda)$.

Thanks to [Gec93] and [Lus84, Thm. 8.6], we know that any cuspidal character $\lambda \in \text{Irr}(\mathbf{L}^F)$ extends to its inertia group $N_{\mathbf{G}^F}(\mathbf{L})_{\lambda}$. Hence we may fix a so-called *extension map* Λ with respect to $\mathbf{L}^F \triangleleft N_{\mathbf{G}^F}(\mathbf{L})$, meaning that for each cuspidal character $\lambda \in \text{Irr}(\mathbf{L}^F)$, $\Lambda(\lambda)$ is an extension of λ to $N_{\mathbf{G}^F}(\mathbf{L})_{\lambda}$. We denote by $W(\lambda) := N_{\mathbf{G}^F}(\mathbf{L})_{\lambda}/\mathbf{L}^F$ the so-called relative Weyl group. After choosing a square root $\sqrt{p} \in \overline{\mathbf{Q}} \subseteq \overline{\mathbf{Q}}_{\ell}$ of p , we have a canonical bijection $\text{Irr}(W(\lambda)) \rightarrow \mathcal{E}(\mathbf{G}^F, \mathbf{L}_J, \lambda)$, see [Car93, Chap. 10], which we denote by $\eta \mapsto \mathbf{R}_{\mathbf{L}}^{\mathbf{G}}(\lambda)_{\eta}$.

The group $W(\lambda)$ is a semidirect product $R(\lambda) \rtimes C(\lambda)$ of a Weyl group $R(\lambda)$ with a root system Φ_{λ} and the stabilizer $C(\lambda)$ in $W(\lambda)$ of a simple system in Φ_{λ} , see [Car93, Chapter 10]. Note that $W(\lambda)$ in general is not necessarily a reflection group but $R(\lambda)$ is. Given $\sigma \in \mathcal{G}$ and $\eta \in \text{Irr}(W(\lambda))$ we define $\eta^{(\sigma)} \in \text{Irr}(W(\lambda^{\sigma}))$ as in [SF19, §3.5]. Let $w_0 \in \mathbf{W}$ be the longest element with respect to S . For any $w \in \mathbf{W}^F$ we write $\text{ind}(w) := |\mathbf{U} \cap \mathbf{U}^{w_0 w}|$, where $\mathbf{U} = \mathbf{U}^F$ and $\mathbf{U} \leq \mathbf{B}$ is the unipotent radical of \mathbf{B} . With this, we now recall [SF19, Thm. 3.8], which will be key to our proofs of Theorems A(i) and B.

Theorem 3.2 (Schaeffer Fry, [SF19, Thm. 3.8]). *Assume $\sigma \in \mathcal{G}$ and $(\mathbf{L}, \lambda) \in \text{Cusp}(\mathbf{G}, F)$ is a cuspidal pair. Let Λ be an extension map with respect to $\mathbf{L}^F \triangleleft N_{\mathbf{G}^F}(\mathbf{L})$. Define:*

- $\delta_{\lambda, \sigma}$ to be the linear character of $W(\lambda)$ such that $\Lambda(\lambda)^{\sigma} = \delta_{\lambda, \sigma} \Lambda(\lambda^{\sigma})$;
- $\delta'_{\lambda, \sigma} \in \text{Irr}(W(\lambda))$ to be the character such that $\delta'_{\lambda, \sigma}(w) = \delta_{\lambda, \sigma}(w)$ for $w \in C(\lambda)$ and $\delta'_{\lambda, \sigma}(w) = 1$ for $w \in R(\lambda)$; and
- $\gamma_{\lambda, \sigma}$ to be the function on $W(\lambda)$ such that $\gamma_{\lambda, \sigma}(w) = \frac{\sqrt{\text{ind}(w_2)}^{\sigma}}{\sqrt{\text{ind}(w_2)}}$, where $w = w_1 w_2$ for $w_1 \in R(\lambda)$ and $w_2 \in C(\lambda)$.

Then for any $\eta \in \text{Irr}(W(\lambda))$, we have

$$(\mathbf{R}_{\mathbf{L}}^{\mathbf{G}}(\lambda)_{\eta})^{\sigma} = \mathbf{R}_{\mathbf{L}}^{\mathbf{G}}(\lambda^{\sigma})_{\eta'},$$

where $\eta' \in \text{Irr}(W(\lambda)) = \text{Irr}(W(\lambda^{\sigma}))$ is defined by $\eta'(w) = \gamma_{\lambda, \sigma}(w) \delta'_{\lambda, \sigma}(w^{-1}) \eta^{(\sigma)}(w)$ for each $w \in W(\lambda)$.

On the group $W(\lambda)$

3.3. For each $w \in \mathbf{W}$, we fix a representative $n_w \in N_{\mathbf{G}}(\mathbf{T})$ such that if $w \in \mathbf{W}^F$, then $n_w \in N_{\mathbf{G}^F}(\mathbf{T})$. If $J \subseteq \Delta$ is an F -stable subset of simple roots, then we have an isomorphism $N_{\mathbf{W}^F}(J) \rightarrow N_{\mathbf{G}^F}(\mathbf{L}_J)/\mathbf{L}_J^F$, given by $w \mapsto n_w \mathbf{L}_J^F$, where $N_{\mathbf{W}^F}(J) = \{w \in \mathbf{W}^F \mid wJ = J\}$. Given a cuspidal pair $(\mathbf{L}_J, \lambda) \in \text{Cusp}(\mathbf{G}, F)$, the group $W(\lambda) \leq N_{\mathbf{G}^F}(\mathbf{L}_J)/\mathbf{L}_J^F$ can thus be identified with the subgroup $N_{\mathbf{W}^F}(J)_\lambda \leq N_{\mathbf{W}^F}(J)$, and hence a subgroup of \mathbf{W} . We will make this identification implicitly in what follows.

Now assume $\iota : \mathbf{G} \rightarrow \tilde{\mathbf{G}}$ is an F -equivariant regular embedding of \mathbf{G} into a group with connected centre and let us identify \mathbf{G} with its image $\iota(\mathbf{G})$. Given $J \subseteq \Delta$, we have a corresponding F -stable Levi $\tilde{\mathbf{L}}_J = \mathbf{L}_J \cdot Z(\tilde{\mathbf{G}})$ of $\tilde{\mathbf{G}}$. If $\lambda \in \text{Irr}(\mathbf{L}_J^F)$ is a cuspidal character, then any character $\tilde{\lambda} \in \text{Irr}(\tilde{\mathbf{L}}_J^F)$ covering λ is also cuspidal. This is standard with respect to the maximal torus $\tilde{\mathbf{T}} = \mathbf{T} \cdot Z(\tilde{\mathbf{G}})$ and Borel $\tilde{\mathbf{B}} = \mathbf{B} \cdot Z(\tilde{\mathbf{G}})$ of $\tilde{\mathbf{G}}$. Note that the Weyl group $\tilde{\mathbf{W}} = N_{\tilde{\mathbf{G}}}(\tilde{\mathbf{T}})/\tilde{\mathbf{T}}$ is naturally isomorphic to \mathbf{W} . We let $W(\tilde{\lambda}) = N_{\mathbf{G}^F}(\mathbf{L}_J)_{\tilde{\lambda}}/\mathbf{L}_J^F$, which is a subgroup of $W(\lambda)$ by [Bon06, Cor. 12.5(c)].

Lemma 3.4. *Assume $(\mathbf{L}, \lambda) \in \text{Cusp}(\mathbf{G}, F)$ is a cuspidal pair and $\tilde{\lambda} \in \text{Irr}(\tilde{\mathbf{L}})$ covers λ . Then $R(\lambda) = W(\tilde{\lambda})$ and there is an injective homomorphism $C(\lambda) \rightarrow H^1(F, Z(\mathbf{G}))$ where $Z(\mathbf{G}) = Z(\mathbf{G})/Z^\circ(\mathbf{G})$.*

Proof. Let $\tilde{W}(\tilde{\lambda}) = \tilde{R}(\tilde{\lambda}) \rtimes \tilde{C}(\tilde{\lambda})$ be as in 3.1 with respect to $\tilde{\mathbf{G}}^F$. By a result of Lusztig, $\tilde{C}(\tilde{\lambda}) = \{1\}$, see [Lus84, Thm. 8.6]. For the first statement, it suffices to show that $\tilde{R}(\tilde{\lambda}) = R(\lambda)$, where we identify $\tilde{W}(\tilde{\lambda})$ and $W(\tilde{\lambda})$. For this, let $\mathbf{T} \leq \mathbf{P} \leq \mathbf{G}$ be an F -stable parabolic subgroup with Levi complement $\mathbf{T} \leq \mathbf{M} \leq \mathbf{P}$. We assume $\mathbf{L} \subsetneq \mathbf{M}$ and \mathbf{M} is minimal with this property. As above, we set $\tilde{\mathbf{M}} = \mathbf{M} \cdot Z(\tilde{\mathbf{G}})$, so $\tilde{\mathbf{L}} \subsetneq \tilde{\mathbf{M}}$.

If $W_{\mathbf{M}^F}(\mathbf{L}) := N_{\mathbf{M}^F}(\mathbf{L})/\mathbf{L}^F$ is trivial then there is nothing to consider. So suppose $W_{\mathbf{M}^F}(\mathbf{L})$ is non-trivial, and hence so is $W_{\tilde{\mathbf{M}}^F}(\tilde{\mathbf{L}}) := N_{\tilde{\mathbf{M}}^F}(\tilde{\mathbf{L}})/\tilde{\mathbf{L}}^F$. By minimality, $W_{\mathbf{M}^F}(\mathbf{L}) \cong W_{\tilde{\mathbf{M}}^F}(\tilde{\mathbf{L}})$ is cyclic of order 2. We will assume that $W_{\mathbf{M}^F}(\mathbf{L})_\lambda = N_{\mathbf{M}^F}(\mathbf{L})_\lambda/\mathbf{L}^F$ is non-trivial. Then $R_{\mathbf{L}}^{\mathbf{M}}(\lambda) = \lambda_1 + \lambda_2$ with $\lambda_1(1) \leq \lambda_2(1)$ and $\lambda_i \in \text{Irr}(\mathbf{M}^F)$. The unique non-trivial element $1 \neq w \in W_{\mathbf{M}^F}(\mathbf{L})_\lambda$ is a generator of $R(\lambda)$ if and only if $\lambda_1(1) \neq \lambda_2(1)$.

As Harish-Chandra induction is compatible with restriction and conjugation, we have

$$\text{Res}_{\mathbf{M}^F}^{\tilde{\mathbf{M}}^F} \left(R_{\tilde{\mathbf{L}}}^{\tilde{\mathbf{M}}}(\tilde{\lambda}) \right) = R_{\mathbf{L}}^{\mathbf{M}} \left(\text{Res}_{\tilde{\mathbf{L}}^F}^{\mathbf{M}^F}(\tilde{\lambda}) \right) = \sum_{g \in \tilde{\mathbf{L}}^F/(\tilde{\mathbf{L}}^F)_\lambda} R_{\mathbf{L}}^{\mathbf{M}}(\lambda)^g = \sum_{g \in \tilde{\mathbf{L}}^F/(\tilde{\mathbf{L}}^F)_\lambda} \lambda_1^g + \lambda_2^g. \quad (3.5)$$

If $\lambda_1(1) \neq \lambda_2(1)$ then these characters cannot be conjugate in $\tilde{\mathbf{M}}^F$, so by Clifford's Theorem $R_{\tilde{\mathbf{L}}}^{\tilde{\mathbf{M}}}(\tilde{\lambda}) = \tilde{\lambda}_1 + \tilde{\lambda}_2$ with $\tilde{\lambda}_i \in \text{Irr}(\tilde{\mathbf{M}}^F)$ an irreducible character with degree $[\tilde{\mathbf{L}}^F : (\tilde{\mathbf{L}}^F)_\lambda] \lambda_i(1)$. Hence, $\tilde{\lambda}_1(1) \neq \tilde{\lambda}_2(1)$ so $R(\lambda) \leq R(\tilde{\lambda})$.

Conversely, if $W_{\mathbf{M}^F}(\mathbf{L})_{\tilde{\lambda}} = N_{\mathbf{M}^F}(\mathbf{L})_{\tilde{\lambda}}/\mathbf{L}^F$ is non-trivial, then $R_{\tilde{\mathbf{L}}}^{\tilde{\mathbf{M}}}(\tilde{\lambda}) = \tilde{\lambda}_1 + \tilde{\lambda}_2$ with $\tilde{\lambda}_i \in \text{Irr}(\tilde{\mathbf{M}}^F)$. We must have $W_{\mathbf{M}^F}(\mathbf{L})_\lambda$ is also non-trivial, by [Bon06, Cor. 12.5(c)], so $R_{\mathbf{L}}^{\mathbf{M}}(\lambda) = \lambda_1 + \lambda_2$ as above. We can compare

$$\text{Res}_{\mathbf{M}^F}^{\tilde{\mathbf{M}}^F} \left(R_{\tilde{\mathbf{L}}}^{\tilde{\mathbf{M}}}(\tilde{\lambda}) \right) = \text{Res}_{\mathbf{M}^F}^{\tilde{\mathbf{M}}^F}(\tilde{\lambda}_1 + \tilde{\lambda}_2)$$

with the terms in (3.5). If λ_1 is a constituent of $\text{Res}_{\widetilde{\mathbf{M}}^F}^{\widetilde{\mathbf{M}}^F}(\widetilde{\lambda}_i)$, then $\sum_{g \in \widetilde{\mathbf{L}}^F / (\widetilde{\mathbf{L}}^F)_\lambda} \lambda_1^g$ must be also, as $\widetilde{\mathbf{L}}^F \leq \widetilde{\mathbf{M}}^F$ and the restriction is invariant under $\widetilde{\mathbf{M}}^F$.

If λ_1 and λ_2 were both constituents of $\text{Res}_{\widetilde{\mathbf{M}}^F}^{\widetilde{\mathbf{M}}^F}(\widetilde{\lambda}_1)$, say, then we would have to have $\text{Res}_{\widetilde{\mathbf{M}}^F}^{\widetilde{\mathbf{M}}^F}(\widetilde{\lambda}_2) = 0$, which is impossible. If $\widetilde{\lambda}_1(1) \neq \widetilde{\lambda}_2(1)$ then $\text{Res}_{\widetilde{\mathbf{M}}^F}^{\widetilde{\mathbf{M}}^F}(\widetilde{\lambda}_i) = \sum_{g \in \widetilde{\mathbf{L}}^F / (\widetilde{\mathbf{L}}^F)_\lambda} \lambda_i^g$ so $\widetilde{\lambda}_i(1) = [\widetilde{\mathbf{L}}^F : (\widetilde{\mathbf{L}}^F)_\lambda] \lambda_i(1)$ and $\lambda_1(1) \neq \lambda_2(1)$. This gives $R(\widetilde{\lambda}) \leq R(\lambda)$.

For the final statement, note that if $g \in N_{\mathbf{G}^F}(\mathbf{L})_\lambda$, then $\widetilde{\lambda}^g \in \text{Irr}(\widetilde{\mathbf{L}}^F)$ also covers λ . Thus $\widetilde{\lambda}^g = \widetilde{\lambda}\theta$ where θ is the inflation of a linear character of $\widetilde{\mathbf{L}}^F / \mathbf{L}^F Z(\widetilde{\mathbf{L}})^F \cong H^1(F, Z(\mathbf{G}))$. This gives a homomorphism $W(\lambda) \rightarrow H^1(F, Z(\mathbf{G}))$ with kernel $W(\widetilde{\lambda}) = R(\lambda)$. \blacksquare

On the Characters $\gamma_{\lambda, \sigma}$ and $\delta_{\lambda, \sigma}$

In the latter parts of this paper, we will need to carefully study the characters $\delta_{\lambda, \sigma}$ and $\gamma_{\lambda, \sigma}$ introduced in Theorem 3.2. Here we introduce a few general statements that will be used later on. For the rest of this section, we fix a cuspidal pair $(\mathbf{L}, \lambda) \in \text{Cusp}(\mathbf{G}, F)$ and a Galois automorphism $\sigma \in \mathcal{G}$.

Lemma 3.6. *Assume $(\mathbf{L}, \lambda) \in \text{Cusp}(\mathbf{G}, F)$ and $\sigma \in \mathcal{G}$. Then $\gamma_{\lambda, \sigma}$ is a character of $W(\lambda)$, and is moreover trivial if and only if at least one of the following holds:*

- q is a square;
- the length $l(w_2)$, with respect to (\mathbf{W}, \mathbf{S}) , is even for each $w_2 \in C(\lambda)$; or
- σ fixes \sqrt{p} .

Otherwise, $\gamma_{\lambda, \sigma}(w) = (-1)^{l(w_2)}$, where $w = w_1 w_2$ with $w_1 \in R(\lambda)$ and $w_2 \in C(\lambda)$.

Proof. By [Car93, §2.9] we have $\text{ind}(w_2) = q^{l(w_2)}$ with $l(w_2)$ as above. Therefore, $\gamma_{\lambda, \sigma}$ is a character in the case of finite reductive groups. In particular, $\gamma_{\lambda, \sigma}(w) = 1$ when $\sqrt{q^{l(w_2)}}$ is fixed by σ and $\gamma_{\lambda, \sigma}(w) = -1$ otherwise. \blacksquare

3.7. When $\mathbf{L} = \mathbf{T}$, we have $N_{\mathbf{G}^F}(\mathbf{T})$ is generated by \mathbf{T}^F and the group $\langle n_\alpha(\pm 1) \mid \alpha \in \Phi \rangle$, see [MS16, Lem. 3.2]. Let R_λ be the subgroup of the stabilizer $N_{\mathbf{G}^F}(\mathbf{T})_\lambda$ of λ in $N_{\mathbf{G}^F}(\mathbf{T})$ generated by \mathbf{T}^F and $\langle n_\alpha(-1) \mid \alpha \in \Phi_\lambda \rangle$, so that $R_\lambda / \mathbf{T}^F \cong R(\lambda)$. Here we use the notation of the Chevalley generators as in [GLS98, §1.12]. The next lemma concerns the restriction of $\Lambda(\lambda)$ to R_λ .

Lemma 3.8. *Assume that \mathbf{G} is simple of simply connected type, not of type A_n , and let $\lambda \in \text{Irr}(\mathbf{T}^F)$. Then the following hold:*

- (i) any extension $\widehat{\lambda}$ of λ to R_λ satisfies that $\widehat{\lambda}(x) \in \{\pm 1\}$ for all $x \in \langle n_\alpha(-1) \mid \alpha \in \Phi_\lambda \rangle$;
- (ii) if $\lambda^\sigma = \lambda$, then $\widehat{\lambda}^\sigma = \widehat{\lambda}$ for any extension $\widehat{\lambda}$ of λ to R_λ . In particular, $R(\lambda) \leq \text{Ker}(\delta_{\lambda, \sigma})$.

Proof. Let $\widehat{\lambda}$ be an extension to R_λ . By [MS16, Lem. 5.1], Φ_λ is comprised of $\alpha \in \Phi$ such that $\lambda(h_\alpha(t)) = 1$ for each $t \in \mathbb{F}_q^\times$. In particular, this means that if $\alpha \in \Phi_\lambda$, then $\lambda(h_\alpha(-1)) = 1$. But since $n_\alpha(-1)^2 = h_\alpha(-1)$, it follows that $\widehat{\lambda}(n_\alpha(-1)) \in \{\pm 1\}$, proving (i). For (ii), notice that

$$\widehat{\lambda}^\sigma(tx) = \widehat{\lambda}^\sigma(t)\widehat{\lambda}^\sigma(x) = \lambda^\sigma(t)\widehat{\lambda}(x) = \lambda(t)\widehat{\lambda}(x) = \widehat{\lambda}(tx),$$

for $tx \in R_\lambda$ with $t \in \mathbf{T}^F$ and $x \in \langle n_\alpha(-1) \mid \alpha \in \Phi_\lambda \rangle$, since $\widehat{\lambda}(x) \in \{\pm 1\}$ and $\lambda^\sigma = \lambda$. \blacksquare

Square Roots

3.9. Let p be an odd prime. In light of 3.6, we will want to study the action of \mathcal{G} on \sqrt{p} . Let us be more explicit about our choice of \sqrt{p} . We fix an injective group homomorphism $\jmath : \mathbb{Q}/\mathbb{Z} \rightarrow \overline{\mathbb{Q}}^\times$ and let $\tilde{\jmath} : \mathbb{Q} \rightarrow \overline{\mathbb{Q}}^\times$ be the composition with the natural quotient map $\mathbb{Q} \rightarrow \mathbb{Q}/\mathbb{Z}$. We have $i = \tilde{\jmath}(1/4)$ is an element such that $i^2 = -1$. We also denote this by $\sqrt{-1}$. Letting $\omega = (-1)^{\frac{p-1}{2}}$, there then exists a unique square root \sqrt{p} satisfying $\sqrt{\omega}\sqrt{p} = \sum_{n=1}^{p-1} \left(\frac{n}{p}\right) \zeta_p^n$, where $\left(\frac{n}{p}\right)$ denotes the Legendre symbol and $\zeta_p := \tilde{\jmath}\left(\frac{1}{p}\right)$, see [Bon06, §36]. We set $\sqrt{\omega p} := \sqrt{\omega}\sqrt{p}$.

Now, let $\sigma \in \mathcal{G}$ and let $k \in \mathbb{Z}$ be an integer coprime to p such that $\xi^\sigma = \xi^k$ for all p th roots of unity $\xi \in \overline{\mathbb{Q}}$. Then we have

$$\sqrt{\omega p}^\sigma = \sum_{n=1}^{p-1} \left(\frac{n}{p}\right) \zeta_p^{kn} = \left(\frac{k}{p}\right) \cdot \sum_{n=1}^{p-1} \left(\frac{kn}{p}\right) \zeta_p^{kn} = \left(\frac{k}{p}\right) \sqrt{\omega p}. \quad (3.10)$$

4. Galois Automorphisms Relevant for the McKay–Navarro Conjecture

4.1. In this section, we turn our attention to the case of the specific Galois automorphisms that appear in the context of the McKay–Navarro conjecture, with the aim of recording several statements that will also likely be useful for future work with the conjecture for finite reductive groups. Let ℓ be a prime and denote by $\mathcal{H} := \mathcal{H}_\ell$ the subgroup of elements $\sigma \in \mathcal{G}$ such that there is some integer $r \geq 0$ satisfying $\zeta^\sigma = \zeta^{\ell^r}$ for all roots of unity ζ of order coprime to ℓ . Throughout, we fix ℓ and $\sigma \in \mathcal{H}$ and understand r to be this integer corresponding to σ .

Assume $p \neq \ell$ is an odd prime. We now discuss the action of \mathcal{H} on \sqrt{p} in order to describe the character $\gamma_{\lambda, \sigma}$ in more detail in the case that $\sigma \in \mathcal{H}$. Note that $\left(\frac{\ell^r n}{p}\right) = \left(\frac{\ell}{p}\right)^r \left(\frac{n}{p}\right)$. Then (3.10) implies $\sqrt{\omega p}^\sigma = \left(\frac{\ell}{p}\right)^r \sqrt{\omega p}$.

The Case ℓ Odd

4.2. First, suppose that ℓ is odd. Note that if r is even, then $\ell^r \equiv 1 \pmod{4}$, so σ fixes 4th roots of unity and $\left(\frac{\ell}{p}\right)^r = 1$ since $\left(\frac{\ell}{p}\right) \in \{\pm 1\}$. If r is odd, then $\ell^r \equiv \ell \pmod{4}$. We also remark that when $\ell \neq p$ are odd primes, we have

$$\left(\frac{\ell}{p}\right) = (-1)^{(p-1)(\ell-1)/4} \left(\frac{p}{\ell}\right).$$

Together with (3.10) this yields the following:

Lemma 4.3. *Assume $\ell \neq p$ are both odd and $\sigma \in \mathcal{H}$. Let $\omega = (-1)^{\frac{p-1}{2}}$, so $p \equiv \omega \pmod{4}$.*

- (i) *If r is even, then $\sqrt{p}^\sigma = \sqrt{p}$ and $\sqrt{\omega p}^\sigma = \sqrt{\omega p}$.*
- (ii) *If r is odd, then $\sqrt{\omega p}^\sigma = \left(\frac{\ell}{p}\right) \sqrt{\omega p}$, so $\sqrt{p}^\sigma = (-1)^{(p-1)(\ell-1)/4} \left(\frac{\ell}{p}\right) \sqrt{p} = \left(\frac{p}{\ell}\right) \sqrt{p}$.*

We can now apply this to the setup of 3.1. In particular, we have \mathbf{G}^F is a finite reductive group defined over \mathbb{F}_q with q a power of p . Combining Lemmas 3.6 and 4.3, we have the following.

Lemma 4.4. *Assume $\ell \neq p$ are both odd and $(\mathbf{L}, \lambda) \in \text{Cusp}(\mathbf{G}, F)$ is a cuspidal pair. Let $\sigma \in \mathcal{H}$ and $w = w_1 w_2 \in W(\lambda)$ with $w_1 \in R(\lambda)$ and $w_2 \in C(\lambda)$.*

- (i) *If q is a square or r is even, then $\gamma_{\lambda, \sigma} = 1$.*
- (ii) *If q is not a square and r is odd, then*

$$\gamma_{\lambda, \sigma}(w) = \left(\frac{p}{\ell}\right)^{l(w_2)} = \begin{cases} (-1)^{l(w_2)} \left(\frac{\ell}{p}\right)^{l(w_2)} & \text{if } p \equiv \ell \equiv 3 \pmod{4} \\ \left(\frac{\ell}{p}\right)^{l(w_2)} & \text{otherwise,} \end{cases}$$

where $l(w_2)$ is the length of w_2 in the Weyl group (\mathbf{W}, \mathbf{S}) of \mathbf{G} .

4.5. The particular case that $\ell \mid (q - 1)$ has been studied in various contexts related to (refinements of) the McKay conjecture, see, e.g., [MS16; CSSF20], due to the relatively nice nature of height-zero characters in this case. For this reason, we remark that if ℓ is a prime dividing $(q - 1)$, we have $q \equiv 1 \pmod{\ell}$ is a square modulo ℓ , and hence either q is a square or p is a square modulo ℓ . Then Lemma 4.4 yields:

Corollary 4.6. *Assume ℓ and p are odd and $\ell \mid (q - 1)$. Then for any cuspidal pair $(\mathbf{L}, \lambda) \in \text{Cusp}(\mathbf{G}, F)$ and $\sigma \in \mathcal{H}$, the character $\gamma_{\lambda, \sigma}$ is trivial.*

The Case $\ell = 2$

4.7. We now focus on the situation when $\ell = 2$ and analyse the character $\gamma_{\lambda, \sigma}$ in this case. We begin by noting that

$$\left(\frac{2}{p}\right) = \begin{cases} 1 & \text{if } p \equiv \pm 1 \pmod{8}, \\ -1 & \text{if } p \equiv \pm 3 \pmod{8}, \end{cases}$$

so comparing with (3.10) yields:

Lemma 4.8. *Let $\sigma \in \mathcal{H}$ with $\ell = 2$ and let p be an odd prime. Let $\omega = (-1)^{\frac{p-1}{2}}$, so $p \equiv \omega \pmod{4}$.*

- (i) *If $p \equiv \pm 1 \pmod{8}$, then $\sqrt{\omega p}^\sigma = \sqrt{\omega p}$.*
- (ii) *If $p \equiv \pm 3 \pmod{8}$, then $\sqrt{\omega p}^\sigma = (-1)^r \sqrt{\omega p}$.*

In [SF19, Lem. 4.10], the character $\gamma_{\lambda, \sigma}$ is described explicitly in the case of a specific choice of σ . We now record an extension of that statement to all of \mathcal{H} .

Lemma 4.9. *Assume $\ell = 2$, p is odd, and $(\mathbf{L}, \lambda) \in \text{Cusp}(\mathbf{G}, F)$. Let $\sigma \in \mathcal{H}$ and $w = w_1 w_2 \in W(\lambda)$ with $w_1 \in R(\lambda)$ and $w_2 \in C(\lambda)$.*

- (i) *If $q \equiv 1 \pmod{8}$ then $\gamma_{\lambda, \sigma}(w) = 1$.*

- (ii) If $q \equiv -1 \pmod{8}$ then $\gamma_{\lambda,\sigma}(w) = \begin{cases} 1 & \text{if } i^\sigma = i, \\ (-1)^{\mathfrak{l}(w_2)} & \text{if } i^\sigma = -i. \end{cases}$
- (iii) If $q \equiv 3 \pmod{8}$ then $\gamma_{\lambda,\sigma}(w) = \begin{cases} 1 & \text{if } i^\sigma = (-1)^r i, \\ (-1)^{\mathfrak{l}(w_2)} & \text{if } i^\sigma = (-1)^{r-1} i. \end{cases}$
- (iv) If $q \equiv -3 \pmod{8}$ then $\gamma_{\lambda,\sigma}(w) = (-1)^{r \cdot \mathfrak{l}(w_2)}$.

Here $\mathfrak{l}(w_2)$ denotes the length of w_2 in the Weyl group (\mathbf{W}, \mathbf{S}) of \mathbf{G} .

Proof. This follows from Lemma 3.6 and Lemma 4.8. ■

4.10. We conclude with one further remark. When $\ell = 2$, the power r defined by $\sigma \in \mathcal{H}$ is even if and only if σ fixes third roots of unity. Hence in this case, the action of σ on \sqrt{p} and descriptions of $\gamma_{\lambda,\sigma}$ obtained in Lemmas 4.8 and 4.9 can alternatively be described entirely in terms of $p \pmod{8}$ and the action of σ on third and fourth roots of unity. The same will be true in Corollary 13.7 below, where we describe $\gamma_{\lambda,\sigma}\delta_{\lambda,\sigma}$ in the case of $\mathrm{Sp}(V)$.

5. Power Maps and Regular Unipotent Elements

5.1. In this section, we return to the setup of 3.1, so that \mathbf{G} is a connected reductive algebraic group. We fix an integer $k \in \mathbb{Z}$ coprime to p . Then the power map $g \mapsto g^k$ on \mathbf{G} induces a permutation of the set of unipotent conjugacy classes $\mathrm{Cl}_{\mathrm{uni}}(\mathbf{G})$. In fact, this map is known to be the identity. In other words, any unipotent element is *rational*. This was shown, for instance, by Tiep–Zalesski [TZ04, Thm. 4.3], with other proofs given by Lusztig [Lus09a, Prop. 2.5(a)] and Liebeck–Seitz [LS12, Cor. 3].

We now consider the analogous question over \mathbb{F}_q . Let $\mathcal{O} \in \mathrm{Cl}_{\mathrm{uni}}(\mathbf{G})^F$ be an F -stable unipotent class. Then the power map induces a permutation $\pi_k : \mathrm{Cl}_{\mathbf{G}^F}(\mathcal{O}^F) \rightarrow \mathrm{Cl}_{\mathbf{G}^F}(\mathcal{O}^F)$ of the set of \mathbf{G}^F -conjugacy classes contained in the fixed point set \mathcal{O}^F . If \mathbf{G} is simple and simply connected, then [TZ04, Thm. 1.7] describes exactly when every unipotent element of \mathbf{G}^F is rational, which means π_k is the identity for any class $\mathcal{O} \in \mathrm{Cl}_{\mathrm{uni}}(\mathbf{G})^F$.

Here we wish to describe the actual permutation. Unlike [TZ04], we will focus just on the case of good characteristic. Note that after [TZ04, Thm. 1.5], we know this permutation is a product of transpositions. In this section, we consider the case of regular unipotent elements. This will be used to settle the general case for symplectic and special orthogonal groups in Section 9. If $\mathbf{G} = \mathrm{SO}(V)$, then every element of \mathbf{G}^F is rational by [TZ04, Thm. 1.9], even strongly rational, so the main case of interest for us is when $\mathbf{G} = \mathrm{Sp}(V)$. However, our argument below also covers the case of $\mathrm{SO}(V)$.

For any $x \in \mathbf{G}^F$ we denote by $H^1(F, A_{\mathbf{G}}(x))$ the F -conjugacy classes of $A_{\mathbf{G}}(x)$. The F -class of $gC_{\mathbf{G}}^{\circ}(x) \in A_{\mathbf{G}}(x)$ is denoted by $[gC_{\mathbf{G}}^{\circ}(x)]$. If $\mathcal{L} : \mathbf{G} \rightarrow \mathbf{G}$ is the Lang map, defined by $\mathcal{L}(g) = g^{-1}F(g)$, then we have a bijection $\mathrm{Cl}_{\mathbf{G}^F}(\mathcal{O}^F) \rightarrow H^1(F, A_{\mathbf{G}}(x))$ given by ${}^9x \mapsto [\mathcal{L}(g)C_{\mathbf{G}}^{\circ}(x)]$, where \mathcal{O} is the \mathbf{G} -class of x . We leave the proof of the following to the reader.

Φ	$A_{2m+\epsilon}$	$B_{2m+\epsilon}$	C_n	$D_{2m+\epsilon}$	E_7
$\check{\rho}_\Delta$	$\epsilon\check{\omega}_m$	$(m+\epsilon)\check{\omega}_1$	$\check{\omega}_n$	$(m+\epsilon)\check{\omega}_1$	$\check{\omega}_7$

Table 1: Representative of the image of $\check{\rho}_\Delta$ in the fundamental group $\mathbb{Z}\check{\Omega}/\mathbb{Z}\check{\Phi}$.

Lemma 5.2. *Assume $\mathcal{O} \in \text{Cl}_{\text{uni}}(\mathbf{G})^F$ is an F -stable unipotent class and $\mathbf{u} \in \mathcal{O}^F$. If $g \in \mathbf{G}$ is such that $\mathbf{u}^k = {}^g\mathbf{u}$, then we have a commutative diagram*

$$\begin{array}{ccc} H^1(F, \mathbf{A}_{\mathbf{G}}(\mathbf{u})) & \longrightarrow & \text{Cl}_{\mathbf{G}^F}(\mathcal{O}^F) \\ \downarrow \tau & & \downarrow \pi_k \\ H^1(F, \mathbf{A}_{\mathbf{G}}(\mathbf{u})) & \longrightarrow & \text{Cl}_{\mathbf{G}^F}(\mathcal{O}^F) \end{array}$$

where $\tau([\mathbf{a}C_{\mathbf{G}}^\circ(\mathbf{u})]) = [g^{-1}\mathbf{a}F(g)C_{\mathbf{G}}^\circ(\mathbf{u})]$. Moreover, if there is an F -stable torus $\mathbf{S} \leq \mathbf{G}$ for which the natural map $\mathbf{S} \rightarrow \mathbf{A}_{\mathbf{G}}(\mathbf{u})$ is surjective, then $\tau([\mathbf{a}C_{\mathbf{G}}^\circ(\mathbf{u})]) = [\mathbf{a}\mathcal{L}(g)C_{\mathbf{G}}^\circ(\mathbf{u})]$.

5.3. Let $X = X(\mathbf{T})$ and $\check{X} = \check{X}(\mathbf{T})$ be the character and cocharacter groups of \mathbf{T} with the usual perfect pairing $\langle -, - \rangle : X \times \check{X} \rightarrow \mathbb{Z}$. If $\mathbb{Z}_{(p)}$ is the localisation of \mathbb{Z} at the prime ideal (p) , then we may choose an isomorphism $\iota : \mathbb{Z}_{(p)}/\mathbb{Z} \rightarrow \mathbb{F}^\times$. With respect to this choice, we have a surjective homomorphism of abelian groups $\tilde{\iota} : \mathbb{Q}\check{X} \rightarrow \mathbf{T}$, as in [Bon05, §3], where $\mathbb{Q}\check{X} = \mathbb{Q} \otimes_{\mathbb{Z}} \check{X}$ is a \mathbb{Q} -vector space. If $\mathbb{Q}X = \mathbb{Q} \otimes_{\mathbb{Z}} X$ then $\langle -, - \rangle$ extends naturally to a non-degenerate bilinear form $\mathbb{Q}X \times \mathbb{Q}\check{X} \rightarrow \mathbb{Q}$.

For each root $\alpha \in \Phi$ we have a corresponding coroot $\check{\alpha} \in \check{X}$. Let $\mathbb{Q}\check{\Phi} \subseteq \mathbb{Q}\check{X}$ be the subspace spanned by $\check{\Phi} = \{\check{\alpha} \mid \alpha \in \Phi\}$. Then we have a set of fundamental dominant coweights $\check{\Omega} = \{\check{\omega}_\alpha \mid \alpha \in \Delta\} \subseteq \mathbb{Q}\check{\Phi}$ defined such that $\langle \alpha, \check{\omega}_\beta \rangle = \delta_{\alpha,\beta}$. By [Bou02, §1, no. 10] we have a vector

$$\check{\rho}_\Delta := \sum_{\alpha \in \Delta} \check{\omega}_\alpha = \frac{1}{2} \sum_{\alpha \in \Phi^+} \check{\alpha} \in \mathbb{Z}\check{\Omega} \subseteq \mathbb{Q}\check{\Phi}.$$

In Table 1, we describe the image of $\check{\rho}_\Delta$ in the fundamental group $\check{\Pi} := \mathbb{Z}\check{\Omega}/\mathbb{Z}\check{\Phi}$ when Φ is irreducible. Here we use the notation in the plates of [Bou02]. For the cases G_2, F_4, E_6 , and, E_8 we have $\check{\rho}_\Delta$ is always contained in $\mathbb{Z}\check{\Phi}$.

Proposition 5.4. *Assume p is good for \mathbf{G} and $\mathbf{u} \in \mathbf{B}^F$ is a regular unipotent element with \mathcal{O} the \mathbf{G} -class containing \mathbf{u} . If $r \in \mathbb{Z}$ is such that $\iota(r/(q-1) + \mathbb{Z}) = k \pmod{p}$, then there is an element $g \in \mathbf{B}$ such that ${}^g\mathbf{u} = \mathbf{u}^k$ and $\mathcal{L}(g)C_{\mathbf{G}}^\circ(\mathbf{u}) = zC_{\mathbf{G}}^\circ(\mathbf{u})$, where $z := \tilde{\iota}(r\check{\rho}_\Delta)$ is contained in $Z(\mathbf{G})$. If τ is as in Lemma 5.2, then $\tau([\mathbf{a}C_{\mathbf{G}}^\circ(\mathbf{u})]) = [\mathbf{a}zC_{\mathbf{G}}^\circ(\mathbf{u})]$.*

Proof. Let $\mathbf{U} \leq \mathbf{B}$ be the unipotent radical of \mathbf{B} . For each $\alpha \in \Delta$ we pick a closed embedding $x_\alpha : \mathbb{F} \rightarrow \mathbf{U}$ onto the corresponding root subgroup. Denote by $\mathbf{U}_{\text{der}} \triangleleft \mathbf{U}$ the derived subgroup of \mathbf{U} and let $\bar{\mathbf{u}} = \prod_{\alpha \in \Delta} x_\alpha(c_\alpha)\mathbf{U}_{\text{der}}$, with $c_\alpha \in \mathbb{F}$, be the image of \mathbf{u} in $\mathbf{U}/\mathbf{U}_{\text{der}}$. Clearly $\bar{\mathbf{u}}^k = \prod_{\alpha \in \Delta} x_\alpha(kc_\alpha)\mathbf{U}_{\text{der}}$. If $t = \tilde{\iota}(r/(q-1) \cdot \check{\rho}_\Delta)$, then we have $\alpha(t) = k$ for all $\alpha \in \Delta$. Note that $\check{\rho}_\Delta$ is invariant under all graph automorphisms and

$$\mathcal{L}(t) = \tilde{\iota} \left(\frac{(q-1)r}{(q-1)} \cdot \check{\rho}_\Delta \right) = \tilde{\iota}(r\check{\rho}_\Delta) = z.$$

Moreover, $z \in Z(\mathbf{G})$ because $\alpha(z) = \iota(\langle \alpha, r\check{\rho}_\Delta \rangle + \mathbb{Z}) = 1$ for any root $\alpha \in \Phi$, as $\langle \alpha, r\check{\rho}_\Delta \rangle = r\langle \alpha, \check{\rho}_\Delta \rangle \in \mathbb{Z}$.

It is clear that ${}^t\bar{u} = \bar{u}^k$, so $t^{-1}(u^k)tu^{-1} \in \mathbf{U}_{\text{der}}$. As u is regular, $t^{-1}(u^k)tu^{-1} = ({}^v u)u^{-1}$ for some $v \in \mathbf{U}$, see the proof of [DM20, Prop. 12.2.2], so ${}^t v u = u^k$. As $\mathcal{L}(t) \in Z(\mathbf{G})$, we have $\mathcal{L}(tv) = \mathcal{L}(v)\mathcal{L}(t) \in C_{\mathbf{G}}(u)$, which implies $\mathcal{L}(v) \in C_{\mathbf{U}}(u) \leq C_{\mathbf{G}}^\circ(u)$ by [DM20, Prop. 12.2.7]. Hence $C_{\mathbf{G}}^\circ(u)\mathcal{L}(tv) = C_{\mathbf{G}}^\circ(u)z$. \blacksquare

5.5. Let us maintain the assumptions of Proposition 5.4. As was already used above, the natural map $Z(\mathbf{G}) \rightarrow C_{\mathbf{G}}(u)$ factors through an F -equivariant isomorphism $\mathcal{Z}(\mathbf{G}) \rightarrow A_{\mathbf{G}}(u)$, where $\mathcal{Z}(\mathbf{G}) = Z(\mathbf{G})/Z^\circ(\mathbf{G})$. This defines a bijection $H^1(F, \mathcal{Z}(\mathbf{G})) \rightarrow H^1(F, A_{\mathbf{G}}(u))$. As $\mathcal{Z}(\mathbf{G})$ is abelian we have $H^1(F, \mathcal{Z}(\mathbf{G})) = \mathcal{Z}(\mathbf{G})/\mathcal{L}(\mathcal{Z}(\mathbf{G}))$. Hence, we have u and u^k are \mathbf{G}^F -conjugate if and only if $z \in \mathcal{L}(\mathcal{Z}(\mathbf{G}))$, where $z = \tilde{\iota}(r\check{\rho}_\Delta)$ is as in Proposition 5.4.

Recall that $Q\check{X} = Q\check{\Phi} \oplus Q\check{\Phi}^\perp$ where $\check{\Phi}^\perp = \{y \in \check{X} \mid \langle x, y \rangle = 0 \text{ for all } x \in \check{\Phi}\}$. We thus have a natural projection map $Q\check{X} \rightarrow Q\check{\Phi}$ and we denote by $\mathcal{Z}\check{\Phi} \subseteq \check{X}_{\text{ss}} \subseteq \mathcal{Z}\check{\Omega}$ the image of \check{X} . If \mathbf{G} is semisimple then $\check{X}_{\text{ss}} = \check{X}$. The map $\tilde{\iota}$ defines an isomorphism $\text{Tor}_{p'}(\mathcal{Z}\check{\Omega}/\check{X}_{\text{ss}}) \rightarrow \mathcal{Z}(\mathbf{G})$ where $\text{Tor}_{p'}$ denotes the p' -torsion subgroup. As $2\check{\rho}_\Delta \in \mathcal{Z}\check{\Phi}$ we see that $z^2 = 1$ and we have $z = 1$ if and only if at least one of the following holds: $p = 2$, $\check{\rho}_\Delta \in \check{X}_{\text{ss}}$, $\log_p(q)$ is even, or $k \pmod{p}$ is a square in \mathbb{F}_p .

Clearly $z = 1$ if either of the first two conditions hold. So assume neither of these two conditions hold. Then $z = 1$ if and only if $2 \mid r$. As $k \pmod{p} \in \mathbb{F}_p$, we have $\tilde{\iota}(s/(p-1)) = k \pmod{p}$ for some $1 \leq s < p$. If $a = \log_p(q)$ then

$$r = s \cdot \frac{q-1}{p-1} = s(p^{a-1} + \cdots + p + 1).$$

Hence, as $p \neq 2$ we have $2 \mid r$ if and only if either a is even or $2 \mid s$. This last condition is equivalent to $k \pmod{p} \in \mathbb{F}_p$ being a square in \mathbb{F}_p .

For an example, suppose \mathbf{G} is a simple group of type A_{n-1} then $\check{\Pi}$ is a cyclic group of order n and $\check{\Pi}_{\mathbf{G}} := \mathcal{Z}\check{\Omega}/\check{X}_{\text{ss}}$ is a cyclic group of order $m \mid n$, which we view additively. The Lang map on $\mathcal{Z}(\mathbf{G}) \cong Z(\mathbf{G})$ is identified with the map $\mathcal{L} : \check{\Pi}_{\mathbf{G}} \rightarrow \check{\Pi}_{\mathbf{G}}$ given by $\mathcal{L}(c) = (q-1)c$ if F is split and $\mathcal{L}(c) = -(q+1)c$ if F is twisted. Identifying $\mathcal{L}(\check{\Pi}_{\mathbf{G}})$ with $\check{\Pi}_{\mathbf{G}}/\text{Ker}(\mathcal{L})$ we see easily that $|\mathcal{L}(\check{\Pi}_{\mathbf{G}})| = m/\text{gcd}(m, q \pm 1)$. Hence, if $z \neq 1$, in which case $2 \mid m$ and $p \neq 2$, we have u is \mathbf{G}^F -conjugate to u^k if and only if $|\mathcal{L}(\check{\Pi}_{\mathbf{G}})|$ is even.

Given this discussion and the data in Table 1 the other simple groups are easily dealt with in this way. This seems to simplify the discussion in [TZ04, §5] and the proof of [TZ04, Thm. 1.7] in the case of good characteristic, especially in the case of twisted groups. We will need the following whose proof is easy and left to the reader. We simply note that when $\mathbf{G} = \text{SO}(V)$ we have $\check{\omega}_1 \in \check{X}_{\text{ss}} = \check{X}$.

Lemma 5.6. *Assume p is good for \mathbf{G} and $u \in \mathbf{G}^F$ is a regular unipotent element. If $\mathbf{G} = \text{GL}(V)$ or $\text{SO}(V)$, then u is rational in \mathbf{G}^F . If $\mathbf{G} = \text{Sp}(V)$, then u and u^k are \mathbf{G}^F -conjugate if and only if $k \pmod{p} \in \mathbb{F}_q$ is a square.*

6. A Variation of GGCs for Disconnected Groups

6.1. In this section, we assume \mathbf{G} is a (possibly disconnected) reductive algebraic group with Frobenius endomorphism $F : \mathbf{G} \rightarrow \mathbf{G}$. If $u \in \mathbf{G}^{\circ F}$ is a unipotent element, then one has a corresponding GGC $\Gamma_u = \Gamma_u^{\mathbf{G}^{\circ F}}$ of the finite group $\mathbf{G}^{\circ F}$. We would like to construct such characters for the group \mathbf{G}^F . One option is the *induced* GGC $\text{Ind}_{\mathbf{G}^{\circ F}}^{\mathbf{G}^F}(\Gamma_u)$. However, we would like to consider a different construction, which instead involves extending Γ_u and then inducing to \mathbf{G}^F .

If $r = p^a$ with $a \geq 0$, then we denote by $F_r : \mathbb{F} \rightarrow \mathbb{F}$ the map defined by $F_r(k) = k^r$. Let $\text{Frob}_1(\mathbf{G})$ denote the automorphism group of \mathbf{G} as an algebraic group. If $r > 1$, then let $\text{Frob}_r(\mathbf{G})$ denote the set of Frobenius endomorphisms on \mathbf{G} endowing \mathbf{G} with an \mathbb{F}_r -structure. We let $\text{Frob}(\mathbf{G}) = \bigcup_{a \geq 0} \text{Frob}_{p^a}(\mathbf{G})$. Moreover, $\text{Frob}(\mathbf{G}, F) \subseteq \text{Frob}(\mathbf{G})$ denotes those $\sigma \in \text{Frob}(\mathbf{G})$ commuting with F , and $\text{Frob}_r(\mathbf{G}, F) = \text{Frob}(\mathbf{G}, F) \cap \text{Frob}_r(\mathbf{G})$. If $\sigma \in \text{Frob}(\mathbf{G})$, then we have a corresponding F_r -semilinear endomorphism $\sigma : \mathfrak{g} \rightarrow \mathfrak{g}$ of the Lie algebra \mathfrak{g} of \mathbf{G}° , which we also denote by σ , see [Tay18, 11.2, 11.3].

6.2. In what follows, we will freely use the notation and terminology of [BDT19] particularly that of [BDT19, §6]. In particular, we assume throughout that \mathbf{G}° is proximate and that $\mathcal{K} = (\phi_{\text{spr}}, \kappa, \chi_p)$ is a Kawanaka datum with respect to which our GGCs will be defined. After the discussion in [Tay18, §11], we may, and will, assume that for any $\sigma \in \text{Frob}_r(\mathbf{G})$ with $r = p^a \geq 1$, the following hold: $\phi_{\text{spr}} \circ \sigma = \sigma \circ \phi_{\text{spr}}$, and $\kappa(\sigma(X), \sigma(Y)) = F_r(\kappa(X, Y))$.

We fix a unipotent element $u \in \mathbf{G}^{\circ F}$ and a corresponding Dynkin cocharacter $\lambda \in \mathcal{D}_u(\mathbf{G}^\circ)^F$. Let $\mathbf{L} = C_{\mathbf{G}}(\lambda)$ be the centraliser of the cocharacter. Then $\mathbf{L}^\circ = C_{\mathbf{G}^\circ}(\lambda)$ is a Levi subgroup of \mathbf{G}° . If $\mathcal{U}(\mathbf{G}^\circ) \subseteq \mathbf{G}^\circ$ is the closed set of unipotent elements, then we have a function $\eta_u : \mathcal{U}(\mathbf{G}^\circ)^F \rightarrow \overline{\mathbb{Q}}_\ell$ whose restriction to $\mathbf{U}(\lambda, -2)^F$, denoted by $\eta_{u, \lambda}$, is a linear character. This character satisfies ${}^x \eta_{u, \lambda} = \eta_{x u, x \lambda}$ for all $x \in \mathbf{G}^F$. We have a corresponding character $\zeta_{u, \lambda} \in \text{Irr}(\mathbf{U}(\lambda, -1)^F)$ defined uniquely by the condition that

$$\text{Ind}_{\mathbf{U}(\lambda, -2)^F}^{\mathbf{U}(\lambda, -1)^F}(\eta_{u, \lambda}) = q^{\dim \mathfrak{g}(\lambda, -1)/2} \zeta_{u, \lambda}.$$

We then have $\Gamma_u = \text{Ind}_{\mathbf{U}(\lambda, -1)^F}^{\mathbf{G}^{\circ F}}(\zeta_{u, \lambda})$.

6.3. Recall that $\ell \neq p$ is a prime. We denote by $C_{\mathbf{G}}(\lambda)_{\zeta_{u, \lambda}}^F$ the stabiliser of $\zeta_{u, \lambda}$ in $C_{\mathbf{G}}(\lambda)^F$. We will assume that $S_{u, \lambda} \leq C_{\mathbf{G}}(\lambda)_{\zeta_{u, \lambda}}^F$ is a fixed Sylow ℓ -subgroup. The group $C_{\mathbf{G}}(\lambda)$ normalises $\mathbf{U}(\lambda, -1)$, so $\mathbf{U}(\lambda, -1)^F \cdot S_{u, \lambda}$ is a group. In fact, this group is a semidirect product, since $\mathbf{U}(\lambda, -1)^F$ is a p -group. Moreover, $\zeta_{u, \lambda}$ has a unique extension $\widehat{\zeta}_{u, \lambda} \in \text{Irr}(\mathbf{U}(\lambda, -1)^F S_{u, \lambda})$ such that $o(\widehat{\zeta}_{u, \lambda}) = o(\zeta_{u, \lambda})$ by [Isa18, Thm. 1.5]. Here, if G is a finite group and $\chi \in \text{Irr}(G)$ is an irreducible character, then $o(\chi)$ denotes the order of the corresponding determinant character. We now define a character

$$\widehat{\Gamma}_{u, \ell} = \text{Ind}_{\mathbf{U}(\lambda, -1)^F S_{u, \lambda}}^{\mathbf{G}^F}(\widehat{\zeta}_{u, \lambda}).$$

Note that in the special case $\mathbf{G} = \mathbf{G}^\circ$, we have $\widehat{\Gamma}_u$ is a summand of Γ_u , which is part of the ideas considered in [BDT19]. More generally, we have the following.

Lemma 6.4. *For any unipotent element $u \in \mathbf{G}^{\circ F}$ we have*

$$\text{Res}_{\mathbf{G}^{\circ F}}^{\mathbf{G}^F}(\widehat{\Gamma}_{u,\ell}) = \sum_{g \in \mathbf{G}^F / \mathbf{G}^{\circ F} S_{u,\lambda}} \Gamma_{gug^{-1}}.$$

Moreover, if $C_{\mathbf{G}}(u)_{\ell}^F \leq C_{\mathbf{G}}(u)^F$ is a Sylow ℓ -subgroup, then $C_{\mathbf{G}}(u)_{\ell}^F \leq \mathbf{G}^{\circ F} S_{u,\lambda}$.

Proof. The first statement follows from the usual Mackey formula for finite groups, together with [Tay18, Prop. 11.10]. If $x \in C_{\mathbf{G}}(u)^F$ then ${}^x\lambda \in \mathcal{D}_u(\mathbf{G}^{\circ})^F$. By [BDT19, Lem. 3.6], and a standard application of the Lang–Steinberg Theorem, there exists $h \in C_{\mathbf{G}^{\circ}}(u)^F$ such that $\lambda = h^x\lambda$. Therefore, if $L = C_{\mathbf{G}}(\lambda)$ then $g = hx \in C_L(u)^F \leq L_{\zeta_{u,\lambda}}^F$ which implies $C_{\mathbf{G}}(u)^F \leq C_{\mathbf{G}^{\circ}}(u)^F \cdot L_{\zeta_{u,\lambda}}^F$. Using conjugacy of Sylow subgroups in the quotient $C_{\mathbf{G}}(u)^F / C_{\mathbf{G}^{\circ}}(u)^F$ gives the second statement. \blacksquare

Remark 6.5. We will be interested in applying this construction in the special case where $\mathbf{G} = \text{GO}(V)$ is an orthogonal group and $p \neq 2$ is odd. In this case, $\mathbf{G}/\mathbf{G}^{\circ}$ has order 2. If the centraliser $C_{\mathbf{G}}(u)$ of $u \in \mathbf{G}^{\circ F}$ is not contained in \mathbf{G}° then $\mathbf{G}^F = \mathbf{G}^{\circ F} C_{\mathbf{G}}(u)_2^F$ and Lemma 6.4 shows that $\widehat{\Gamma}_{u,2}$ is an extension of the GGGC Γ_u .

The following extends [Tay18, Prop. 11.10] to give the action of the automorphism group on these characters.

Proposition 6.6. *Recall our assumption that \mathbf{G}° is proximate, p is good for \mathbf{G} , and $\ell \neq p$ is a prime. For any unipotent element $u \in \mathbf{G}^{\circ F}$ and $\sigma \in \text{Frob}(\mathbf{G}, F)$, we have ${}^{\sigma}\widehat{\Gamma}_{u,\ell} = \widehat{\Gamma}_{\sigma(u),\ell}$.*

Proof. Recall that the restriction of σ to \mathbf{G}^F is an automorphism. It is easy to see that $\sigma(\mathbf{U}(\lambda, -1)) = \mathbf{U}(\sigma \cdot \lambda, -1)$, where $\sigma \cdot \lambda$ is as in [Tay18, 11.6], and we get that ${}^{\sigma}\zeta_{u,\lambda} = \zeta_{\sigma(u),\sigma \cdot \lambda}$ by arguing as in [Tay18, Prop. 11.10]. Moreover, $\sigma(C_{\mathbf{G}}(\lambda)_{\zeta_{u,\lambda}}^F) = C_{\mathbf{G}}(\sigma \cdot \lambda)_{\zeta_{\sigma(u),\sigma \cdot \lambda}}^F$ and there exists an element $g \in C_{\mathbf{G}}(\sigma \cdot \lambda)_{\zeta_{\sigma(u),\sigma \cdot \lambda}}^F$ such that ${}^g\sigma(S_{u,\lambda}) = S_{\sigma(u),\sigma \cdot \lambda}$. Now certainly

$${}^{\sigma}\widehat{\Gamma}_{u,\ell} = {}^{g\sigma}\widehat{\Gamma}_{u,\ell} = \text{Ind}_{\mathbf{U}(\sigma \cdot \lambda, -1)^F S_{\sigma(u),\sigma \cdot \lambda}}^{\mathbf{G}^F} ({}^{g\sigma}\widehat{\zeta}_{u,\lambda})$$

and ${}^{g\sigma}\widehat{\zeta}_{u,\lambda}$ is an extension of $\zeta_{\sigma(u),\sigma \cdot \lambda}$. By the unicity of the extension, ${}^{g\sigma}\widehat{\zeta}_{u,\lambda} = \widehat{\zeta}_{\sigma(u),\sigma \cdot \lambda}$ and the result follows. \blacksquare

We now want to extend [SFT18, Prop. 4.10], which gives the action of \mathcal{G} on GGGCs. Understanding the effect of \mathcal{G} on $\widehat{\Gamma}_u$ seems to be more complicated and appears to be related to stronger notions of conjugacy, such as being strongly real. The following, though weaker than the result in [SFT18], will be sufficient for our purposes.

Proposition 6.7. *Recall our assumption that \mathbf{G}° is proximate, p is good for \mathbf{G} , and $\ell \neq p$ is a prime. Assume $\sigma \in \mathcal{G}$ is a Galois automorphism and $k \in \mathbb{Z}$ is an integer coprime to p such that $\xi^{\sigma} = \xi^k$ for all p th roots of unity $\xi \in \overline{\mathbb{Q}}_{\ell}^{\times}$. If $u \in \mathbf{G}^{\circ F}$ is a unipotent element and u^k is $\mathbf{G}^{\circ F}$ -conjugate to u , then $\widehat{\Gamma}_{u,\ell}^{\sigma} = \widehat{\Gamma}_{u,\ell}$.*

Proof. From the proof of [SFT18, Prop. 4.10], there exists an element $x \in \mathbf{L}^{\circ}$ such that $\phi_{\text{spr}}({}^x u) = k\phi_{\text{spr}}(u) \in \mathfrak{g}(\lambda, 2)_{\text{reg}}$ and ${}^x u$ is $\mathbf{G}^{\circ F}$ -conjugate to u^k . By assumption, ${}^{g^x} u = u$

for some $g \in \mathbf{G}^{\circ\mathbb{F}}$. The natural map $C_{\mathbf{L}^\circ}(\mathfrak{u})/C_{\mathbf{L}^\circ}^\circ(\mathfrak{u}) \rightarrow C_{\mathbf{G}^\circ}(\mathfrak{u})/C_{\mathbf{G}^\circ}^\circ(\mathfrak{u})$ is an isomorphism. Hence, a quick application of the Lang–Steinberg Theorem in the group $C_{\mathbf{L}^\circ}^\circ(\mathfrak{u})$ implies that, after possibly replacing x by $x\mathfrak{a}$ with $\mathfrak{a} \in C_{\mathbf{L}^\circ}(\mathfrak{u})$, we may assume that $x \in \mathbf{L}^{\circ\mathbb{F}}$. By Proposition 6.6, we have $\widehat{\Gamma}_{x\mathfrak{u},\ell}^\sigma = \widehat{\Gamma}_{\mathfrak{u}^k,\ell}^\sigma$ so it suffices to show that $\widehat{\Gamma}_{\mathfrak{u},\ell}^\sigma = \widehat{\Gamma}_{x\mathfrak{u},\ell}^\sigma$.

Arguing as in [SFT18, Prop. 4.10], we see that $\eta_{\mathfrak{u}}^\sigma = \eta_{x\mathfrak{u}} = {}^x\eta_{\mathfrak{u}}$. It follows that $\zeta_{\mathfrak{u},\lambda}^\sigma = \zeta_{x\mathfrak{u},\lambda} = {}^x\zeta_{\mathfrak{u},\lambda}$. Being a Galois conjugate, the character $\zeta_{\mathfrak{u},\lambda}^\sigma = {}^x\zeta_{\mathfrak{u},\lambda}$ has the same stabiliser as $\zeta_{\mathfrak{u},\lambda}$ in $C_{\mathbf{G}}(\lambda)^{\mathbb{F}}$. This implies x normalises the stabiliser of $\zeta_{\mathfrak{u},\lambda}$ in $C_{\mathbf{G}}(\lambda)^{\mathbb{F}}$, so $S_{\mathfrak{u},\lambda}^{xg} = S_{\mathfrak{u},\lambda}$ for some $g \in C_{\mathbf{G}}(\lambda)^{\mathbb{F}}_{\zeta_{\mathfrak{u},\lambda}}$. By the unicity of the extension, we have $\widehat{\zeta}_{\mathfrak{u},\lambda}^{\sigma xg} = \widehat{\zeta}_{\mathfrak{u},\lambda}$ so

$$\widehat{\Gamma}_{\mathfrak{u},\ell}^\sigma = \widehat{\Gamma}_{\mathfrak{u},\ell}^{\sigma xg} = \text{Ind}_{\mathbf{U}(\lambda,-1)^{\mathbb{F}} \cdot S_{\mathfrak{u},\lambda}^{xg}}^{\mathbf{G}^{\mathbb{F}}}(\widehat{\zeta}_{\mathfrak{u},\lambda}^{\sigma xg}) = \widehat{\Gamma}_{\mathfrak{u},\ell}^\sigma. \quad \blacksquare$$

7. Classical Groups

From now on, we assume that $p \neq 2$.

7.1. We fix a finite dimensional \mathbb{F} -vector space V , with $\dim(V) > 1$, and let $\mathcal{B} : V \times V \rightarrow \mathbb{F}$ be a non-degenerate symmetric or alternating bilinear form. We denote by $\text{GL}(V \mid \mathcal{B})$, resp., $\text{SL}(V \mid \mathcal{B})$, the subgroup of isometries of \mathcal{B} in the general linear group, resp., special linear group. If \mathcal{B} is symmetric, then $\text{GO}(V) := \text{GL}(V \mid \mathcal{B})$ is the orthogonal group and $\text{SO}(V) := \text{SL}(V \mid \mathcal{B}) = \text{GO}(V)^\circ$ is the special orthogonal group. If \mathcal{B} is alternating, then $\text{Sp}(V) := \text{GL}(V \mid \mathcal{B}) = \text{SL}(V \mid \mathcal{B})$ is the symplectic group.

For calculations, it will be convenient to make concrete choices for the form \mathcal{B} . We define a totally ordered set

$$\mathcal{J} = \begin{cases} \{1 \prec \dots \prec n \prec 0 \prec -n \prec \dots \prec -1\} & \text{if } \dim(V) = 2n + 1 \text{ is odd} \\ \{1 \prec \dots \prec n \prec -n \prec \dots \prec -1\} & \text{if } \dim(V) = 2n \text{ is even} \end{cases} \quad (7.2)$$

and choose a corresponding basis $\mathcal{V} = (v_i \mid i \in \mathcal{J}) \subset V$ ordered by \prec . For any integer $m \in \mathbb{Z}$, let $\text{sgn}(m) = 1$ if $m \geq 0$ and $\text{sgn}(m) = -1$ if $m < 0$. Let $\epsilon \in \{0, 1\}$ be such that $\mathcal{B}(v, w) = (-1)^\epsilon \mathcal{B}(w, v)$ for all $v, w \in V$. Then we assume that $\mathcal{B}(v_i, v_j) = \text{sgn}(i)^\epsilon \cdot \delta_{i,-j}$ for all $i, j \in \mathcal{J}$

Tori and Normalisers

7.3. Let $\mathbf{G} = \text{GL}(V \mid \mathcal{B})$. Via our choice of basis \mathcal{V} , we will interchangeably describe elements of $\text{GL}(V)$ either as linear maps or matrices. We will denote by $\mathbf{T} \leq \mathbf{B} \leq \mathbf{G}^\circ$ the subgroups of diagonal and upper triangular matrices. For $i \in \mathcal{J}$ we let $\check{\xi}_i : \mathbb{F}^\times \rightarrow \text{GL}(V)$ be the homomorphism defined such that $\check{\xi}_i(\zeta)v_j = \zeta^{\delta_{i,j}}v_j$ for any $\zeta \in \mathbb{F}^\times$ and $j \in \mathcal{J}$. We have $\mathbf{T} = \mathbf{T}_1 \times \dots \times \mathbf{T}_n$, where \mathbf{T}_i is the image of $\mathbf{d}_i : \mathbb{F}^\times \rightarrow \mathbf{T}$ defined by setting $\mathbf{d}_i(\zeta) = \check{\xi}_i(\zeta)\check{\xi}_{-i}(\zeta^{-1})$.

Denote by $\mathfrak{S}_{\mathcal{J}}$ the symmetric group on \mathcal{J} . Let $W_n \leq \mathfrak{S}_{\mathcal{J}}$ be the subgroup generated by $\mathbf{S} = \{s_1, \dots, s_{n-1}, s_n\}$, where $s_i = (i, i+1)(-i, -i-1)$ for $1 \leq i \leq n-1$ and $s_n = (n, -n)$. The pair (W_n, \mathbf{S}) is a Coxeter system of type B_n . We define elements:

- $t_m = (m, -m) = s_m \cdots s_{n-1} s_n s_{n-1} \cdots s_m$ if $1 \leq m \leq n$,
- $u_m = (m, -m)(n, -n) = s_m \cdots s_{n-2} (s_n s_{n-1} s_n s_{n-1}) s_{n-2} \cdots s_m$ if $1 \leq m < n$.

The subgroup $W'_n \leq W_n$ generated by $S' = \{s_1, \dots, s_{n-1}, s_n s_{n-1} s_n\}$ is a Coxeter group of type D_n . We have a further subgroup $W''_n \leq W'_n$ of type B_{n-1} , generated by $\{s_1, \dots, s_{n-2}, u_{n-1}\}$. When $n = 1$, we have $W''_n = W'_n$ is trivial. Let $\tilde{S} = S \cup \{s_0\}$, where $s_0 = t_1$. Assume $0 \leq a \leq n$ and let $b = n - a$. Then we have a reflection subgroup $W_a \times W_b \leq W_n$ of type $B_a \times B_b$ with Coxeter generators $\tilde{S} \setminus \{s_a\}$.

7.4. Recall that for any $\sigma \in \mathfrak{S}_J$, we have a corresponding transformation $p_\sigma \in GL(V)$ such that $p_\sigma(v_i) = v_{\sigma(i)}$ for all $i \in J$. The groups $\mathbf{W} = N_G(\mathbf{T})/\mathbf{T}$ and $\mathbf{W}^\circ = N_{G^\circ}(\mathbf{T})/\mathbf{T}$ denote the Weyl groups of \mathbf{G} and \mathbf{G}° . For each $s \in S$, we define an element $n_s \in N_G(\mathbf{T})$ as follows. We let $n_{s_i} := p_{s_i}$ for any $1 \leq i < n$ and

$$n_{s_n} := \begin{cases} p_{s_n} \check{\epsilon}_n(-1) & \text{if } \mathbf{G} = \mathrm{Sp}(V), \\ p_{s_n} \check{\epsilon}_0(-1) & \text{if } \mathbf{G} = \mathrm{GO}(V) \text{ and } \dim(V) \text{ is odd} \\ p_{s_n} & \text{if } \mathbf{G} = \mathrm{GO}(V) \text{ and } \dim(V) \text{ is even.} \end{cases}$$

The map defined by $n_s \mathbf{T} \mapsto s$, for $s \in S$, extends to a unique isomorphism $\mathbf{W} \rightarrow W_n$. We will implicitly identify \mathbf{W} and W_n in this way.

By Matsumoto's Theorem [GP00, Thm. 1.2.2], we obtain a unique element $n_w = n_{s_1} \cdots n_{s_k} \in N_G(\mathbf{T})$, where $w = s_1 \cdots s_k$, with $s_i \in S$, is a reduced expression for w . For $\mathbf{G} = \mathrm{GO}(V)$, the map $w \mapsto n_w$ is a homomorphism $\mathbf{W} \rightarrow N_G(\mathbf{T})$, i.e., the Weyl group lifts. When $\dim(V)$ is even and $\mathbf{G} = \mathrm{GO}(V)$, the group \mathbf{W}° is mapped onto W'_n .

Subspace Subgroups and Conformal Groups

7.5. Assume $0 \leq m \leq n$ is an integer and let $\mathcal{J}_1 = \{1, \dots, m, -m, \dots, -1\}$ and $\mathcal{J}_0 = \mathcal{J} \setminus \mathcal{J}_1$. We have corresponding subspaces $V_i \subseteq V$ spanned by $\{v_k \mid k \in \mathcal{J}_i\}$ and tori $\mathbf{S}_i = \prod_{k \in \mathcal{J}_i \setminus \{0\}} \mathbf{T}_k$ with $i \in \{0, 1\}$. The decomposition $V = V_1 \oplus V_0$ is an orthogonal decomposition of V and we have a corresponding subgroup $\mathbf{M}_1 \times \mathbf{M}_0 \leq \mathbf{G}$, where $\mathbf{M}_i = GL(V_i \mid \mathcal{B})$. Note that the group $\mathbf{L}_J = \mathbf{S}_1 \times \mathbf{M}_0^\circ$ is a standard Levi subgroup of \mathbf{G}° with $J \subseteq \Delta$. Identifying $\mathbf{W}_i = N_{\mathbf{M}_i}(\mathbf{S}_i)/\mathbf{S}_i$ with a subgroup of $\mathbf{W} = W_n$, we have $\mathbf{W}_1 \times \mathbf{W}_0 = W_m \times W_{n-m}$ is the subgroup of type $B_m \times B_{n-m}$ in 7.3. We recall the following well-known result.

Lemma 7.6. *Assume $\mathbf{L}_J = \mathbf{S}_1 \times \mathbf{M}_0^\circ$ is a standard Levi subgroup as in 7.5 such that $1 \leq m \leq n - 2$. Then we have $N_{\mathbf{W}^\circ}(J) \cong W_m$ is a Weyl group of type B_m with generators*

$$\begin{cases} \{s_1, \dots, s_{m-1}, u_m\} & \text{when } \mathbf{G} = \mathrm{GO}(V) \text{ and } \dim(V) \text{ is even,} \\ \{s_1, \dots, s_{m-1}, t_m\} & \text{otherwise.} \end{cases}$$

7.7. Now assume $\dim(V)$ is even. In this case, we let $\tilde{\mathbf{G}} = \mathbf{G} \cdot Z(GL(V))$ be the corresponding conformal group and similarly $\tilde{\mathbf{G}}^\circ = \mathbf{G}^\circ \cdot Z(GL(V))$. The inclusion $\mathbf{G}^\circ \rightarrow \tilde{\mathbf{G}}^\circ$ is a regular embedding, as in 3.3, and the group $\tilde{\mathbf{T}} = \mathbf{T} \cdot Z(GL(V))$ is a maximal

torus of $\widetilde{\mathbf{G}}^\circ$. We define a closed embedding $\mathbf{z} : \mathbb{F}^\times \rightarrow \widetilde{\mathbf{T}}$ by setting $\mathbf{z}(\zeta) = \check{\varepsilon}_1(\zeta) \cdots \check{\varepsilon}_n(\zeta)$. Writing \mathbf{Z} for the image of \mathbf{z} , we have $\widetilde{\mathbf{T}} = \mathbf{T} \times \mathbf{Z}$. Moreover, we have $\widetilde{\mathbf{M}}_0 = \mathbf{M}_0 \cdot \mathbf{Z}$ and $\widetilde{\mathbf{M}}_0^\circ = \mathbf{M}_0^\circ \cdot \mathbf{Z}$ are also isomorphic to conformal groups, and $\widetilde{\mathbf{L}}_J = \mathbf{S}_1 \times \widetilde{\mathbf{M}}_0^\circ$. We note that for any $1 \leq k \leq n$, we have

$${}^t_k \mathbf{z}(\zeta) = \mathbf{d}_k(\zeta^{-1}) \mathbf{z}(\zeta) \quad \text{and} \quad {}^u_k \mathbf{z}(\zeta) = \mathbf{d}_k(\zeta^{-1}) \mathbf{d}_n(\zeta^{-1}) \mathbf{z}(\zeta).$$

Furthermore, for $\prod_{i=1}^n \mathbf{d}_i(\zeta_i) \in \mathbf{T}$ and $1 \leq k \leq n$, we have

$${}^t_k \left(\left(\prod_{i=1}^n \mathbf{d}_i(\zeta_i) \right) \mathbf{z}(\zeta) \right) = \mathbf{d}_k(\zeta_k^{-1} \zeta^{-1}) \left(\prod_{i \neq k} \mathbf{d}_i(\zeta_i) \right) \mathbf{z}(\zeta).$$

Automorphisms and Frobenius Endomorphisms

7.8. If $q = p^\alpha$ is a power of p with $\alpha \geq 0$ an integer, then we denote by $F_q : V \rightarrow V$ the unique semi-linear map fixing the basis \mathcal{V} and satisfying $F_q(av) = a^q v$ for all $a \in \mathbb{F}$. Precomposing with F_q gives a Frobenius endomorphism $F_q : \text{GL}(V) \rightarrow \text{GL}(V)$. We also denote by F_q the restriction of this map to any F_q -stable subgroup. We will let $\gamma_1 \in \text{Aso}(\mathbf{G})$ be the identity unless $\mathbf{G} = \text{GO}(V)$ and $\dim(V)$ is even, in which case $\gamma_1 = \iota_g$ where $g = n_{s_n} \in \text{N}_{\mathbf{G}}(\mathbf{T})$. Moreover, we let $\gamma_2 := \tau_\alpha$ be as in (i) of Lemma 2.1, where $\alpha : \mathbf{G}/\mathbf{G}^\circ \rightarrow \mathbf{G}^\circ \cap Z(\mathbf{G})$ is the unique isomorphism when it exists and the identity otherwise. Note γ_2 is the identity on the connected component.

Now let $\Gamma(\mathbf{G}) = \langle F_p \rangle \times \langle \gamma_1, \gamma_2 \rangle \leq \text{Aso}(\mathbf{G})$ be the submonoid generated by these elements. For $q = p^\alpha$ with $\alpha \geq 0$, we let $\Gamma_q(\mathbf{G}) := F_q \langle \gamma_1, \gamma_2 \rangle \subseteq \Gamma(\mathbf{G})$ be the coset containing F_q . We also define $\Gamma(\mathbf{G}^\circ) := \langle F_p \rangle \times \langle \gamma_1 \rangle \leq \text{Aso}(\mathbf{G})$ and $\Gamma_q(\mathbf{G}^\circ) := F_q \langle \gamma_1 \rangle \subseteq \Gamma(\mathbf{G})$. Up to conjugation by \mathbf{G}° , any Frobenius endomorphism endowing \mathbf{G} with an \mathbb{F}_q -rational structure is contained in $\Gamma_q(\mathbf{G})$. We say $F \in \Gamma_q(\mathbf{G})$ is *split* if $F = F_q$ and *twisted* otherwise. Let $F \in \Gamma_q(\mathbf{G})$ and denote by $\Gamma(\mathbf{G}, F)$ the image of $C_{\Gamma(\mathbf{G})}(F)$ in $\text{Aut}(\mathbf{G}^F)$, similarly by $\Gamma(\mathbf{G}^\circ, F)$ the image of $C_{\Gamma(\mathbf{G}^\circ)}(F)$ in $\text{Aut}(\mathbf{G}^{\circ F})$.

Lemma 7.9. *If $\mathbf{G} = \text{GL}(V | \mathcal{B})$ is a classical group, then we have decompositions*

$$\text{Aso}(\mathbf{G}^\circ) = \text{Inn}(\mathbf{G}^\circ) \rtimes \Gamma(\mathbf{G}^\circ) \quad \text{and} \quad \text{Aso}(\mathbf{G}) = \text{Inn}(\mathbf{G}^\circ) \rtimes \Gamma(\mathbf{G}).$$

Moreover, if $F \in \Gamma_q(\mathbf{G})$ then we also have decompositions

$$\text{Aso}(\mathbf{G}^\circ, F) = \text{Inn}(\mathbf{G}^\circ, F) \rtimes \Gamma(\mathbf{G}^\circ, F) \quad \text{and} \quad \text{Aso}(\mathbf{G}, F) = \text{Inn}(\mathbf{G}^\circ, F) \rtimes \Gamma(\mathbf{G}, F).$$

Finally, $\text{Aut}(\mathbf{G}^F) = \text{K}(\mathbf{G}^{\circ F}) \rtimes \text{Aso}(\mathbf{G}, F)$.

Proof. The statement for $\text{Aso}(\mathbf{G}^\circ)$ is well known, see [GLS98, Thm. 1.15.7]. The restriction maps $\text{Aso}(\mathbf{G}) \rightarrow \text{Aso}(\mathbf{G}^\circ)$ and $\text{Aut}(\mathbf{G}^F) \rightarrow \text{Aut}(\mathbf{G}^{\circ F})$ are surjective. The kernels are calculated using Lemma 2.1. ■

Remark 7.10. When $\mathbf{G} = \text{GO}(V)$, we have $\Gamma_q(\mathbf{G})$ has cardinality four. Therefore, $\text{GO}(V)$ has four distinct \mathbb{F}_q -rational forms whereas $\text{SO}(V)$ has only two.

Dual groups and Dual automorphisms

7.11. Let us now assume that $\mathbf{G} = \mathrm{SO}(V)$ is a special orthogonal group. We let $V^* \subseteq V$ be the subspace spanned by $\mathcal{V} \setminus \{v_0\}$, with \mathcal{V} as in (7.2). The dual of \mathbf{G} is then the group $\mathrm{SL}(V^* | \mathcal{B}^*)$ where \mathcal{B}^* is the form from 7.1 satisfying $\mathcal{B}^*(v, w) = (-1)^{\dim(V)} \mathcal{B}^*(w, v)$ for all $v, w \in V^*$. In particular, $\mathbf{G} = \mathbf{G}^*$ if $\dim(V)$ is even. Recalling from [Tay18, 5.3] the notion of duality between isogenies, we have the following.

Lemma 7.12. *Assume $\mathbf{G} = \mathrm{SO}(V)$ is a special orthogonal group and let $*$: $\Gamma(\mathbf{G}) \rightarrow \Gamma(\mathbf{G}^*)$ be the unique monoid isomorphism satisfying $F_p^* = F_p$ and $\gamma_1^* = \gamma_1$. Then for any $\sigma \in \Gamma(\mathbf{G})$, we have $\sigma^* \in \Gamma(\mathbf{G}^*)$ is an asogeny dual to σ .*

8. Action of Automorphisms on Unipotent Characters

From now on, we assume that \mathbf{G} is either $\mathrm{GL}(V | \mathcal{B})$ or $\mathrm{SL}(V | \mathcal{B})$ and $F \in \Gamma_q(\mathbf{G})$ is a Frobenius endomorphism, with q a power of $p \neq 2$, as in Section 7.

We recall that the set $\mathcal{E}(\mathbf{G}^F, 1) \subseteq \mathrm{Irr}(\mathbf{G}^F)$ of unipotent characters consists of those irreducible characters $\chi \in \mathrm{Irr}(\mathbf{G}^F)$ whose restriction to $\mathbf{G}^{\circ F}$ contains a unipotent character.

Lemma 8.1. *Assume $\mathbf{G} = \mathrm{GO}(V)$ is an orthogonal group with $\dim(V) \geq 5$. If $\chi \in \mathcal{E}(\mathbf{G}^F, 1)$ is a unipotent character, then $\chi^\sigma = \chi$ for any $\sigma \in \mathrm{Aut}(\mathbf{G}^F)$.*

Proof. Let $\chi \in \mathcal{E}(\mathbf{G}^F, 1)$ be a unipotent character and let $\psi_1 \in \mathcal{E}(\mathbf{G}^{\circ F}, 1)$ be an irreducible constituent of the restriction $\mathrm{Res}_{\mathbf{G}^{\circ F}}^{\mathbf{G}^F}(\chi)$. As $Z(\mathbf{G})^F = Z(\mathbf{G}^\circ)^F \leq \mathbf{G}^{\circ F}$, it follows that $\omega_\chi = \omega_{\psi_1}$, which is trivial by Proposition 2.4(i). Thus if $\sigma \in \langle \gamma_2 \rangle \mathbf{K}(\mathbf{G}^F)$, then the statement holds by Lemma 2.1(iii).

As γ_1 is inner, it suffices to consider the case where $\sigma \in \mathrm{Inn}(\mathbf{G}^\circ, F) \langle F_p \rangle$. We have $\mathrm{Res}_{\mathbf{G}^{\circ F}}^{\mathbf{G}^F}(\chi)$ is either irreducible or the sum of two distinct irreducible constituents. If $\mathrm{Res}_{\mathbf{G}^{\circ F}}^{\mathbf{G}^F}(\chi) = \psi_1 + \psi_2$ with $\psi_1 \neq \psi_2$ irreducible, then $\chi = \mathrm{Ind}_{\mathbf{G}^{\circ F}}^{\mathbf{G}^F}(\psi_1)$. It is well known that $\psi_1^\sigma = \psi_1$, so $\chi^\sigma = \chi$ in this case. Now consider the case where $\mathrm{Res}_{\mathbf{G}^{\circ F}}^{\mathbf{G}^F}(\chi) = \psi_1$ is irreducible. Then $\mathrm{Ind}_{\mathbf{G}^{\circ F}}^{\mathbf{G}^F}(\psi_1) = \chi + \theta\chi$, where θ is the inflation of the non-trivial character of $\mathbf{G}^F/\mathbf{G}^{\circ F}$. Since σ fixes θ , it suffices to show that one character lying over $\psi := \psi_1$ is fixed by σ .

Denote by S the quotient group $\mathrm{OP}'(\mathbf{G}^F)/Z(\mathrm{OP}'(\mathbf{G}^F))$, which is a finite simple group. Recall that the unipotent character ψ restricts irreducibly to $\mathrm{OP}'(\mathbf{G}^F)$ and has $Z(\mathrm{OP}'(\mathbf{G}^F))$ in its kernel. Let $\eta \in \mathrm{Irr}(S)$ be the deflation of $\mathrm{Res}_{\mathrm{OP}'(\mathbf{G}^F)}^{\mathbf{G}^{\circ F}}(\psi)$ to S . If $H \leq \mathrm{Aut}(S)$ is the stabiliser of η , then by [Mal08, Thm. 2.4] there exists a character $\tilde{\eta} \in \mathrm{Irr}(H)$ extending η . If $\alpha : \mathbf{G}^F \rightarrow \mathrm{Aut}(S)$ is the natural map, then $\langle \alpha(\mathbf{G}^F), \sigma \rangle \leq H$. Then inflating the restriction $\mathrm{Res}_{\alpha(\mathbf{G}^F)}^H(\tilde{\eta})$ yields a σ -invariant irreducible character of \mathbf{G}^F extending ψ . ■

8.2. We record here an elementary observation that will be used in the next proof and later sections. For $i \in \{1, 2\}$, let G_i be a finite group with a subgroup $N_i \leq G_i$ of index two. Let $G = G_1 \times G_2$ and $N = N_1 \times N_2$ and assume we have a subgroup

$N \leq G^\circ \leq G$. For any subgroup $H \leq G$, we set $\bar{H} = HN/N$, so $\bar{G} = \bar{G}_1 \times \bar{G}_2 \cong C_2 \times C_2$ is a Klein four group and $\bar{G}^\circ \leq \bar{G}$.

Let $\chi \in \text{Irr}(G^\circ)$ be an irreducible character and let $\chi_i \in \text{Irr}(N_i)$ be such that $\chi_1 \boxtimes \chi_2 \in \text{Irr}(N)$ is a constituent of the restriction $\text{Res}_N^{\bar{G}}(\chi)$. The group \bar{G} acts naturally by conjugation on $\text{Irr}(N)$ and

$$\bar{G}_{\chi_1 \boxtimes \chi_2} = (\bar{G}_1)_{\chi_1} \times (\bar{G}_2)_{\chi_2}.$$

If \bar{G}° is the diagonal embedding of C_2 into \bar{G} , then one of the following holds:

- $\bar{G}_{\chi_1 \boxtimes \chi_2}^\circ \neq 1$ and $\chi = \text{Res}_{\bar{G}^\circ}^{\bar{G}}(\tilde{\chi}_1 \boxtimes \tilde{\chi}_2)$ with $\tilde{\chi}_i \in \text{Irr}(G_i)$ extending χ_i ,
- $\bar{G}_{\chi_1 \boxtimes \chi_2}^\circ = 1$ and $\chi = \text{Ind}_N^{G^\circ}(\chi_1 \boxtimes \chi_2)$.

For the following we recall also that the group $(G/G^\circ)^\vee$ of linear characters of G/G° also acts naturally on $\text{Irr}(G)$ by tensoring with the inflation.

Proposition 8.3. *Assume $\mathbf{G} = \text{GO}(V)$ and $s \in \mathbf{G}^F$ is a semisimple element such that $s^2 = 1$. We identify $A_{\mathbf{G}^{\circ F}}(s)$ with $C_{\mathbf{G}^\circ}(s)^F / C_{\mathbf{G}^\circ}^\circ(s)^F$. Let $\gamma_1 \in \text{Aso}(\mathbf{G}, F)$ be as in 7.8. If $\chi \in \mathcal{E}(C_{\mathbf{G}^\circ}(s)^F, 1)$ is a unipotent character, then the following hold:*

- (i) $\chi^{F_p} = \chi$,
- (ii) if $s \neq 1$ then $\chi^{\gamma_1} = \chi$ if and only if $A_{\mathbf{G}^{\circ F}}(s)_\chi^\vee \neq A_{\mathbf{G}^{\circ F}}(s)^\vee$.

Proof. We have $V = V_1 \oplus V_{-1}$, where V_ζ is the ζ -eigenspace of s . Moreover, $C_{\mathbf{G}}(s) = \text{GO}(V_1) \times \text{GO}(V_{-1})$. The quotient $A_{\mathbf{G}^F}(s) \cong C_{\mathbf{G}}(s)^F / C_{\mathbf{G}}^\circ(s)^F$ is a Klein four group and $A_{\mathbf{G}^{\circ F}}(s)$ is the diagonally embedded subgroup. Let $\chi_1 \boxtimes \chi_2 \in \mathcal{E}(C_{\mathbf{G}}^\circ(s)^F, 1)$ be an irreducible constituent of $\text{Res}_{C_{\mathbf{G}}^\circ(s)^F}^{C_{\mathbf{G}}(s)^F}(\chi)$. By the discussion in 8.2, either χ is the restriction of a character of $C_{\mathbf{G}}(s)^F$, which is F_p -fixed by Lemma 8.1, or is the induction of $\chi_1 \boxtimes \chi_2$, which is known to be F_p -fixed. Hence, χ is F_p -fixed, giving (i). Statement (ii) follows from the description above. \blacksquare

9. Power Maps and Classical Groups

9.1. Recall that a partition of an integer $N \geq 0$ is a weakly decreasing sequence of positive integers $\mu = (\mu_1, \mu_2, \dots)$ such that $\sum_{i \in \mathbb{N}} \mu_i = N$. For any $m \in \mathbb{N}$, we set $r_m(\mu) := |\{i \in \mathbb{N} \mid \mu_i = m\}|$. If $\epsilon \in \{0, 1\}$, we denote by $\mathcal{P}_\epsilon(N)$ the set of all partitions μ of N such that $r_m(\mu) \equiv 0 \pmod{2}$ whenever $m \equiv \epsilon \pmod{2}$. For $\mu \in \mathcal{P}(N)$, we set

$$\alpha(\mu, \epsilon) := |\{m \in \mathbb{N} \mid r_m(\mu) \neq 0 \text{ and } m \equiv 1 + \epsilon \pmod{2}\}|,$$

$$\delta(\mu, \epsilon) := \begin{cases} 1 & \text{if } N \text{ is even and } r_m(\mu) \text{ is odd for some } m \equiv 1 + \epsilon \pmod{2}, \\ 0 & \text{otherwise.} \end{cases}$$

Let $\mathbf{G} = \text{GL}(V \mid \mathcal{B})$ with $N = \dim(V) \geq 0$ and $\epsilon \in \{0, 1\}$ such that $\mathcal{B}(v, w) = (-1)^\epsilon \mathcal{B}(w, v)$ for all $v, w \in V$. We have a bijection $\mathcal{P}_\epsilon(N) \rightarrow \text{Cl}_{\text{uni}}(\mathbf{G})$, denoted by

$\mu \mapsto \mathcal{O}_\mu$, such that μ gives the sizes of the Jordan blocks in the Jordan normal form of any $u \in \mathcal{O}_\mu$ in its natural action on V . Note that any unipotent class $\mathcal{O}_\mu \in \text{Cl}_{\text{uni}}(\mathbf{G})$ is contained in \mathbf{G}° but may not be a single \mathbf{G}° -conjugacy class. Recall that a unipotent element $u \in \mathbf{G}^\circ$ is said to be \mathbf{G}° -distinguished if any torus of $C_{\mathbf{G}}^\circ(u) = C_{\mathbf{G}^\circ}^\circ(u)$ is contained in $Z^\circ(\mathbf{G}^\circ) = Z^\circ(\mathbf{G})$.

Lemma 9.2. *Let $\pi : \mathbf{G}^\circ \rightarrow \mathbf{G}_{\text{ad}}$ be an adjoint quotient of \mathbf{G}° . Then π defines a bijection $\text{Cl}_{\text{uni}}(\mathbf{G}^\circ) \rightarrow \text{Cl}_{\text{uni}}(\mathbf{G}_{\text{ad}})$ between the sets of unipotent classes. If $u \in \mathcal{O}_\mu$, then $A_{\mathbf{G}}(u)$, $A_{\mathbf{G}^\circ}(u)$, and $A_{\mathbf{G}_{\text{ad}}}(\pi(u))$ are all elementary abelian 2-groups whose respective orders are the maximum of 1 and: $2^{\alpha(\mu, \epsilon)}$, $2^{\alpha(\mu, \epsilon) - 1}$, and $2^{\alpha(\mu, \epsilon) - 1 - \delta(\mu, \epsilon)}$.*

Proof. This follows immediately from [LS12, Thm. 3.1]. ■

Lemma 9.3. *If $u \in \mathbf{G}$ is a \mathbf{G}° -distinguished unipotent element, then the subset $A \subseteq C_{\mathbf{G}}(u)$ of all semisimple elements is a subgroup and $C_{\mathbf{G}}(u) = C_{\mathbf{G}}^\circ(u) \times A$. Moreover, the group $\mathbf{R} = C_{\mathbf{G}}(A)$ is reductive, $u \in \mathbf{R}$ is regular unipotent, and $A = Z(\mathbf{R})$.*

Proof. Let $\epsilon \in \{0, 1\}$ be such that $\mathcal{B}(v, w) = (-1)^\epsilon \mathcal{B}(w, v)$ for all $v, w \in V$. Let $\mu := (n_1, \dots, n_t) \in \mathcal{P}_\epsilon(\mathbb{N})$ be a partition such that $n_1 > \dots > n_t$ and $n_i \equiv 1 + \epsilon \pmod{2}$ for all $1 \leq i \leq t$. Correspondingly, we can choose an orthogonal decomposition $V_1 \oplus \dots \oplus V_t$ of V such that $\dim(V_i) = n_i$. We also have a subgroup $\mathbf{R} = \mathbf{R}_1 \times \dots \times \mathbf{R}_t$ of \mathbf{G} where $\mathbf{R}_i = \text{GL}(V_i | \mathcal{B})$ and we identify \mathcal{B} with its restriction to V_i .

Fix a regular unipotent element $u \in \mathbf{R}$. Then $u \in \mathcal{O}_\mu$ is \mathbf{G}° -distinguished and all distinguished unipotent elements arise in this way, see [LS12, Prop. 3.5]. Recall that $\mathbf{R}_i = \text{GO}(V_i)$, not $\text{SO}(V_i)$, when $\epsilon = 0$. We have $Z(\mathbf{R})$ is an elementary abelian 2-group and $C_{\mathbf{G}}(u) = C_{\mathbf{G}}^\circ(u) \times Z(\mathbf{R})$. Indeed, $C_{\mathbf{G}}^\circ(u) \cap Z(\mathbf{R})$ is trivial, as $C_{\mathbf{G}}^\circ(u)$ is unipotent and $|Z(\mathbf{R})| = 2^t = 2^{\alpha(\mu, \epsilon)} = |A_{\mathbf{G}}(u)|$. Finally, it is clear that $\mathbf{R} = C_{\mathbf{G}}(A)$ and $A = Z(\mathbf{R})$. ■

9.4. We now assume that $u \in \mathbf{G}^{\text{F}}$ is any unipotent element. Fix an F-stable maximal torus $\mathbf{S} \leq C_{\mathbf{G}}^\circ(u)$ and consider the subgroup $\mathbf{M} = C_{\mathbf{G}}(\mathbf{S})$, which is necessarily F-stable. We have an isomorphism

$$\mathbf{M} \cong \text{GL}(U_1) \times \dots \times \text{GL}(U_d) \times \text{GL}(V_0 | \mathcal{B}) \tag{9.5}$$

for some decompositions $V = V_1 \oplus V_0$, as in 7.5, and $V_1 = (U_1 \oplus U'_1) \oplus \dots \oplus (U_d \oplus U'_d)$, where U_i and U'_i are totally isotropic subspaces. The element u is \mathbf{M}° -distinguished, where $\mathbf{M}^\circ = C_{\mathbf{G}^\circ}(\mathbf{S})$ is an F-stable Levi subgroup of \mathbf{G}° . We now show that u is regular in a reductive subgroup of \mathbf{G} . This idea also appears in work of Testerman [Tes95, §3].

Proposition 9.6. *Assume $u \in \mathbf{G}^{\text{F}}$ is a unipotent element and $\mathbf{S} \leq C_{\mathbf{G}}^\circ(u)$ is an F-stable maximal torus. Let $\mathbf{M} = C_{\mathbf{G}}(\mathbf{S})$ and denote by $A \subseteq C_{\mathbf{M}}(u)$ the subset of semisimple elements. Then u is regular unipotent in the F-stable reductive subgroup $\mathbf{R} = C_{\mathbf{G}}(A)$ and $A = Z(\mathbf{R})$. Moreover, the natural map $Z(\mathbf{R}) \rightarrow A_{\mathbf{R}}(u)$ is surjective and the natural map $A_{\mathbf{M}}(u) \rightarrow A_{\mathbf{G}}(u)$ is injective.*

Proof. Note that if $\mathbf{H} = \text{GL}(V)$, then a unipotent element $v \in \mathbf{H}$ is distinguished if and only if it is regular. For such an element, $Z(\mathbf{H}) \leq C_{\mathbf{H}}(v)$ gives the subset of semisimple

elements, which is clearly a subgroup. After Lemma 9.3 and (9.5), it follows immediately that $A = Z(\mathbf{R})$ is a subgroup, $\mathbf{R} = C_G(A)$ is reductive, and $u \in \mathbf{R}$ is regular.

The natural maps $Z(\mathbf{R})/Z^\circ(\mathbf{R}) \rightarrow A_{\mathbf{R}}(u) \rightarrow A_{\mathbf{M}}(u)$ are isomorphisms, which follows from Lemma 9.3. Now, as $\mathbf{M} = C_G(\mathbf{S})$ is the centralizer of a torus, a standard argument shows that the map $A_{\mathbf{M}}(u) \rightarrow A_{\mathbf{G}}(u)$ is injective, see [Spa85, 1.4]. Hence $A_{\mathbf{R}}(u) \rightarrow A_{\mathbf{G}}(u)$ is injective. \blacksquare

Corollary 9.7. *If $\mathbf{G} = \mathrm{GO}(V)$, then any unipotent element $u \in \mathbf{G}^{\circ F}$ is rational in $\mathbf{G}^{\circ F}$.*

Proof. The group \mathbf{R}° is a direct product of general linear and special orthogonal groups. As $u \in \mathbf{R}^\circ$ is a regular element, it follows from Lemma 5.6 that u and u^k are conjugate in $\mathbf{R}^{\circ F}$ and hence also in $\mathbf{G}^{\circ F}$. \blacksquare

Corollary 9.8. *Assume $\mathbf{G} = \mathrm{Sp}(V)$ with $\dim(V) = 2n$. We have $\mathbf{R} = \mathbf{R}^\circ$ is a direct product of symplectic and general linear groups. If $g \in \mathbf{R}$ is such that ${}^g u = u^k$, then the map τ of Lemma 5.2 is given by $\tau([C_G^\circ(u)a]) = [C_G^\circ(u)a\mathcal{L}(g)]$. In particular, if $u \in \mathcal{O}_\mu$ with $\mu \in \mathcal{P}_1(2n)$, then u is \mathbf{G}^F -conjugate to u^k if and only if one of the following hold:*

- (i) $r_{2m}(\mu)$ is even for all $1 \leq m \leq n$,
- (ii) $k \pmod{p} \in \mathbb{F}_q$ is a square.

Proof. It is known that there exists an F -stable maximal torus $\mathbf{S} \leq \mathbf{G}$ for which the natural map $\mathbf{S} \rightarrow A_{\mathbf{G}}(u)$ is surjective, see the proof of [BDT19, Prop. 9.3] for instance. The statements now follow from Proposition 9.6. \blacksquare

10. Cuspidal Characters of Orthogonal Groups

10.1. In this section, we will give analogues of the calculations in [Tay18, §14] for the special orthogonal groups. For this, we introduce a little notation. We refer the reader to [Tay18] for more details. For any integers $m, \delta \in \mathbb{N}_0$, we let

$$\Delta(\delta, m) = \begin{cases} 1 & \text{if } \delta = 0 \text{ and } m \neq 0, \\ 0 & \text{otherwise.} \end{cases}$$

Given integers $r, n \in \mathbb{N}_0$ and $d \in \{0, 1\}$, we will denote by $\mathbb{X}_{n,d}^r := \mathbb{X}_{n,d}^{r,0}$ the sets of symbols defined in [LS85, §1], see [Tay18, 13.4].

For any n , we let ${}^\dagger \mathbb{X}_{n,0}^0 \subseteq \mathbb{X}_{n,0}^0$ be the set of symbols $\begin{bmatrix} a_1 & \cdots & a_m \\ b_1 & \cdots & b_m \end{bmatrix}$ such that $\sum_{i=1}^m a_i > \sum_{i=1}^m b_i$. For compatibility, we also let ${}^\dagger \mathbb{X}_{n,1}^0 = \mathbb{X}_{n,1}^0$. If $e \in \mathbb{N}$, then we will need the following *special symbols*

$$s_{e,1} = \begin{bmatrix} 0 & 1 & \cdots & e-1 & e \\ 1 & 2 & \cdots & e & \end{bmatrix} \in {}^\dagger \mathbb{X}_{e(e+1),1}^0 \quad \text{and} \quad s_{e,0} = \begin{bmatrix} 1 & 2 & \cdots & e \\ 0 & 1 & \cdots & e-1 \end{bmatrix} \in {}^\dagger \mathbb{X}_{e^2,0}^0.$$

Recall we have a bijection ${}^\dagger \mathbb{X}_{n,1}^0 \rightarrow \mathrm{Irr}(W_n)$ and an injection ${}^\dagger \mathbb{X}_{n,0}^0 \rightarrow \mathrm{Irr}(W'_n)$, defined as in [Lus84, 4.5], where $W'_n \leq W_n$ are as in 7.3.

Let $\mathbf{G} = \mathrm{SO}(V)$ be a special orthogonal group with $\dim(V) = 2n + \delta$ and $\delta \in \{0, 1\}$. We have a regular embedding $\iota : \mathbf{G} \rightarrow \tilde{\mathbf{G}}$, where $\tilde{\mathbf{G}} = \mathbf{G} \cdot \mathrm{Z}(\mathrm{GL}(V))$ is the corresponding conformal group, as in 7.7. We denote by $\tilde{\mathbf{G}}^*$ a dual group of $\tilde{\mathbf{G}}$ and by $\iota^* : \tilde{\mathbf{G}}^* \rightarrow \mathbf{G}^*$ a surjective homomorphism dual to the embedding. The dual group \mathbf{G}^* is as in 7.11. By the classification of quasi-isolated semisimple elements, we have a semisimple element $s \in \mathbf{G}^*$ is quasi-isolated if and only if $s^2 = 1$, see [Bon05, Prop. 4.11, Exmp. 4.10]. If $V_{\pm 1}^*(s)$ is the (± 1) -eigenspace for s , then $\dim(V_{\pm 1}^*(s))$ is even and $C_{\mathbf{G}^*}^{\circ}(s) = \mathrm{SL}(V_1^*(s) \mid \mathcal{B}^*) \times \mathrm{SL}(V_{-1}^*(s) \mid \mathcal{B}^*)$.

Now, if $s \in \mathbf{G}^{*\mathrm{F}^*}$ is F^* -fixed and $\tilde{s} \in \tilde{\mathbf{G}}^{*\mathrm{F}^*}$ is such that $\iota^*(\tilde{s}) = s$, then ι^* restricts to a surjective homomorphism $C_{\tilde{\mathbf{G}}^*}(\tilde{s})^{\mathrm{F}^*} \rightarrow C_{\mathbf{G}^*}^{\circ}(s)^{\mathrm{F}^*}$. Conjugating \tilde{s} into $\tilde{\mathbf{T}}^*$, we identify the Weyl group $\tilde{\mathbf{W}}^*(s)$ of $C_{\tilde{\mathbf{G}}^*}(\tilde{s})$ with a subgroup of $\tilde{\mathbf{W}}^*$. Using inflation, we identify the unipotent characters of these groups. The following may be extracted from the work of Lusztig in a way entirely analogous to that of [Tay18, Lem. 14.3], so we omit the details. Here $n_{\tilde{\chi}}$, for $\tilde{\chi} \in \mathrm{Irr}(\tilde{\mathbf{G}}^{\mathrm{F}})$, is defined as in [Tay18, 14.1]

Lemma 10.2 (Lusztig). *Assume $\mathbf{G} = \mathrm{SO}(V)$ and let $\dim(V) = 2n + \delta$ with $\delta \in \{0, 1\}$. Let $\tilde{s} \in \tilde{\mathbf{G}}^{*\mathrm{F}^*}$ be a semisimple element whose image $s = \iota^*(\tilde{s}) \in \mathbf{G}^{*\mathrm{F}^*}$ is quasi-isolated with $2a = \dim(V_1^*(s))$ and $2b = \dim(V_{-1}^*(s))$. If $\tilde{\chi} \in \mathcal{E}(\tilde{\mathbf{G}}^{\mathrm{F}}, \tilde{s})$ is a cuspidal character, then the following hold:*

- (i) $\tilde{\chi}$ is the unique cuspidal character in $\mathcal{E}(\tilde{\mathbf{G}}^{\mathrm{F}}, \tilde{s})$ and $a = e(e + \delta)$ and $b = f(f + \delta)$, for some non-negative integers e and f , and either $e + \delta \geq 2$ or $f + \delta \geq 2$,
- (ii) $\tilde{\chi}$ is contained in the family $\mathcal{E}(\tilde{\mathbf{G}}^{\mathrm{F}}, \tilde{s}, \tilde{\mathcal{C}}) \subseteq \mathcal{E}(\tilde{\mathbf{G}}^{\mathrm{F}}, \tilde{s})$, where $\mathrm{Irr}(\tilde{\mathbf{W}}^*(\tilde{s}) \mid \tilde{\mathcal{C}})$ contains the special character $\mathcal{S}_{e,\delta} \boxtimes \mathcal{S}_{f,\delta} \in \dagger \mathbb{X}_{a,\delta}^0 \times \dagger \mathbb{X}_{b,\delta}^0$,
- (iii) $n_{\tilde{\chi}} = 2^{e+f-\Delta(\delta,e)-\Delta(\delta,f)}$.

10.3. Let \mathbf{G}_{ad} be the adjoint group of the same type as \mathbf{G} , and let $\pi : \tilde{\mathbf{G}} \rightarrow \mathbf{G}_{\mathrm{ad}}$ be an adjoint quotient. Then $\pi \circ \iota : \mathbf{G} \rightarrow \mathbf{G}_{\mathrm{ad}}$ is also an adjoint quotient of \mathbf{G} . The maps $\pi \circ \iota$ and π define bijections $\mathrm{Cl}_{\mathrm{uni}}(\mathbf{G}) \rightarrow \mathrm{Cl}_{\mathrm{uni}}(\mathbf{G}_{\mathrm{ad}}) \leftarrow \mathrm{Cl}_{\mathrm{uni}}(\tilde{\mathbf{G}})$ between the unipotent classes of these groups. Hence we can label the unipotent classes as in Section 9. Moreover, we have $|\mathcal{A}_{\tilde{\mathbf{G}}}(\mathbf{u})| = |\mathcal{A}_{\mathbf{G}_{\mathrm{ad}}}(\pi(\mathbf{u}))|$ is given by Lemma 9.2.

Proposition 10.4. *Assume $\mathbf{G} = \mathrm{SO}(V)$ and $\tilde{s} \in \tilde{\mathbf{G}}^{*\mathrm{F}^*}$ is a semisimple element whose image $s = \iota^*(\tilde{s}) \in \mathbf{G}^*$ is quasi-isolated. If $\tilde{\chi} \in \mathcal{E}(\tilde{\mathbf{G}}^{\mathrm{F}}, \tilde{s})$ is a cuspidal irreducible character and $\mathbf{u} \in \mathcal{O}_{\tilde{\chi}}^*$ is a representative of the wave front set, then we have $n_{\tilde{\chi}} = |\mathcal{A}_{\tilde{\mathbf{G}}}(\mathbf{u})|$.*

Proof. Let $\dim(V) = 2n + \delta$ with $\delta \in \{0, 1\}$. We may identify $E = \mathcal{S}_{e,\delta} \boxtimes \mathcal{S}_{f,\delta}$ with an irreducible character of $\tilde{\mathbf{W}}^*(s)$. According to [Lus09b, 4.5(a), 6.3(b)], we have $j_{\tilde{\mathbf{W}}^*(s)}^{\tilde{\mathbf{W}}^*}(E) = \mathcal{S}_{e,\delta} \oplus \mathcal{S}_{f,\delta} \in \mathbb{X}_{n,\delta}^0$ with the j -induction defined in [GP00, §5.2] and the addition of symbols defined as in [LS85], see [Tay18, 13.6]. If $\Lambda_{0,\delta}^2$ is the unique element of $\mathbb{X}_{0,\delta}^2$ then we get a symbol $\mathrm{Spr}(j_{\tilde{\mathbf{W}}^*(s)}^{\tilde{\mathbf{W}}^*}(E)) = \mathcal{S}_{e,\delta} \oplus \mathcal{S}_{f,\delta} \oplus \Lambda_{0,\delta}^2 \in \mathbb{X}_{n,\delta}^2$. From this symbol, we may extract the partition of a unipotent class, as in [GM00, §2], which will be exactly the class $\mathcal{O}_{\tilde{\chi}}^*$. By symmetry, it suffices to treat the case where $0 \leq e \leq f$. We set $k = f - e \geq 0$.

Assume first that $\delta = 1$. Then the symbol $\text{Spr}(j_{\widetilde{\mathbf{W}}^*(s)}^{\widetilde{\mathbf{W}}^*}(E))$ is exactly the same as that written in the first case occurring in the proof of [Tay18, 14.4]. The unipotent class corresponding to this symbol is parameterised by the partition $\lambda = 2\mu + 1$, where

$$\mu = (k + 2e, \dots, k + 1, k, k - 1, k - 1, \dots, 1, 1, 0, 0).$$

As λ contains $k + (2e + 1) = e + f + 1$ distinct odd numbers, we have $n_{\widetilde{\chi}} = 2^{e+f} = |\Lambda_{\widetilde{\mathbf{G}}}(\mathbf{u})|$ by Lemmas 9.2 and 10.2.

Now assume $\delta = 0$. We have $\text{Spr}(j_{\widetilde{\mathbf{W}}^*(s)}^{\widetilde{\mathbf{W}}^*}(E))$ is the symbol

$$\begin{bmatrix} 0 & 1 & \dots & k-1 & k & k+1 & \dots & k+e-1 \\ 1 & 4 & \dots & 3k-2 & 3k+2 & 3k+6 & \dots & 3k+4e-2 \\ 0 & 3 & \dots & 3k-3 & 3k & 3k+4 & \dots & 3k+4e-4 \end{bmatrix}.$$

The unipotent class corresponding to this symbol is parameterised by the partition $\lambda = 2\mu + 1$, where

$$\mu = (2k + 4e - 1, \dots, k + 1, k, k - 1, k - 1, \dots, 1, 1, 0, 0).$$

As λ contains $k + 2e = e + f$ distinct odd numbers and at least one odd number occurs an odd number times, we have $n_{\widetilde{\chi}} = 2^{e+f-\Delta(0,e)-\Delta(0,f)} = |\Lambda_{\widetilde{\mathbf{G}}}(\mathbf{u})|$. \blacksquare

10.5. Recall from 7.8 that we have defined a subgroup $\Gamma(\mathbf{G}, F) \leq \text{Aut}(\mathbf{G}^F)$ of graph and field automorphism. We denote by $\mathcal{G}(\mathbf{G}, F)$ the direct product $\Gamma(\mathbf{G}, F) \times \mathcal{G}$, which acts on $\text{Irr}(\mathbf{G}^F)$ through the actions of $\text{Aut}(\mathbf{G}^F)$ and \mathcal{G} .

Theorem 10.6. *Assume $\mathbf{G} = \text{SO}(V)$ and $s \in \mathbf{G}^{*F}$ is quasi-isolated. Then for any $\sigma \in \text{Aso}(\mathbf{G}, F)$, we have $\mathcal{E}(\mathbf{G}^F, s)^\sigma = \mathcal{E}(\mathbf{G}^F, s)$. Moreover, any cuspidal character $\chi \in \mathcal{E}(\mathbf{G}^F, s)$ satisfies $\mathcal{G}(\mathbf{G}, F)_\chi = \mathcal{G}(\mathbf{G}, F)$.*

Proof. That the series $\mathcal{E}(\mathbf{G}^F, s)$ is σ -stable is shown as in [Tay18, Thm. 14.6], appealing to [SFT18, Lem. 3.4] when $\sigma \in \mathcal{G}$. Let $\tilde{s} \in \widetilde{\mathbf{G}}^*$ be a semisimple element such that $\iota(\tilde{s}) = s$ and let $\tilde{\chi} \in \mathcal{E}(\widetilde{\mathbf{G}}^F, \tilde{s})$ be a character covering χ . The constituents of $\text{Res}_{\widetilde{\mathbf{G}}^F}^{\widetilde{\mathbf{G}}^F}(\tilde{\chi})$ are the cuspidal characters in $\mathcal{E}(\mathbf{G}^F, s)$, so $\text{Res}_{\widetilde{\mathbf{G}}^F}^{\widetilde{\mathbf{G}}^F}(\tilde{\chi})$ is σ -stable. If χ extends to $\widetilde{\mathbf{G}}^F$, then it is the unique cuspidal character contained in $\mathcal{E}(\mathbf{G}^F, s)$ so must be $\mathcal{G}(\mathbf{G}, F)$ -invariant.

Assume now that χ does not extend to $\widetilde{\mathbf{G}}^F$. Then we must have $\dim(V)$ is even and $\text{Res}_{\widetilde{\mathbf{G}}^F}^{\widetilde{\mathbf{G}}^F}(\tilde{\chi}) = \chi_1 + \chi_2$ with $\chi_1 \neq \chi_2$ distinct irreducible characters. As $\chi \in \{\chi_1, \chi_2\}$, we must show that $\chi_i^\sigma = \chi_i$. The arguments in [Tay13, §2], together with Lemma 9.2, show that $Z(\mathbf{G})$ embeds in $\Lambda_{\mathbf{G}}(\mathbf{u})$. Arguing as in [Tay13, Prop. 5.4], we may find two unipotent elements $u_1, u_2 \in \mathcal{O}_\chi^{*F}$ such that $\langle \Gamma_{u_i}, \chi_j \rangle = \delta_{i,j}$. We claim that the character Γ_{u_i} is $\mathcal{G}(\mathbf{G}, F)$ -invariant. From this it follows that χ_i is $\mathcal{G}(\mathbf{G}, F)$ -invariant.

Firstly, we see that Γ_{u_i} is \mathcal{G} -invariant by Corollary 9.7 and [SFT18, Prop. 4.10]. The action of $\text{Aso}(\mathbf{G}, F)$ on Γ_{u_i} is described by [Tay18, 11.10]. The same argument as used in [Tay18, 13.2] shows that Γ_{u_i} is F_p -invariant. We now just need to show that Γ_{u_i} is $\text{GO}(V)^F$ -invariant. By Lemma 9.2, and the description of $\mathcal{O}_\chi^* = \mathcal{O}_{\tilde{\chi}}^*$ given in the proof of

Proposition 10.4, we have $C_{\mathrm{GO}(V)^F}(u)$ is not contained in $C_{\mathbf{G}^F}(u)$. As \mathbf{G}^F has index 2 in $\mathrm{GO}(V)^F$, we have the \mathbf{G}^F -class of u is $\mathrm{GO}(V)^F$ -invariant. \blacksquare

10.7. Let $\mathbf{G} = \mathrm{GO}(V)$ be an orthogonal group. We let $\mathcal{G}(\mathbf{G}, F) = \Gamma(\mathbf{G}, F) \times \mathcal{G}$ as above. Recall that a character $\chi \in \mathrm{Irr}(\mathbf{G}^F)$ is said to be cuspidal if the restriction $\mathrm{Res}_{\mathbf{G}^{\circ F}}^{\mathbf{G}^F}(\chi)$ contains a cuspidal irreducible character of $\mathbf{G}^{\circ F}$. In the following sections, we will need to have some control over cuspidal characters of $\mathrm{GO}(V)$ to construct certain extensions. Using the characters constructed in Section 6, we get the following.

Theorem 10.8. *Assume $\mathbf{G} = \mathrm{GO}(V)$. Let $s \in \mathbf{G}^{\circ*F}$ be a quasi-isolated semisimple element. If $\chi \in \mathrm{Irr}(\mathbf{G}^F)$ is a cuspidal character lying over $\chi^\circ \in \mathcal{E}(\mathbf{G}^{\circ F}, s)$ with $s^2 = 1$, then χ is $\mathcal{G}(\mathbf{G}, F)$ -invariant.*

Proof. By Theorem 10.6, the character χ° is $\mathcal{G}(\mathbf{G}^\circ, F)$ -invariant so extends to \mathbf{G}^F . Hence χ is one such extension. All other extensions are of the form $\theta\chi$, where θ is the inflation of an irreducible character of $\mathbf{G}^F/\mathbf{G}^{\circ F}$. Hence, it suffices to show that just one extension of χ° is $\mathcal{G}(\mathbf{G}, F)$ -invariant.

Now, we may find a GGGC Γ_u of $\mathbf{G}^{\circ F}$, with $u \in \mathcal{O}_\chi^{*F}$ as in the proof of Theorem 10.6, such that $\langle \Gamma_u, \chi \rangle = 1$. If $C_{\mathbf{G}}(u)_2^F \leq C_{\mathbf{G}}(u)^F$ is a Sylow 2-subgroup, then $C_{\mathbf{G}}(u)_2^F$ is not contained in $\mathbf{G}^{\circ F}$, so by Lemma 6.4, the character $\widehat{\Gamma}_{u,2}$ of \mathbf{G}^F extends Γ_u . It must contain only one of the two possible extensions of χ° . Moreover, arguing as in the proof of Theorem 10.6, it follows from Propositions 6.6 and 6.7 that $\widehat{\Gamma}_{u,2}$ is $\mathcal{G}(\mathbf{G}, F)$ -invariant. Hence, the unique extension of χ° that it contains must also be $\mathcal{G}(\mathbf{G}, F)$ -invariant. \blacksquare

11. Automorphisms and Quasi-Isolated Series

11.1. Recall the notation for cuspidal pairs and Harish-Chandra series from Section 3. If $s \in \mathbf{G}^{\circ*F}$ is a semisimple element, then we let $\mathrm{Cusp}_s(\mathbf{G}^\circ, F) \subseteq \mathrm{Cusp}(\mathbf{G}^\circ, F)$ be the set of cuspidal pairs (\mathbf{L}, λ) such that $\mathcal{E}(\mathbf{G}^{\circ F}, \mathbf{L}, \lambda) \subseteq \mathcal{E}(\mathbf{G}^{\circ F}, s)$, see [Bon06, Thm. 11.10]. Then we have

$$\mathcal{E}(\mathbf{G}^{\circ F}, s) = \bigsqcup_{(\mathbf{L}, \lambda) \in \mathrm{Cusp}_s(\mathbf{G}^\circ, F)/\mathbf{W}^F} \mathcal{E}(\mathbf{G}^{\circ F}, \mathbf{L}, \lambda),$$

where the pairs are taken up to the natural action of \mathbf{W}^F by conjugation. In the following sections, as in [Tay18, §15], we will study the actions of $\mathrm{Aut}(\mathbf{G}^F)$ and \mathcal{G} on the Harish-Chandra series above. The proof of the following is identical to that of [Tay18, 15.4].

Lemma 11.2. *Let $\mathbf{G} = \mathrm{GL}(V \mid \mathcal{B})$. Assume $s \in \mathbf{G}^{\circ*F}$ is a semisimple element with $s^2 = 1$ and let $(\mathbf{L}, \lambda) \in \mathrm{Cusp}_s(\mathbf{G}^\circ, F)$ be a cuspidal pair. Then $\mathbf{L} = \mathbf{Z}^\circ(\mathbf{L}) \times \mathbf{L}_{\mathrm{der}}$ and there exists an orthogonal decomposition $V = V_1 \oplus V_0$ as in 7.5 such that the following hold:*

- (i) $\mathbf{L}_{\mathrm{der}} = \mathbf{M}_0^\circ$ and $\mathbf{Z}^\circ(\mathbf{L}) = \mathbf{S}_1 \leq \mathbf{M}_1^\circ$ is a maximally split torus of \mathbf{M}_1° ,
- (ii) $\lambda = \lambda_1 \boxtimes \psi$ with $\lambda_1 \in \mathrm{Irr}(\mathbf{S}_1^F)$, $\psi \in \mathcal{E}(\mathbf{M}_0^{\circ F}, s_0)$ a cuspidal character, $s_0^2 = 1$, and $\lambda_1^2 = 1$.

11.3. Now assume $(\mathbf{L}, \lambda) \in \text{Cusp}_s(\mathbf{G}^\circ, F)$ is as in Lemma 11.2. Using the notation of Section 7, we let $\tau = p_{t_1} p_{t_n} \in N_{\mathbf{G}^\circ}(\mathbf{T})$. An argument with Weyl groups along the lines of that in [Tay18, 15.6] yields that

$$N_{\mathbf{G}^\circ}(\mathbf{L}) = (N_{\mathbf{M}_1^\circ}(\mathbf{S}_1) \times \mathbf{M}_0^\circ) \langle \tau \rangle \leq N_{\mathbf{G}}(\mathbf{L}) = N_{\mathbf{M}_1}(\mathbf{S}_1) \times \mathbf{M}_0.$$

With this in place, we have the following analogue of [Tay18, 15.6].

Lemma 11.4. *Assume \mathbf{G} , s , and (\mathbf{L}, λ) , are as in Lemma 11.2. Then for any extension $\Lambda(\lambda) \in \text{Irr}(N_{\mathbf{G}^\circ F}(\mathbf{L})_\lambda)$ of λ , the following hold:*

- (i) $\Lambda(\lambda)$ extends to a character of the form $\Lambda_1(\lambda_1) \boxtimes \Lambda_0(\psi)$ in $N_{\mathbf{G}^F}(\mathbf{L})_\lambda = N_{\mathbf{M}_1^F}(\mathbf{S}_1)_{\lambda_1} \times \mathbf{M}_0^F$, where $\Lambda_1(\lambda_1)$ and $\Lambda_0(\psi)$ are extensions of λ_1 and ψ to $N_{\mathbf{M}_1^F}(\mathbf{S}_1)_{\lambda_1}$ and \mathbf{M}_0^F , respectively;
- (ii) $\Lambda(\lambda)$ is $\Gamma(\mathbf{G}^\circ, F)$ -invariant.

Proof. By [Tay18, 15.6] we can assume $\mathbf{G} = \text{GO}(V)$. To prove part (i), it suffices to know that λ_1 and ψ extend, using 8.2. The group \mathbf{S}_1 has a $\Gamma(\mathbf{G}^\circ, F)$ -stable complement in $N_{\mathbf{M}_1}(\mathbf{S}_1)$ given by the permutation matrices. As λ_1 is linear, it extends trivially to $N_{\mathbf{M}_1^F}(\mathbf{S}_1)_{\lambda_1}$ and this must be $\Gamma(\mathbf{G}^\circ, F)$ -invariant. That ψ has an invariant extension is Theorem 10.8. \blacksquare

Proof (of Theorem A.(ii)). Let $\mathbf{G} = \text{SO}(V)$ and assume $(\mathbf{L}, \lambda) \in \text{Cusp}_s(\mathbf{G}, F)$ is as in Lemma 11.2. After Lemma 11.4, we see that λ has an F_p -invariant extension $\Lambda(\lambda)$. Since F_p is the identity on $W(\lambda)$, the statement follows using [MS16, 5.6], as in [Tay18, 15.6]. \blacksquare

12. Galois Automorphisms: Quasi-Isolated Series of $\text{SO}(V)$

12.1. Let $\mathbf{G} = \text{GL}(V \mid \mathcal{B})$ and let $s \in \mathbf{G}^{\circ * F^*}$ be a quasi-isolated semisimple element. We begin by introducing some notation to be used in this and the next section. Slightly abusing notation, we let $W^\circ(\lambda) = N_{\mathbf{G}^\circ F}(\mathbf{L})_\lambda / \mathbf{L}^F$ and $W(\lambda) = N_{\mathbf{G}^F}(\mathbf{L})_\lambda / \mathbf{L}^F$. Let Λ , Λ_1 , and Λ_0 be extension maps with respect to $\mathbf{L}^F \triangleleft N_{\mathbf{G}^\circ F}(\mathbf{L})$, $\mathbf{S}_1^F \triangleleft N_{\mathbf{M}_1^F}(\mathbf{S}_1)$, and $\mathbf{M}_0^{\circ F} \triangleleft \mathbf{M}_0^F$, respectively, such that $\Lambda(\lambda) = \text{Res}_{N_{\mathbf{G}^\circ F}(\mathbf{L})_\lambda}^{N_{\mathbf{G}^F}(\mathbf{L})_\lambda} (\Lambda_1(\lambda_1) \boxtimes \Lambda_0(\psi))$ as in Lemma 11.4. Then we similarly let $W(\lambda_1) = N_{\mathbf{M}_1^F}(\mathbf{S}_1)_{\lambda_1} / \mathbf{S}_1^F$ and $W^\circ(\lambda_1) = N_{\mathbf{M}_1^{\circ F}}(\mathbf{S}_1)_{\lambda_1} / \mathbf{S}_1^F$. We also have $W(\lambda) = W(\lambda_1) \times \mathbf{M}_0^F / \mathbf{M}_0^{\circ F}$, since the cuspidal character ψ is invariant under \mathbf{M}_0^F . Let $W^\circ(\lambda) = R(\lambda) \rtimes C(\lambda)$ and $W^\circ(\lambda_1) = R(\lambda_1) \rtimes C(\lambda_1)$ be decompositions as in 3.1.

For any $\sigma \in \mathcal{G}$, Gallagher's theorem implies that there exist linear characters $\delta_{\lambda_1, \sigma} \in \text{Irr}(W(\lambda_1))$ and $\delta_{\psi, \sigma} \in \text{Irr}(\mathbf{M}_0^F)$ such that $\Lambda_1(\lambda_1)^\sigma = \delta_{\lambda_1, \sigma} \Lambda_1(\lambda_1^\sigma)$ and similarly $\Lambda_0(\psi)^\sigma = \delta_{\psi, \sigma} \Lambda_0(\psi^\sigma)$. Moreover, for $\delta_{\lambda, \sigma}$ such that $\Lambda(\lambda)^\sigma = \delta_{\lambda, \sigma} \Lambda(\lambda^\sigma)$ as in Theorem 3.2, we have

$$\delta_{\lambda, \sigma} = \text{Res}_{W^\circ(\lambda)}^{W(\lambda)} (\delta_{\lambda_1, \sigma} \boxtimes \delta_{\psi, \sigma}).$$

Further, recall that the characters $\gamma_{\lambda, \sigma}$ and $\delta'_{\lambda, \sigma}$ of Theorem 3.2 are characters of $C(\lambda)$.

With this, we have mostly reduced ourselves to the case of principal series characters and cuspidal characters. The rest of this section is devoted to completing the proof of Theorem A. We first consider the case of principal series characters.

Proposition 12.2. *Assume $\mathbf{G} = \mathrm{GO}(V)$, $F \in \Gamma_q(\mathbf{G}^\circ)$, and $\lambda \in \mathrm{Irr}(\mathbf{T}^F)$ is a character such that $\lambda^2 = 1$. Then each member of $\mathcal{E}(\mathbf{G}^{\circ F}, \mathbf{T}, \lambda)$ is invariant under \mathcal{G} .*

Proof. Note that in this case, $\lambda = \lambda_1$. As in 7.3, we have $\mathbf{T} = \mathbf{T}_1 \times \cdots \times \mathbf{T}_n$. Each \mathbf{T}_i is F -stable, so $\mathbf{T}^F = \mathbf{T}_1^F \times \cdots \times \mathbf{T}_n^F$. Acting with \mathbf{W}^F , we can assume that $\lambda = 1_{\mathbf{T}_1^F} \boxtimes \cdots \boxtimes 1_{\mathbf{T}_a^F} \boxtimes \varepsilon_{\mathbf{T}_{a+1}^F} \boxtimes \cdots \boxtimes \varepsilon_{\mathbf{T}_{n-1}^F} \boxtimes \lambda_n$ for some $0 \leq a < n$, with $\lambda_n \in \{1_{\mathbf{T}_n^F}, \varepsilon_{\mathbf{T}_n^F}\}$. If F is split, then we will also assume that $\lambda_n = \varepsilon_{\mathbf{T}_n^F}$. Let $b = n - a$.

We will assume that $\dim(V)$ is even, as the statement in the case that $\dim(V)$ is odd follows from a simplified version of the argument. Note that $\mathbf{W}^F \leq \mathbf{W} = W_n$ is the subgroup W_n'' of type B_{n-1} , as in 7.3, when F is twisted. In the notation of 7.5, we have $W(\lambda) \leq \mathbf{W}_1 \times \mathbf{W}_0$ is $W_a \times W_b$ if F is split and $W_a \times W_b''$, of type $B_a \times B_{b-1}$, if F is twisted. Moreover, using Lemma 3.4 and 7.7, $R(\lambda) \leq \mathbf{W}_1 \times \mathbf{W}_0$ is the subgroup $W'_a \times W'_b$ of type $D_a \times D_b$ if F is split and $W'_a \times W_b''$, of type $D_a \times B_{b-1}$, if F is twisted. In all cases, the group $C(\lambda)$ is generated by u_1 , in the notation of 7.3, which has order two and even length in \mathbf{W} . Then for each $\sigma \in \mathcal{G}$, Lemma 3.6 implies $\gamma_{\lambda, \sigma} = 1$, and we further have $\Lambda_\lambda(n_{u_1})^2 = \lambda(1) = 1$ for any extension $\Lambda_\lambda := \Lambda(\lambda)$ of λ to $N_{\mathbf{G}^{\circ F}}(\mathbf{T})_\lambda$. Then $\Lambda_\lambda(n_{u_1}) \in \{\pm 1\}$ is fixed by σ , and hence $\delta'_{\lambda, \sigma} = 1$.

It remains to consider the values of characters of $W^\circ(\lambda)$ and its corresponding Hecke algebra. If F is split, then the sequence of subgroups $R(\lambda) \leq W^\circ(\lambda) \leq W(\lambda)$ fits the setup of 8.2. In particular, each irreducible character of $W^\circ(\lambda)$ either extends to $W(\lambda)$ or is induced from $R(\lambda)$. Moreover, there is a corresponding sequence of subalgebras of the Hecke algebra, and the same statement must be true for the representations of these algebras, by [GP00, §10.4.2]. Using this and the fact that $R(\lambda)$ and $W(\lambda)$ are products of type B and D groups, it follows that all irreducible representations of the algebras corresponding to $R(\lambda)$ and $W(\lambda)$ are defined over \mathbb{Q} . This implies that all irreducible representations of the Hecke algebra $\mathrm{End}_{\mathbb{G}}(R_{\mathbf{T}}^{\mathbb{G}}(\lambda))$ are also defined over \mathbb{Q} . When F is twisted, a similar argument applies. In particular, we have $\eta = \eta^\sigma = \eta^{(\sigma)}$ for all $\eta \in \mathrm{Irr}(W(\lambda))$ and all $\sigma \in \mathcal{G}$. This completes the proof, using Theorem 3.2. \blacksquare

Proof (of Theorem A.(i)). Let $\mathbf{G} = \mathrm{GO}(V)$ and $\sigma \in \mathcal{G}$. As in the proof of Proposition 12.2, we will just treat the case $\dim(V)$ even. After Proposition 12.2, it suffices to prove the statement for all members of a Harish-Chandra series $\mathcal{E}(\mathbf{G}^{\circ F}, \mathbf{L}, \lambda)$ with $(\mathbf{L}, \lambda) \in \mathrm{Cusp}_s(\mathbf{G}, F)$ a cuspidal pair such that $\mathbf{L} \neq \mathbf{T}$. We maintain the notation of 12.1-12.1. By Theorem 10.8, the character $\delta_{\psi, \sigma}$ is trivial. Moreover, it follows from the proof of Proposition 12.2 that $\delta_{\lambda_1, \sigma}$ is trivial on $C(\lambda_1)$. We let $m = \dim(\mathbf{S}_1)$. As $\mathbf{L} \neq \mathbf{T}$, we have $0 \leq m \leq n - 2$. Then we similarly see that $\delta_{\lambda_1, \sigma}$ is trivial on $n_{\mathbf{t}_m}$, which generates $W(\lambda_1)/W^\circ(\lambda_1)$, implying $\delta'_{\lambda, \sigma}$ must be trivial.

As in the case of Proposition 12.2, we may arrange that $\lambda_1 = 1_{\mathbf{T}_1^F} \boxtimes \cdots \boxtimes 1_{\mathbf{T}_a^F} \boxtimes \varepsilon_{\mathbf{T}_{a+1}^F} \boxtimes \cdots \boxtimes \varepsilon_{\mathbf{T}_m^F}$ for some $0 \leq a \leq m$. Let $b = m - a$. Identifying $N_{\mathbf{G}^{\circ F}}(\mathbf{L})/\mathbf{L}^F$ with the Coxeter group W_m , as in Lemma 7.6, we have the group $W^\circ(\lambda)$ is the subgroup $W_a \times W_b$ of

7.3. Using the notation of 7.7, we let $\tilde{\lambda} = \lambda_1 \boxtimes \tilde{\psi} \in \text{Irr}(\tilde{\mathbf{L}}_J^F)$ with $\tilde{\psi} \in \text{Irr}(\tilde{\mathbf{M}}_0^{\circ F})$ a character covering ψ .

By Lemma 3.4, and the description of the action of u_m in 7.7, we see that $W^\circ(\tilde{\lambda}) = R(\lambda) = W_a \times W'_b \leq W^\circ(\lambda)$ is of type $B_a \times D_b$. Hence $C(\lambda)$ is generated by u_m as in Lemma 7.6. The element u_m has even length in \mathbf{W} , so $\gamma_{\lambda, \sigma}$ is trivial by Lemma 3.6. Finally, as $W(\lambda)$ is a product of type B Weyl groups, we have $\eta^{(\sigma)} = \eta$ for all $\eta \in \text{Irr}(W(\lambda))$, so the result follows from Theorem 3.2. ■

Proof (of Theorem A.(iii)). We may again assume $\dim(V)$ is even and $F = \mathbb{F}_q \in \Gamma_q(\mathbf{G})$ is split and $\gamma = \gamma_1$ is as in 7.8. Let $(\mathbf{L}, \lambda) \in \text{Cusp}(\mathbf{G}, F)$ be a cuspidal pair and set $\tilde{\mathbf{L}} = \mathbf{L} \cdot Z(\tilde{\mathbf{G}}) \leq \tilde{\mathbf{G}}$. Suppose $\tilde{\lambda} \in \text{Irr}(\tilde{\mathbf{L}}^F)$ covers λ . Identifying $\tilde{W}(\tilde{\lambda})$ with $W(\tilde{\lambda}) \leq W(\lambda)$, we have by [Bon06, Thm. 13.9] that

$$\text{Res}_{\mathbf{G}^F}^{\tilde{\mathbf{G}}^F} \left(R_{\tilde{\mathbf{L}}}^{\tilde{\mathbf{G}}}(\tilde{\lambda})_{\tilde{\eta}} \right) = R_{\mathbf{L}}^{\mathbf{G}}(\lambda)_{\text{Ind}_{W(\tilde{\lambda})}^{W(\lambda)} \tilde{\eta}}$$

for any $\tilde{\eta} \in \text{Irr}(W(\tilde{\lambda}))$. Here we extend the map $\text{Irr}(W(\lambda)) \rightarrow \text{Irr}(\mathbf{G}^F)$, given by $\eta \mapsto R_{\mathbf{L}}^{\mathbf{G}}(\lambda)_\eta$, linearly to all class functions.

Consider first the case where $\mathbf{L} = \mathbf{T}$. As above we arrange that $\lambda = 1_{\mathbf{T}_1^F} \boxtimes \cdots \boxtimes 1_{\mathbf{T}_a^F} \boxtimes \varepsilon_{\mathbf{T}_{a+1}^F} \boxtimes \cdots \boxtimes \varepsilon_{\mathbf{T}_n^F}$ for some $1 < a < n$. Note we assume $s \neq \pm 1$. Using the above identity, as well as [MS16, 5.6], the statement holds in this case. Note, by Lusztig's classification this series contains all the characters $\chi \in \mathcal{E}(\mathbf{G}^F, s)$ satisfying the condition $\text{Inn}(\mathbf{G}, F)_\chi = \text{Inn}(\mathbf{G}, F)$. Now assume $\mathbf{L} \neq \mathbf{T}$. The graph automorphism γ_1 induces the identity on $W(\lambda)$ so the statement again follows from [MS16, 5.6]. ■

13. Galois Automorphisms: Quasi-Isolated Series of $\text{Sp}(V)$

13.1. In this section, we consider the action of \mathcal{G} on quasi-isolated series when $\mathbf{G} = \text{Sp}(V)$. As in 12.1, we need to understand the action of \mathcal{G} on principal series characters and cuspidal characters. The following treats the cuspidal case and extends [Tay18, 14.6] to the setting of Galois automorphisms.

Theorem 13.2. *Assume $\mathbf{G} = \text{Sp}(V)$ and $s \in \mathbf{G}^{*F^*}$ is a quasi-isolated semisimple element. Let $\sigma \in \mathcal{G}$ be a Galois automorphism, let $k \in \mathbb{Z}$ be an integer coprime to p such that $\xi^\sigma = \xi^k$ for all p th roots of unity $\xi \in \overline{\mathbb{Q}}^\times$, and let $\omega = (-1)^{(p-1)/2}$. If $\chi \in \mathcal{E}(\mathbf{G}^F, s)$ is a cuspidal character, then we have $\chi^\sigma = \chi$ if and only if one of the following holds:*

- (i) $s = 1$, so that χ is unipotent,
- (ii) $k \pmod{p} \in \mathbb{F}_q$ is a square.

In particular, we have $\mathcal{Q}(\chi) = \mathbb{Q}$ if $s = 1$ or q is square, and $\mathcal{Q}(\chi) = \mathbb{Q}(\sqrt{\omega p})$ otherwise.

Proof. Unipotent characters of classical groups are rational-valued, so we may assume $s \neq 1$. Assume \mathcal{O}_χ^* is the wave-front set of χ , as in [Tay18, 14.1]. Let $\dim(V) = 2n$ and let $\mu \in \mathcal{P}_1(2n)$ be the partition parameterising the unipotent class \mathcal{O}_χ^* , under the bijection in 9.1. From the proof of [Tay18, 14.4], we see that $r_{2m}(\mu) = 1$ for some $0 < m \leq n$.

Then arguments along the lines of those used in Theorem 10.6, instead appealing to Corollary 9.8 and [SFT18, Prop. 4.10], complete the proof of the first statement. The statement about character fields follows from this, combined with (3.10) and 5.5. ■

Remark 13.3. This implies that the Harish-Chandra series in 11.1 can be permuted non-trivially by the action of \mathcal{G} , unlike the case of the action of $\text{Aut}(\mathbf{G}^F)$.

Lemma 13.4. *Assume $\mathbf{G} = \text{Sp}(V)$. Let $\lambda \in \text{Irr}(\mathbf{T}^F)$ be a nontrivial character such that $\lambda^2 = 1$ and let $\sigma \in \mathcal{G}$. Then*

- (i) $\eta = \eta^\sigma = \eta^{(\sigma)}$ for every $\eta \in \text{Irr}(W(\lambda))$;
- (ii) $l(w_2)$ is odd for $1 \neq w_2 \in C(\lambda)$;
- (iii) if $q \equiv 1 \pmod{4}$, then any extension of λ to $N_{\mathbf{G}^F}(\mathbf{T})_\lambda$ is fixed by σ ; in particular, $\delta_{\lambda,\sigma} = \delta'_{\lambda,\sigma}$ is trivial;
- (iv) if $q \equiv 3 \pmod{4}$, then there is a σ -invariant extension of λ to $N_{\mathbf{G}^F}(\mathbf{T})_\lambda$ if and only if σ fixes 4th roots of unity. In particular, $\delta_{\lambda,\sigma} = \delta'_{\lambda,\sigma}$ is trivial if and only if σ fixes fourth roots of unity.

Proof. As in the proof of Proposition 12.2, we arrange that $\lambda = 1_{\mathbf{T}_1^F} \boxtimes \cdots \boxtimes 1_{\mathbf{T}_a^F} \boxtimes \varepsilon_{\mathbf{T}_{a+1}^F} \boxtimes \cdots \boxtimes \varepsilon_{\mathbf{T}_n^F}$ for some $0 \leq a < n$. Note we have $b = n - a > 0$ because λ is assumed to be nontrivial. Taking $m = a$ in 7.5, we have $W(\lambda) = \mathbf{W}_1 \times \mathbf{W}_0$ is a Weyl group of type $B_a \times B_b$. This gives (i). Letting $\tilde{\lambda} = \lambda \boxtimes 1_{\mathbf{Z}^F} \in \text{Irr}(\tilde{\mathbf{T}}^F)$, we see using Lemma 3.4 that $R(\lambda) = W(\tilde{\lambda}) = W_a \times W'_b$ and $C(\lambda) = \langle s_n \rangle$, which proves (ii).

Let $c := n_{s_n} \in N_{\mathbf{G}^F}(\mathbf{T})_\lambda$ represent $s_n \in C(\lambda)$, and let $\xi \in \mathbf{T}_n^F \cong \mathbb{F}_q^\times$ be a generator of the cyclic group. Since $c^2 = \mathbf{d}_n(-1)$ and $-1 = \xi^{(q-1)/2}$, we see that for any extension $\Lambda_\lambda := \Lambda(\lambda)$ of the linear character λ to $N_{\mathbf{G}^F}(\mathbf{T})_\lambda$, we have

$$\Lambda_\lambda(c)^2 = \Lambda_\lambda(c^2) = \varepsilon_{\mathbf{T}_n^F}(\xi^{(q-1)/2}) = \varepsilon_{\mathbf{T}_n^F}(\xi)^{(q-1)/2} = (-1)^{(q-1)/2}. \quad (13.5)$$

By combining this with Lemma 3.8, we see that if $q \equiv 1 \pmod{4}$, then all values of Λ_λ are in $\{\pm 1\}$, giving (iii). On the other hand, if $q \equiv 3 \pmod{4}$, then (13.5) implies Λ_λ must take primitive fourth roots of unity as values on c , proving (iv). ■

13.6. When $\lambda \in \text{Irr}(\mathbf{T}^F)$ is a nontrivial character satisfying $\lambda^2 = 1$, combining Lemmas 3.6 and 13.4 yields a complete description of $\gamma_{\lambda,\sigma}$ and $\delta_{\lambda,\sigma}$, which can be viewed as characters of $C(\lambda) \cong C_2$. In particular, note that these depend only on q and σ , and we obtain the following.

Corollary 13.7. *Assume $\mathbf{G} = \text{Sp}(V)$. Let $\omega \in \{\pm 1\}$ be such that $p \equiv \omega \pmod{4}$. Let $\lambda \in \text{Irr}(\mathbf{T}^F)$ be a nontrivial character such that $\lambda^2 = 1$ and let $\sigma \in \mathcal{G}$. Identifying $\text{Irr}(C(\lambda))$ with $\{\pm 1\}$, the following hold:*

- (i) if q is a square, then $\gamma_{\lambda,\sigma} \delta_{\lambda,\sigma} = 1$;
- (ii) if q is not a square, then $\gamma_{\lambda,\sigma} \delta_{\lambda,\sigma} = \alpha$, where $\alpha \in \{\pm 1\}$ is such that $\sqrt{\omega p}^\sigma = \alpha \sqrt{\omega p}$.

With this in place, we may now give a proof of Theorem B.

Proof (of Theorem B). Theorem B follows immediately from Theorem 13.2, Lemma 13.4, and Corollary 13.7, using Theorem 3.2 and Lemma 11.2. ■

14. Galois Automorphisms Relevant for the McKay–Navarro Conjecture II

14.1. We now return to the setting of Section 4 and consider those Galois automorphisms relevant to the McKay–Navarro conjecture in the context of symplectic groups. Such explicit statements will be particularly useful in future work. As in Section 4, we assume $\sigma \in \mathcal{H}$ is fixed and $r \geq 0$ is the integer such that $\zeta^\sigma = \zeta^{\ell^r}$ for all roots of unity $\zeta \in \overline{\mathbb{Q}}^\times$ of order coprime to ℓ .

Lemma 14.2. *Assume $\mathbf{G} = \mathrm{Sp}(V)$ and $\ell \neq p$ are both odd. Let $\lambda \in \mathrm{Irr}(\mathbf{T}^F)$ be a nontrivial character such that $\lambda^2 = 1$. Identifying $\mathrm{Irr}(\mathbf{C}(\lambda))$ with the group $\{\pm 1\}$, we have:*

- (i) *If q is square or r is even, then $\gamma_{\lambda,\sigma}\delta_{\lambda,\sigma} = 1$.*
- (ii) *If q is not square and r is odd, then $\gamma_{\lambda,\sigma}\delta_{\lambda,\sigma} = \left(\frac{\ell}{p}\right)$.*

In particular, note that if $\ell \mid (q-1)$, then $\gamma_{\lambda,\sigma}\delta_{\lambda,\sigma} = 1$ unless $q \equiv \ell \equiv 3 \pmod{4}$ and r is odd, in which case $\gamma_{\lambda,\sigma}\delta_{\lambda,\sigma} = \delta_{\lambda,\sigma} = -1$.

Proof. This follows from Lemma 4.3 and Corollary 13.7. ■

Lemma 14.3. *Assume $\mathbf{G} = \mathrm{Sp}(V)$ and $\ell = 2$. Let $\lambda \in \mathrm{Irr}(\mathbf{T}^F)$ be a nontrivial character such that $\lambda^2 = 1$. Then $\gamma_{\lambda,\sigma}\delta_{\lambda,\sigma} \in \mathrm{Irr}(\mathbf{C}(\lambda)) \cong \{\pm 1\}$ satisfies:*

- *If $q \equiv \pm 1 \pmod{8}$, then $\gamma_{\lambda,\sigma}\delta_{\lambda,\sigma} = 1$.*
- *If $q \equiv \pm 3 \pmod{8}$, then $\gamma_{\lambda,\sigma}\delta_{\lambda,\sigma} = (-1)^r$.*

Proof. This follows from Lemma 4.8 and Corollary 13.7. ■

Remark 14.4. We end this section with some useful remarks on specific situations, which follow from Theorem 13.2 and Lemmas 13.4 and 14.2. Let $\mathbf{G} = \mathrm{Sp}(V)$ with $\ell \neq p$ and p odd.

- (i) For $\ell \mid (q-1)$ odd, we have every member of $\mathcal{E}(\mathbf{G}^F, s)$ with $s \neq 1$ quasi-isolated is fixed by all of \mathcal{H} if and only if at least one of q or ℓ is congruent to 1 modulo 4.
- (ii) There has been much recent interest in the particular case that $\sigma := \sigma_1$ is the Galois automorphism fixing ℓ' -roots of unity and mapping ℓ -power roots of unity to their $\ell+1$ power, see, e.g., [NT19; RSFV20; NRSFV20]. In this situation, we have every member of any $\mathcal{E}(\mathbf{G}^F, s)$ with $s^2 = 1$ is fixed by σ_1 . Hence, the action of σ_1 on $\mathrm{Irr}_{\ell'}(\mathbf{G}^F)$ in this case is determined by the action on irreducible characters of general linear groups, which is well understood, see, e.g. [SV19; SFT18].

15. Automorphisms Not Lifting to Asogenies

15.1. In this section, we assume that $\mathbf{G} = \mathrm{SO}(V)$. We wish to consider the group of automorphisms $\mathbf{K}(\mathbf{G}^F)$ defined in Proposition 2.3. Recall that this group is either trivial or a cyclic group of order 2. It is non-trivial only when $Z(\mathbf{G})^F \leq \mathrm{O}^{\mathrm{p}'}(\mathbf{G}^F)$. The classification of when this happens is well known and is typically calculated using a result of Zassenhaus [Zas62, §2, Thm.] on the spinor norm. The following slightly more general statement is also well known, but we include a proof that is in keeping with the theory of algebraic groups.

Lemma 15.2. *Assume $\mathbf{G} = \mathrm{SO}(V)$ and $n = \lfloor \dim(V)/2 \rfloor$. We let ϵ be 1 if F is split and -1 if F is twisted. Let $0 \leq a < n$ be an integer and set $b = n - a$. Denote by $\mathcal{O} \subseteq \mathbf{G}$ the \mathbf{G} -conjugacy class of $s = \mathbf{d}_{a+1}(-1) \cdots \mathbf{d}_n(-1)$ so that $\dim(V_{-1}(s)) = 2b$, where $V_{-1}(s)$ is the (-1) -eigenspace of s .*

- (i) *If $s \in Z(\mathbf{G})^F$, then s is contained in $\mathrm{O}^{\mathrm{p}'}(\mathbf{G}^F)$ if and only if $q^n \equiv \epsilon \pmod{4}$,*
- (ii) *If $s \notin Z(\mathbf{G})^F$, then $\mathcal{O}' = \mathcal{O} \cap \mathrm{O}^{\mathrm{p}'}(\mathbf{G}^F)$ is a single $\mathrm{O}^{\mathrm{p}'}(\mathbf{G}^F)$ -conjugacy class and it contains s if and only if $q^b \equiv \epsilon \pmod{4}$.*

Proof. Let $\pi : \mathbf{G}_{\mathrm{sc}} \rightarrow \mathbf{G}$ be a simply connected cover of \mathbf{G} . We denote again by F a Frobenius endomorphism of \mathbf{G}_{sc} such that $F \circ \pi = \pi \circ F$. Let $\mathbf{T}_{\mathrm{sc}} = \pi^{-1}(\mathbf{T})$, an F -stable maximal torus of \mathbf{T} . The image $\pi(\mathbf{G}_{\mathrm{sc}}^F)$ is precisely $\mathrm{O}^{\mathrm{p}'}(\mathbf{G}^F)$. If $t \in \mathbf{G}^F$ then $t \in \mathrm{O}^{\mathrm{p}'}(\mathbf{G}^F)$ if and only if $\pi^{-1}(t) \cap \mathbf{G}_{\mathrm{sc}}^F \neq \emptyset$. Note that if it is non-empty then $\pi^{-1}(t) \cap \mathbf{G}_{\mathrm{sc}}^F = \pi^{-1}(t)$ because F fixes $\mathrm{Ker}(\pi)$.

First let us note that as the centraliser of a semisimple element contains a maximal torus, it follows from [DM20, 11.4.11] that any \mathbf{G}^F -conjugacy class of semisimple elements contained in $\mathrm{O}^{\mathrm{p}'}(\mathbf{G}^F)$ is a single $\mathrm{O}^{\mathrm{p}'}(\mathbf{G}^F)$ -class. Now, we claim that at most one \mathbf{G}^F -class in \mathcal{O}^F is contained in $\mathrm{O}^{\mathrm{p}'}(\mathbf{G}^F)$. Fix $\tilde{s} \in \pi^{-1}(s)$ then as $F(s) = s$ we have $\tilde{s}^{-1}F(\tilde{s}) \in \mathrm{Ker}(\pi) \leq Z(\mathbf{G}_{\mathrm{sc}})$. Recall we have a homomorphism $\gamma_s : C_{\mathbf{G}}(s) \rightarrow \mathrm{Ker}(\pi)$ given by $\gamma_s(x) = [\tilde{x}, \tilde{s}]$ where $\tilde{x} \in \pi^{-1}(x)$. This factors through an injective homomorphism $C_{\mathbf{G}}(s)/C_{\mathbf{G}}^{\circ}(s) \rightarrow \mathrm{Ker}(\pi)$, see [Bon05, Cor. 2.8]. Now assume $s' = {}^g s \in \mathbf{G}^F$ for some $g \in \mathbf{G}$ and $\tilde{s}' \in \pi^{-1}(s')$ then $\tilde{s}'^{-1}F(\tilde{s}') = \tilde{s}^{-1}F(\tilde{s})\gamma_s(g^{-1}F(g))$. From this the claim follows easily.

Let $\tilde{\iota}_{\mathrm{sc}} : \mathbf{Q}\check{\Phi} \rightarrow \mathbf{T}_{\mathrm{sc}}$ be a surjective group homomorphism as in 5.3. We use the notation for roots as in [Bou02], which is consistent with our setup. We may take $\tilde{s} = \tilde{\iota}_{\mathrm{sc}}(\frac{1}{2}\check{\omega}_a) \in \pi^{-1}(s)$ as a preimage of s , where

$$\check{\omega}_a = \begin{cases} \check{\alpha}_{a+1} + 2\check{\alpha}_{a+2} + \cdots + (b-2)\check{\alpha}_{n-2} + \frac{1}{2}((b-2)\check{\alpha}_{n-1} + b\check{\alpha}_n) & \dim(V) \text{ even,} \\ \check{\alpha}_{a+1} + 2\check{\alpha}_{a+2} + \cdots + (b-2)\check{\alpha}_{n-2} + (b-1)\check{\alpha}_{n-1} + \frac{b}{2}\check{\alpha}_n & \dim(V) \text{ odd.} \end{cases}$$

A straightforward calculation shows that if F is split then $F(\tilde{s}) = \tilde{s}$ if and only if $\frac{(q-1)b}{4} \in \mathbb{Z}$. Similarly if F is twisted then $F(\tilde{s}) = \tilde{s}$ if and only if $\frac{(q-1)b}{4} \in \frac{1}{2} + \mathbb{Z}$. One readily checks the equivalence of these conditions with those given in (i) and (ii). \blacksquare

15.3. Assume now that $K(\mathbf{G}^F)$ is non-trivial, so that $Z(\mathbf{G})^F \leq \text{Op}'(\mathbf{G}^F)$. Let $s \in \mathbf{G}^{*F^*}$ be a quasi-isolated semisimple element. If $s \in \text{Op}'(\mathbf{G}^F)$, then $K(\mathbf{G}^F)_\chi = K(\mathbf{G}^F)$ for all $\chi \in \mathcal{E}(\mathbf{G}^F, s)$, using Proposition 2.4. Now assume $s \notin \text{Op}'(\mathbf{G}^F)$ and let $z \in Z(\mathbf{G}^*)$. If sz is not \mathbf{G}^* -conjugate to s , then no character in $\mathcal{E}(\mathbf{G}^F, s)$ is $K(\mathbf{G}^F)$ -invariant. Hence, the only case we need to consider is when $n = 2m$ and $C_{\mathbf{G}^*}^\circ(s)$ has type $D_m \times D_m$. In this case, we show $\mathcal{E}(\mathbf{G}^F, s)$ is $K(\mathbf{G}^F)$ -invariant.

Theorem 15.4. *Assume $\mathbf{G} = \text{SO}(V)$ and $\dim(V)$ is even. Let $s \in \mathbf{G}^{*F^*}$ be a quasi-isolated semisimple element and $\chi \in \mathcal{E}(\mathbf{G}^F, s)$ be a cuspidal character. Then*

- (i) $s \in \text{Op}'(\mathbf{G}^{*F^*})$ except when $q \equiv 1 \pmod{4}$ and $\text{SO}(V_{-1}^*(s))^{F^*}$ is of type ${}^2D_m(q)$.

Assume now that $Z(\mathbf{G})^F \leq \text{Op}'(\mathbf{G}^F)$, so that $K(\mathbf{G}^F)$ is non-trivial. Then

- (ii) χ is $K(\mathbf{G}^F)$ -invariant if and only if either $s \in \text{Op}'(\mathbf{G}^{*F^*})$ or $\dim(V_1^*(s)) = \dim(V_{-1}^*(s))$.

Proof. (i). By the classification of unipotent characters, a quasisimple group of type D_n , resp., 2D_n , has a cuspidal unipotent character if and only n is an even, resp., odd, square, see [Lus78, Table II]. This is now a straightforward check using Lemma 15.2.

(ii). Assume $s \notin \text{Op}'(\mathbf{G}^{*F^*})$. Then the central character $\omega_\chi = \omega_s$ is non-trivial by (iii) of Proposition 2.4. It follows from Corollary 2.5 that if sz is not \mathbf{G}^{*F^*} -conjugate to s , with $1 \neq z \in Z(\mathbf{G}^*)^{F^*}$, then $\chi^\sigma \neq \chi$. Now assume one of the conditions in (ii) holds. Then $\mathcal{E}(\mathbf{G}^F, s)$ is $K(\mathbf{G}^F)$ -invariant. Any GGGC is clearly $K(\mathbf{G}^F)$ -invariant, as it is non-zero only on p -elements. Hence, the statement follows as in the proof of Theorem 10.6. ■

Remark 15.5. This shows that when $q \equiv -1 \pmod{4}$, we have all cuspidal characters contained in a series $\mathcal{E}(\mathbf{G}^F, s)$ with s quasi-isolated are $K(\mathbf{G}^F)$ -invariant.

15.6. Now, let $\mathbf{G} = \text{GO}(V)$ with $\dim(V)$ even. We next wish to describe the action of $K(\mathbf{G}^{\circ F})$ on the characters of quasi-isolated series preserved by $K(\mathbf{G}^{\circ F})$. Toward this end, we begin with an analogue of Proposition 6.6 for $K(\mathbf{G}^{\circ F})$. Note that since $\text{Op}'(\mathbf{G}^F) = \text{Op}'(\mathbf{G}^{\circ F})$, we may identify $K(\mathbf{G}^{\circ F})$ with a subgroup of $K(\mathbf{G}^F)$. We keep the notation of 6.2 and 6.3.

Proposition 15.7. *Let $\mathbf{G} = \text{GO}(V)$ with $\dim(V)$ even. Then for any unipotent element $u \in \mathbf{G}^{\circ F}$, we have $\widehat{\Gamma}_{u,2}$ is fixed by $K(\mathbf{G}^F)$.*

Proof. Let $\tau \in K(\mathbf{G}^F)$ be nontrivial. Since $\mathbf{U}(\lambda, -1)^F$ is comprised of p -elements and the Sylow 2-subgroup $S_{u,\lambda}$ of $C_{\mathbf{G}}(\lambda)_{\zeta_{u,\lambda}}^F$ contains $Z(\mathbf{G}^F)$, we see that the subgroup $\mathbf{U}(\lambda, -1)^F \cdot S_{u,\lambda}$ is preserved by τ . Further, $\widehat{\zeta}_{u,\lambda}^\tau$ is an extension of $\zeta_{u,\lambda}$ satisfying $o(\widehat{\zeta}_{u,\lambda}^\tau) = o(\widehat{\zeta}_{u,\lambda}) = o(\zeta_{u,\lambda})$. Hence $\widehat{\zeta}_{u,\lambda}^\tau = \widehat{\zeta}_{u,\lambda}$ by the uniqueness of this property. Then we have

$$\widehat{\Gamma}_{u,2}^\tau = \text{Ind}_{\mathbf{U}(\lambda, -1)^F S_{u,\lambda}}^{\mathbf{G}^F}(\widehat{\zeta}_{u,\lambda}^\tau) = \text{Ind}_{\mathbf{U}(\lambda, -1)^F S_{u,\lambda}}^{\mathbf{G}^F}(\widehat{\zeta}_{u,\lambda}) = \widehat{\Gamma}_{u,2},$$

as claimed. ■

Theorem 15.8. *Let $\mathbf{G} = \text{GO}(V)$ with $\dim(V)$ even, and let $s \in \mathbf{G}^{\circ * F^*}$ be a quasi-isolated semisimple element and $\chi \in \mathcal{E}(\mathbf{G}^{\circ F}, s)$ cuspidal such that χ is $K(\mathbf{G}^{\circ F})$ -invariant. Then χ extends to a $K(\mathbf{G}^F)$ -invariant cuspidal character in $\text{Irr}(\mathbf{G}^F)$.*

Proof. This follows from 15.7, using the same argument as in Theorem 10.6. \blacksquare

15.9. We next consider the action of $K(\mathbf{G}^{\circ F})$ on pairs $(\mathbf{L}, \lambda) \in \text{Cusp}_s(\mathbf{G}^{\circ}, F)$ for s quasi-isolated. Note that since $Z(\mathbf{G})^F \leq \mathbf{L}^F$, the group \mathbf{L}^F is preserved by $K(\mathbf{G}^{\circ F})$. Further, recall that using Corollary 2.5, $\mathcal{E}(\mathbf{G}^{\circ F}, s)$ is fixed by $K(\mathbf{G}^{\circ F})$ if and only if $s \in \text{OP}'(\mathbf{G}^F)$ or $\dim(V_1(s)) = \dim(V_{-1}(s))$. Hence, we wish to determine when the series $\mathcal{E}(\mathbf{G}^{\circ F}, \mathbf{L}, \lambda) \subseteq \mathcal{E}(\mathbf{G}^{\circ F}, s)$ are also preserved in this situation. We keep the notation of Section 11.

Theorem 15.10. *Let $\mathbf{G} = \text{GO}(V)$ with $\dim(V)$ even, and let $s \in \mathbf{G}^{\circ * F^*}$ be a quasi-isolated semisimple element such that $\mathcal{E}(\mathbf{G}^{\circ F}, s)$ is preserved by $K(\mathbf{G}^{\circ F})$. Let $(\mathbf{L}, \lambda) \in \text{Cusp}_s(\mathbf{G}^{\circ}, F)$. Then*

- (i) *If $\epsilon = 1$ or $\mathbf{L} \neq \mathbf{T}$, the character λ is invariant under $K(\mathbf{G}^{\circ F})$ if and only if $s \in \text{OP}'(\mathbf{G}^F)$.*
- (ii) *If $\epsilon = -1$ and $\mathbf{L} = \mathbf{T}$, the character λ is invariant under $K(\mathbf{G}^{\circ F})$ if and only if at least one of the following holds*
 - $s \in \text{OP}'(\mathbf{G}^F)$ or
 - $q \equiv 1 \pmod{4}$ and λ is trivial on \mathbf{T}_n^F .

(iii) *Every character of $\mathcal{E}(\mathbf{G}^{\circ F}, \mathbf{L}, \lambda)$ is fixed by $K(\mathbf{G}^{\circ F})$ if and only if λ is fixed by $K(\mathbf{G}^{\circ F})$.*

Proof. Keep the notation of Section 12, so $\lambda = \lambda_1 \boxtimes \psi$, with λ_1 as in the proof of Proposition 12.2 or Theorem A(i), depending on whether $\mathbf{L} = \mathbf{T}$ or $\mathbf{L} \neq \mathbf{T}$. Let $\sigma \in K(\mathbf{G}^{\circ F})$ be nontrivial.

We begin by showing the “only if” direction of (i) and (ii). For this, assume $s \notin \text{OP}'(\mathbf{G}^{*F^*})$ and that $\dim(V_1(s)) = \dim(V_{-1}(s))$. Write $2b = \dim(V_1(s)) = \dim(V_{-1}(s))$ and let $s = s_1 s_0$ with $s_i \in \mathbf{M}_i^{*F^*}$. For $i \in \{0, 1\}$, write $2a_i := \dim(V_1(s_i))$ and $2b_i := \dim(V_{-1}(s_i))$. Note that by Lemma 15.2, we have $q^b \equiv -\epsilon \pmod{4}$.

We claim that for each $g_0 \in \mathbf{M}_0^{\circ F}$, there is $g_1 \in \mathbf{S}_1^F$ such that $g_1 g_0$ is not in $\text{OP}'(\mathbf{G}^F)$. Note that we would otherwise have $\mathbf{S}_1^F \subseteq \text{OP}'(\mathbf{G}^F)$. If $q \equiv -1 \pmod{4}$, Theorem 15.4 yields that $s_0 \in \text{OP}'(\mathbf{M}_0^{*F^*})$, so our assumption $s \notin \text{OP}'(\mathbf{G}^{*F^*})$ implies that $s_1 \notin \text{OP}'(\mathbf{G}^{*F^*})$. Viewing s_1 instead as an element of $\mathbf{G}^F \cong \mathbf{G}^{*F^*}$, we therefore see $\mathbf{S}_1^F \not\subseteq \text{OP}'(\mathbf{G}^F)$. Similarly, if $q \equiv 1 \pmod{4}$, then $\epsilon = -1$ and by Lemma 15.2, the element $\mathbf{d}_m(-1)$, for example, is an element of \mathbf{S}_1^F but not $\text{OP}'(\mathbf{G}^F)$. This proves the claim.

First, assume $\mathbf{T} \neq \mathbf{L}$ and recall that λ_1 may be chosen to be trivial on \mathbf{T}_i^F for $1 \leq i \leq \alpha_1$ and $\varepsilon_{\mathbf{T}_i^F}$ for $\alpha_1 + 1 \leq i \leq m$. Write z, z_0 , and z_1 for the generators of $Z(\mathbf{G}^F)$, $Z(\mathbf{M}_0^F)$, and $Z(\mathbf{M}_1^F)$, respectively. Let $g_0 \in \mathbf{M}_0^{\circ F}$ such that $\psi(g_0) \neq 0$, and let $g_1 \in \mathbf{S}_1^F$ such that $g = g_1 g_0 \in \mathbf{L}^F$ is not a member of $\text{OP}'(\mathbf{G}^F)$. Then

$$\lambda^\sigma(g) = \lambda(gz) = \psi(g_0 z_0) \lambda_1(g_1 z_1) = \lambda(g) \omega_{s_0}(z_0) (-1)^{(q-1)b_1/2}.$$

Using Lemma 15.2, we see

$$\omega_{s_0}(z_0) = \begin{cases} (-1)^{b_0(q-1)/2} & \text{if } \epsilon = 1, \\ (-1)^{b_0(q-1)/2+1} & \text{if } \epsilon = -1. \end{cases}$$

Then since $q^b \not\equiv \epsilon \pmod{4}$ and $\lambda(g) \neq 0$, we have $\lambda^\sigma(g) \neq \lambda(g)$.

In the case $\mathbf{T} = \mathbf{L}$, we take $\lambda = \lambda_1$, $m = n$, $a_0 = b_0 = 0$, and $a_1 = b_1 = b$. If $\epsilon = 1$, λ is therefore of the form above. If $\epsilon = -1$, the restriction of λ to \mathbf{T}_n^F may be either trivial or $\epsilon_{\mathbf{T}_n^F}$. Let $g \in \mathbf{T}^F$ such that $g \notin \text{Op}'(\mathbf{G}^F)$. Since λ is linear, we know $\lambda(g) \neq 0$, and we have

$$\lambda^\sigma(g) = \lambda(gz) = \begin{cases} \lambda(g)(-1)^{(q-1)(b-1)/2} & \text{if } \epsilon = -1 \text{ and } \lambda \text{ is trivial on } \mathbf{T}_n^F, \\ \lambda(g)(-1)^{(q-1)(b-1)/2}(-1)^{(q-\epsilon)/2} & \text{otherwise.} \end{cases}$$

If $\epsilon = 1$ or λ is nontrivial on \mathbf{T}_n^F , we therefore have $\lambda^\sigma \neq \lambda$, since $q^b \not\equiv \epsilon \pmod{4}$. This completes the “only if” direction of (i). If $\epsilon = -1$ and λ is trivial on \mathbf{T}_n^F , we have $\lambda^\sigma = \lambda$ if and only if $q \equiv 1 \pmod{4}$ or b is odd, giving (ii).

We now show the “if” direction of (iii). That is, we claim that if $\lambda^\sigma = \lambda$, then $\chi^\sigma = \chi$ for each $\chi \in \mathcal{E}(\mathbf{G}^{\circ F}, \mathbf{L}, \lambda)$. Recall that χ may be written $\chi = \mathbf{R}_L^G(\lambda)_\eta$ for some $\eta \in \text{Irr}(W^\circ(\lambda))$, as in 3.1. Let Λ be an extension map with respect to $\mathbf{L}^F \triangleleft \mathbf{N}_{\mathbf{G}^{\circ F}}(\mathbf{L})$, and let $\delta_{\lambda, \sigma} \in \text{Irr}(W^\circ(\lambda))$ be such that $\Lambda(\lambda)^\sigma = \delta_{\lambda, \sigma} \Lambda(\lambda^\sigma) = \delta_{\lambda, \sigma} \Lambda(\lambda)$. Recall that we may write $\Lambda(\lambda) = \text{Res}_{\mathbf{N}_{\mathbf{G}^{\circ F}}(\mathbf{L})_\lambda}^{\mathbf{N}_{\mathbf{G}^F}(\mathbf{L})_\lambda} (\Lambda_1(\lambda_1) \boxtimes \Lambda_0(\psi))$ as in Lemma 11.4. Now, calculations exactly as those above for λ yield that $\Lambda_1(\lambda_1) \boxtimes \Lambda_0(\psi)$ is fixed by σ as well, so $\delta_{\lambda, \sigma} = 1$. Then by [MS16, Theorem 4.6], we have $(\mathbf{R}_L^G(\lambda)_\eta)^\sigma = \mathbf{R}_L^G(\lambda^\sigma)_{\eta^\sigma \cdot \delta_{\lambda, \sigma}^{-1}} = \mathbf{R}_L^G(\lambda)_{\eta^\sigma}$. But note that $\eta^\sigma = \eta$ for all $\eta \in \text{Irr}(W^\circ(\lambda))$, since $Z(\mathbf{G}^{\circ F}) \leq \mathbf{L}^F$, completing the claim.

Recall that by Proposition 2.4, every $\chi \in \mathcal{E}(\mathbf{G}^{\circ F}, s)$ is fixed by $K(\mathbf{G}^{\circ F})$ if $s \in \text{Op}'(\mathbf{G}^F)$. Hence to complete the proof, it now suffices to show that (\mathbf{L}, λ) and $(\mathbf{L}, \lambda)^\sigma$ define distinct Harish-Chandra series when $\lambda \neq \lambda^\sigma$. Note that (\mathbf{L}, λ) is moved by σ if and only if λ is. We claim $(\mathbf{L}, \lambda^\sigma) \neq (\mathbf{L}, \lambda)^w$ for any $w \in \mathbf{W}^{\circ F}$. Otherwise, we have $w \in \mathbf{N}_{\mathbf{G}^{\circ F}}(\mathbf{L})$ and $w\sigma^{-1}$ fixes λ . But from the structure of $\mathbf{N}_{\mathbf{G}^{\circ F}}(\mathbf{L})$ and λ , we see the action of w on λ_1 does not affect the number of i such that $\text{Res}_{\mathbf{T}_i^F}^{\mathbf{S}_i^F} \lambda_1$ is of the form $\epsilon_{\mathbf{T}_i^F}$. Considering the calculations above, we see that $\lambda^{w\sigma^{-1}} = \lambda$ is therefore impossible if $\lambda^\sigma \neq \lambda$. It follows that the series $\mathcal{E}(\mathbf{G}^{\circ F}, \mathbf{L}, \lambda)$ is preserved by σ if and only if λ is fixed by σ . ■

16. Reductions to the Quasi-Isolated Case

16.1. The following discussion, aimed at clarifying the equivariance statement of [Tay18, Thm. 9.5], applies to any connected reductive algebraic group with no restriction on p . We also take this opportunity to upgrade this discussion to incorporate the action of the Galois group \mathcal{G} . However, we first need to define a dual action of the Galois group on semisimple elements of \mathbf{G}^* .

Recall that we fixed an isomorphism $\iota : \mathbb{Z}_{(p)}/\mathbb{Z} \rightarrow \mathbb{F}^\times$ in 5.3 and an injective homomorphism $\jmath : \mathbb{Q}/\mathbb{Z} \rightarrow \overline{\mathbb{Q}}^\times$ in 3.9. The composition $\kappa = \jmath \circ \iota^{-1}$ is an injective group homomorphism $\mathbb{F}^\times \rightarrow \overline{\mathbb{Q}}^\times$. The image $\kappa(\mathbb{F}^\times) \leq \overline{\mathbb{Q}}^\times$ is simply the group of all roots of unity whose order is coprime to p . This is preserved by the action of \mathcal{G} on $\overline{\mathbb{Q}}$, so identifying \mathbb{F}^\times with its image under κ yields an automorphic action of \mathcal{G} on \mathbb{F}^\times .

16.2. Let \check{X}^* be the cocharacter group of the torus \mathbf{T}^* . Recall that we have a group isomorphism $\mathbb{F}^\times \otimes_{\mathbb{Z}} \check{X}^* \rightarrow \mathbf{T}^*$, as in 5.3, that on simple tensors is given by $k \otimes \gamma \mapsto \gamma(k)$.

Under this isomorphism, the action of \mathcal{G} on \mathbb{F}^\times defines an automorphic action \mathcal{G} on \mathbf{T}^* . We denote the resulting homomorphism $\mathcal{G} \rightarrow \text{Aut}(\mathbf{T}^*)$ by $\sigma \mapsto \sigma^*$ and say σ^* is *dual* to σ . To be specific, assume $s \in \mathbf{T}^*$ is an element of order $d > 0$ and let $\zeta_d = \tilde{\jmath}(\frac{1}{d})$ be a primitive d th root of unity. If $\sigma \in \mathcal{G}$, then $\sigma(\zeta_d) = \zeta_d^k$ for some integer $k \in \mathbb{Z}$ coprime to d and we have $\sigma^*(s) = s^k$.

We will freely use the notation of [Tay18], in particular that of [Tay18, §8], but we maintain our preference here for right over left actions. Moreover, the reference tori $(\mathbf{T}_0, \mathbf{T}_0^*)$ of [Tay18] will simply be denoted by $(\mathbf{T}, \mathbf{T}^*)$ to be in keeping with our notation here. Now fix a semisimple element $s \in \mathbf{T}^*$ with $\mathbb{T}_{\mathbf{W}^*}(s, \mathbb{F}^*) \neq \emptyset$ and a coset $\mathfrak{a} \in \mathcal{A}_{\mathbf{W}^*}(s, \mathbb{F}^*) = \mathbf{W}^{\circ}(s) \setminus \mathbb{T}_{\mathbf{W}^*}(s, \mathbb{F}^*)$. After possibly replacing s with a \mathbf{W}^* -conjugate, we will assume that the Levi cover of $\mathbf{C}_{\mathbf{G}^*}(s)$, defined as in [Tay18, Def. 8.11], is standard of the form \mathbf{L}_I^* for some subset $I \subseteq \Delta$.

Let X^* be the character group of \mathbf{T}^* and set $\mathbb{R}X^* = \mathbb{R} \otimes_{\mathbb{Z}} X^*$. Given a vector $v \in \mathbb{R}X^*$, we denote by $[v] = \{kv \mid 0 < k \in \mathbb{R}\}$ the positive half-line spanned by v . Now let $\text{Aso}^*(\mathbf{G}, \mathbb{F}) \leq \text{Aso}(\mathbf{G}, \mathbb{F})$ be the subgroup stabilising the pair (\mathbf{T}, \mathbf{B}) . If $\sigma^* \in \text{Aso}(\mathbf{G}^*, \mathbb{F}^*)$ is dual to $\sigma \in \text{Aso}^*(\mathbf{G}, \mathbb{F})$, then σ^* defines an automorphism of \mathbf{W}^* and a permutation of the set of half-lines $[\Delta^*]$ of the simple roots. To have a uniform notation, we assume that if $\sigma \in \mathcal{G}$, then the dual σ^* induces the identity on \mathbf{W}^* and $[\Delta^*]$. We use the same conventions for σ itself.

16.3. We set $\mathcal{A}(\mathbf{G}, \mathbb{F}) = \text{Aso}(\mathbf{G}, \mathbb{F}) \times \mathcal{G}$ and $\mathcal{A}^*(\mathbf{G}, \mathbb{F}) = \text{Aso}^*(\mathbf{G}, \mathbb{F}) \times \mathcal{G}$. Moreover, we denote by $\mathcal{A}^*(\mathbf{G}, \mathbb{F})_{s, \mathfrak{a}} \leq \mathcal{A}^*(\mathbf{G}, \mathbb{F})$ the subgroup stabilising the series $\mathcal{E}_0(\mathbf{G}^{\mathbb{F}}, s, \mathfrak{a})$. If $\sigma \in \mathcal{A}^*(\mathbf{G}, \mathbb{F})$, then $\mathcal{E}_0(\mathbf{G}^{\mathbb{F}}, s, \mathfrak{a})^\sigma = \mathcal{E}_0(\mathbf{G}^{\mathbb{F}}, \sigma^*(s), \sigma^*(\mathfrak{a}))$. Note that $\sigma^*(\mathfrak{a}) = \mathfrak{a}$ if $\sigma \in \mathcal{G}$. Hence, if $\sigma \in \mathcal{A}^*(\mathbf{G}, \mathbb{F})_{s, \mathfrak{a}}$ then there exists an element $\chi^* \in \mathbf{W}^*$ such that

$$(s, \mathfrak{a}) = ({}^{\chi^*} \sigma^*(s), \chi^* \sigma^*(\mathfrak{a}) \mathbb{F}^*(\chi^{*-1})).$$

There is then a unique element $y^* \in \mathbf{W}_I^*$ such that $[y^* \chi^* \sigma^*(I^*)] = [I^*]$. We set $z_\sigma^* = y^* \chi^*$. The choice of χ^* above is unique up to multiplying by an element of $\mathbf{W}^*(s) \leq \mathbf{W}_I^*$. Therefore, the element z_σ^* is uniquely determined by σ .

The coset $\mathbf{W}_I^* \mathfrak{a}$ contains a unique element $w_1^* \in \mathbb{T}_{\mathbf{W}^*}(I^*, \mathbb{F}^*)$, and this element is fixed in the sense that $z_\sigma^* \sigma^*(w_1^*) \mathbb{F}^*(z_\sigma^{*-1}) = w_1^*$. Dually, we have $\sigma^{-1}(z_\sigma(I)) = I$ and $z_\sigma^{-1} \mathbb{F}(\sigma^{-1}(w_1) z_\sigma) = \mathbb{F}(w_1)$. As in [Tay18, Prop. 9.2], there exists an element $n_\sigma \in \mathbf{N}_{\mathbf{G}}(\mathbf{T}_0)$ representing z_σ such that $n_\sigma^{-1} \mathbb{F}(\sigma^{-1}(n_{w_1}) n_\sigma) = \mathbb{F}(n_{w_1})$. The choice for n_σ is unique up to multiplication by an element of $\mathbf{T}^{\mathbb{F} w_1}$.

As $\text{Aso}(\mathbf{G}, \mathbb{F}) = \text{Inn}(\mathbf{G}^{\mathbb{F}}) \text{Aso}^*(\mathbf{G}, \mathbb{F})$, the natural map $\text{Aso}^*(\mathbf{G}, \mathbb{F}) \rightarrow \text{Out}(\mathbf{G}, \mathbb{F})$ is surjective. We set $\mathcal{O}(\mathbf{G}, \mathbb{F}) = \mathcal{A}(\mathbf{G}, \mathbb{F}) / \text{Inn}(\mathbf{G}^{\mathbb{F}}) \cong \text{Out}(\mathbf{G}, \mathbb{F}) \times \mathcal{G}$. A straightforward calculation shows that we have a well-defined group homomorphism $\mathcal{O}(\mathbf{G}, \mathbb{F})_{s, \mathfrak{a}} \rightarrow \mathcal{O}(\mathbf{L}_I, \mathbb{F} w_1)$ given by $\sigma \mapsto \sigma n_\sigma$, where $\mathcal{O}(\mathbf{G}, \mathbb{F})_{s, \mathfrak{a}} \leq \mathcal{O}(\mathbf{G}, \mathbb{F})$ is the subgroup preserving the series $\mathcal{E}_0(\mathbf{G}^{\mathbb{F}}, s, \mathfrak{a})$. The following makes [Tay18, Thm. 9.5] more precise.

Theorem 16.4. *Let $\mathfrak{a}_1 = \mathfrak{a} w_1^{*-1} \in \mathcal{A}_{\mathbf{W}_I^*}(s, w_1^* \mathbb{F}^*)$. Then we have a bijection $(-1)^{l(w_1)} \mathbf{R}_{I, w_1}^{\mathbf{G}} : \mathcal{E}_0(\mathbf{L}_I^{\mathbb{F} w_1}, s, \mathfrak{a}_1) \rightarrow \mathcal{E}_0(\mathbf{G}^{\mathbb{F}}, s, \mathfrak{a})$ which is $\mathcal{O}(\mathbf{G}, \mathbb{F})_{s, \mathfrak{a}}$ -equivariant with respect to the homomorphism $\mathcal{O}(\mathbf{G}, \mathbb{F})_{s, \mathfrak{a}} \rightarrow \mathcal{O}(\mathbf{L}_I, \mathbb{F} w_1)$ defined above. In particular, $\mathbf{R}_{I, w_1}^{\mathbf{G}}(\psi)^\sigma = \mathbf{R}_{I, w_1}^{\mathbf{G}}(\psi^{\sigma n_\sigma})$ for*

any $\psi \in \mathcal{E}_0(\mathbf{L}_I^{\text{Fw}_1}, s, \alpha_1)$ and $\sigma \in \mathcal{A}(\mathbf{G}, F)$.

Proof. In the case of Galois automorphisms, we use [DM20, Cor. 8.1.6, Prop. 9.1.6] to conclude that $\mathbf{R}_{I, \text{w}_1}^{\mathbf{G}}$ is \mathcal{G} -equivariant. We can now argue as in [Tay18, Thm. 9.5]. \blacksquare

Proofs of Theorems C and D

16.5. We now consider the above discussion in the context of $\mathbf{G} = \text{SL}(V \mid \mathcal{B})$. Up to \mathbf{W}^* -conjugacy, we can assume that $s = s_1 s_0 \in \mathbf{T}^*$ where, for some $1 \leq m \leq n$, we have $s_1 = \mathbf{d}_1(\xi_1) \cdots \mathbf{d}_m(\xi_m)$ and $s_0 = \mathbf{d}_{m+1}(\xi_{m+1}) \cdots \mathbf{d}_n(\xi_n)$ with $\xi_i^2 \neq 1$ for all $1 \leq i \leq m \leq n$ and $\xi_i^2 = 1$ for all $m < i \leq n$. We let $\mathbf{M}_1^* \times \mathbf{M}_0^*$ be a group determined by the integer m , as in 7.5. Then the Levi cover of $\mathbf{C}_{\mathbf{G}^*}(s)$ is a standard Levi subgroup $\mathbf{L}_I^* = \mathbf{L}_{J_1}^* \times \mathbf{L}_{J_0}^*$ with $I = J_1 \sqcup J_0 \subseteq \Delta$ and $\mathbf{L}_{J_1}^* = \mathbf{C}_{\mathbf{M}_1^*}(s)$ and $\mathbf{L}_{J_0}^* = \mathbf{M}_0^{*\circ}$.

It is clear to see that we have a bijection $\mathbf{T}_{\mathbf{W}_1^*}(s_1, F^*) \times \mathbf{T}_{\mathbf{W}_0^*}(s_0, F^*) \rightarrow \mathbf{T}_{\mathbf{W}^*}(s, F^*)$, and in this way we get a bijection $\mathcal{A}_{\mathbf{W}_0^*}(s_0, F^*) \rightarrow \mathcal{A}_{\mathbf{W}^*}(s, F^*)$. We have a unique decomposition $w_1^* = v_1^* v_0^*$ with $v_i^* \in \mathbf{T}_{\mathbf{W}_i^*}(s_i, F^*)$ and we get a corresponding coset $b_i \in \mathcal{A}_{\mathbf{W}_i^*}(s_i, F^*)$ containing v_i^* . It is easy to see that v_i^* is the unique element of the coset $\mathbf{W}_{J_i}^* v_i^*$ contained in $\mathbf{T}_{\mathbf{W}_i^*}(J_i^*, F^*)$. As in the proof of [Tay18, Thm. 16.2], we have a tensor product decomposition

$$\mathcal{E}_0(\mathbf{L}_I^{\text{Fw}_1}, s, \alpha_1) = \mathcal{E}_0(\mathbf{L}_{J_1}^{\text{Fv}_1}, s_1, b_1) \boxtimes \mathcal{E}_0(\mathbf{L}_{J_0}^{\text{Fv}_0}, s_0, b_0). \quad (16.6)$$

16.7. To use Theorem 16.4, we need to describe the image of the homomorphism $\mathcal{O}(\mathbf{G}, F)_{s, \alpha} \rightarrow \mathcal{O}(\mathbf{L}_I, \text{Fw}_1)$, which we do now. For any $\sigma \in \mathcal{A}^*(\mathbf{G}, F)_{s, \alpha}$ we have $\sigma^*(s_0) = s_0$. Hence, it is clear that the element z_σ^* defined in 16.3 is contained in the subgroup \mathbf{W}_1^* . Thus the element n_σ may be chosen inside the group $\mathbf{N}_{\mathbf{M}_1}(\mathbf{S}_1)$, where $\mathbf{S}_1 \leq \mathbf{M}_1$ is the maximal torus defined in 7.5. If $\mathbf{G} = \text{SO}(V)$, then the Weyl group lifts in \mathbf{G} and we may, in fact, choose $n_\sigma = n_{z_\sigma}$. If $\mathbf{G} = \text{Sp}(V)$ then this is not necessarily the case.

If $t \in \mathbf{G}^{*F^*}$ is a semisimple element, then there exists $g \in \mathbf{G}^*$ and $s \in \mathbf{T}^*$ such that $t = {}^g s$. If $g^{-1} F^*(g) \in \mathbf{T}_{\mathbf{G}^*}(s, F^*)$ maps onto $\alpha^* \in \mathcal{A}_{\mathbf{W}^*}(s, F^*)$, then $\mathcal{E}(\mathbf{G}^F, t) = \mathcal{E}_0(\mathbf{G}^F, s, \alpha)$, see [Tay18, Lem. 8.6]. Using this identification, it is easy to translate the statements of Theorems C and D into corresponding statements in the language used above. It is these translated statements that we prove below. We leave the translation to the reader.

Proof (of Theorem C). One calculates easily using 16.7 that the image of $\mathcal{G}_{s, \alpha} := \mathcal{G} \cap \mathcal{O}(\mathbf{G}, F)_{s, \alpha}$ inside $\mathcal{O}(\mathbf{L}_I, \text{Fw}_1)$ is contained in the subgroup $\mathcal{O}(\mathbf{L}_{J_1}, \text{Fv}_1)_{s_1, b_1} \leq \mathcal{O}(\mathbf{L}_I, \text{Fw}_1)$. Theorem 16.4 implies that if $\chi \in \mathcal{E}_0(\mathbf{G}^F, s, \alpha)$, then $\chi = \pm \mathbf{R}_{I, \text{w}_1}(\psi)$ for a unique $\psi = \psi_1 \boxtimes \psi_0 \in \mathcal{E}_0(\mathbf{L}_I^{\text{Fw}_1}, s, \alpha_1)$. The equivariance of this bijection implies that we need only calculate the stabiliser of ψ under the image of $\mathcal{G}_{s, \alpha}$. The group \mathbf{L}_{J_1} is a direct product of general linear groups so a unicity of multiplicities argument shows that ψ_1 is $\mathcal{O}(\mathbf{L}_{J_1}, \text{Fv}_1)_{s_1, b_1}$ -invariant, see [SFT18, Prop. 3.10] for instance. It follows that $(\mathcal{G}_{s, \alpha})_\chi = (\mathcal{G}_{s, \alpha})_{\psi_0}$ so $\mathbf{Q}(\chi) = \mathbf{Q}_s(\psi_0)$. The statement now follows from Theorems A and B, together with the fact that the fields \mathbf{Q}_s and $\mathbf{Q}(\psi_0) \subseteq \mathbf{Q}(\mathfrak{j}(\frac{1}{p}))$ are linearly disjoint since p and d are coprime. \blacksquare

Remark 16.8. Let $\text{Cl}_{\text{ss}}(\mathbf{G}^{*F^*})$ be the set of semisimple conjugacy classes of \mathbf{G}^{*F^*} . We define an action of \mathcal{G} on $\text{Cl}_{\text{ss}}(\mathbf{G}^{*F^*})$ by setting $\sigma^*([{}^g s]) = [{}^g \sigma^*(s)]$, where $s \in \mathbf{T}^*$ and $g \in \mathbf{G}^*$ are such that ${}^g s \in \mathbf{G}^{*F^*}$. One easily checks this is well-defined, given the action $\sigma^*(s)$ via power maps as in 16.2. The stabiliser $\mathcal{G}_s \leq \mathcal{G}$ of $\mathcal{E}(\mathbf{G}^F, s)$ is then the stabiliser of the class $[s] \in \text{Cl}_{\text{ss}}(\mathbf{G}^{*F^*})$ under this action.

Proof (of Theorem D). (i). If $s_0 = 1$, then the series $\mathcal{E}_0(\mathbf{G}^F, s, \mathfrak{a})$ is not γ -invariant because the \mathbf{W}^* -orbit of s is not γ^* -stable, so we can assume that $s_0 \neq 1$. In this case, we have $\gamma^*(s) = s$. The set $\mathcal{A}_{\mathbf{W}^*}(s, F^*)$ has size 2 so clearly $\gamma^*(\mathfrak{a}) = \mathfrak{a}$ as γ^* fixes the coset $\mathbf{W}^{*\circ}(s) \in \mathcal{A}_{\mathbf{W}^*}(s, F^*)$.

The following argument can be extrapolated from [Lus84, §6.21], see also [DM90]. We defer to [Lus84; DM90] for the details. The coset $\mathfrak{a}_1 = \mathfrak{a}w_1^{*-1}$ contains a unique element of minimal length, which we denote by $w_{\mathfrak{a}_1}^* \in \mathfrak{a}_1$. The element $w_{\mathfrak{a}}^* = w_{\mathfrak{a}_1}^* w_1^* \in \mathfrak{a}$ is then the corresponding minimal length element in \mathfrak{a} .

We pick a Jordan decomposition $J_{s, \mathfrak{a}_1}^{\mathbf{L}_I, Fw_1} : \mathcal{E}_0(\mathbf{L}_I^{Fw_1}, s, \mathfrak{a}) \rightarrow \mathcal{E}_0(\mathbf{C}_{\mathbf{L}_I^*}(s)^{w_{\mathfrak{a}_1}^* F^*}, 1)$ so that

$$J_{s, \mathfrak{a}_1}^{\mathbf{L}_I, Fw_1}(\mathbf{R}_{x^* w_{\mathfrak{a}_1}^*}^{\mathbf{L}_I, Fw_1}(s)) = (-1)^{l(w_{\mathfrak{a}_1}^*)} \mathbf{R}_{x^*}^{\mathbf{C}_{\mathbf{G}^*}(s), w_{\mathfrak{a}_1}^* w_1^* F^*}(1)$$

for all $x^* \in \mathbf{W}_I^{*\circ}(s)$. We adorn the superscripts here with the relevant Frobenius endomorphisms for the given group. For the definition of the Deligne–Lusztig character in the disconnected group $\mathbf{C}_{\mathbf{G}^*}(s)$ we refer to [DM90, Prop. 1.3]. There is then a unique bijection $J_{s, \mathfrak{a}}^{\mathbf{G}, F} : \mathcal{E}_0(\mathbf{G}^F, s, \mathfrak{a}) \rightarrow \mathcal{E}_0(\mathbf{C}_{\mathbf{G}^*}(s)^{w_{\mathfrak{a}}^* F^*}, 1)$ satisfying the property

$$J_{s, \mathfrak{a}_1}^{\mathbf{L}_I, Fw_1} = (-1)^{l(w_1^*)} (J_{s, \mathfrak{a}}^{\mathbf{G}, F} \circ \mathbf{R}_{I, w}^{\mathbf{G}, F}).$$

We claim this a Jordan decomposition. Firstly, by transitivity of induction, we have

$$\mathbf{R}_{I, w_1}^{\mathbf{G}, F}(\mathbf{R}_{x^* w_{\mathfrak{a}_1}^*}^{\mathbf{L}_I, Fw_1}(s)) = \mathbf{R}_{x^* w_{\mathfrak{a}_1}^* w_1^*}^{\mathbf{G}, F}(s) = \mathbf{R}_{x^* w_{\mathfrak{a}}^*}^{\mathbf{G}, F}(s)$$

for any $x^* \in \mathbf{W}_I^{*\circ}(s) = \mathbf{W}^{*\circ}(s)$. This implies that

$$J_{s, \mathfrak{a}}^{\mathbf{G}, F}(\mathbf{R}_{x^* w_{\mathfrak{a}}^*}^{\mathbf{G}, F}(s)) = (-1)^{l(w_1^*)} J_{s, \mathfrak{a}_1}^{\mathbf{L}_I, Fw_1}(\mathbf{R}_{x^* w_{\mathfrak{a}_1}^*}^{\mathbf{L}_I, Fw_1}(s)) = (-1)^{l(w_{\mathfrak{a}}^*)} \mathbf{R}_{x^*}^{\mathbf{C}_{\mathbf{G}^*}(s), w_{\mathfrak{a}}^* F^*}(1).$$

That the signs agree is shown in [Lus84]. This shows the bijection is a Jordan decomposition.

After 16.7 we see that the image of γ in $\mathcal{O}(\mathbf{L}_I, Fw_1)$ is the restriction of γ to \mathbf{L}_I . Using the tensor product decomposition it suffices to prove the statement in the case $s = s_0$. But now the statement follows from Proposition 8.3 and (iii) of Theorem A.

(ii). Note that $\text{Aso}^*(\mathbf{G}, F)_{s, \mathfrak{a}} = \text{Inn}(\mathbf{T}, F)\Gamma(\mathbf{G}, F)_{s, \mathfrak{a}}$, where $\Gamma(\mathbf{G}, F)_{s, \mathfrak{a}} \leq \Gamma(\mathbf{G}, F)$ is the stabiliser of the series $\mathcal{E}_0(\mathbf{G}^F, s, \mathfrak{a})$. After 16.7, we see that the image of $\Gamma(\mathbf{G}, F)_{s, \mathfrak{a}}$ is contained in the subgroup $\text{Out}(\mathbf{L}_{J_1}, Fv_1) \times \Gamma(\mathbf{L}_{J_0}, Fv_0) \leq \text{Out}(\mathbf{L}_I, Fw_1)$. As $Z(\mathbf{L}_J)$ is connected, $\text{Out}(\mathbf{L}_J, Fv_1)$ contains no diagonal automorphisms. Hence, the image of $\text{Inn}(\mathbf{T}, F)$ is contained in the subgroup $\text{Inn}(\mathbf{S}_0, Fv_0) \leq \text{Out}(\mathbf{L}_{J_0}, Fv_0)$, where $\mathbf{S}_0 \leq \mathbf{L}_{J_0}$ is the maximal torus defined in 7.5. The statement now follows from the equivariance of the

bijection in Theorem 16.4 and Theorem A. ■

16.9. We note that the proof of Theorem D.(ii) given above also applies in the case $\mathbf{G} = \mathrm{Sp}(V)$, clarifying somewhat the proof in [Tay18, Thm. 16.2]. Now, we end with a discussion of the real-valued characters for these groups.

Corollary 16.10. *Keep the hypothesis of Theorem C.*

- (i) *If $\mathbf{G} = \mathrm{SO}(V)$, then $\chi \in \mathcal{E}(\mathbf{G}^F, s)$ is real-valued if and only if s is \mathbf{G}^{*F^*} -conjugate to s^{-1} .*
- (ii) *If $\mathbf{G} = \mathrm{Sp}(V)$, then $\chi \in \mathcal{E}(\mathbf{G}^F, s)$ is real-valued if and only if s is \mathbf{G}^{*F^*} -conjugate to s^{-1} and either -1 is not an eigenvalue of s or $q \equiv 1 \pmod{4}$.*

Proof. Let us continue the notation used in the proof of Theorem C above. We have $\chi \in \mathcal{E}_0(\mathbf{G}^F, s, \alpha)$ is real if and only if $\mathbb{Q}(\psi_0)$ and \mathbb{Q}_s are both real. But note that \mathbb{Q}_s is real if and only if the element of $\mathrm{Gal}(\mathbb{Q}(\zeta_d)/\mathbb{Q})$ mapping ζ_d to ζ_d^{-1} is a member of $\mathcal{G}_{s, \alpha}$, and hence (s, α) is in the same \mathbf{W}^* -orbit as (s^{-1}, α) . Applying Remark 16.8 completes the proof. ■

Acknowledgements

The first-named author was supported in part by grants from the Simons Foundation (Award No. 351233) and the National Science Foundation (Award No. DMS-1801156).

Part of this work was completed while the authors were in residence at the Mathematical Sciences Research Institute in Berkeley, California during the Spring 2018 semester program on Group Representation Theory and Applications, supported by the National Science Foundation under Grant No. DMS-1440140. The authors thank Radha Kessar for useful discussions concerning Section 8 and Martin Liebeck for sharing a proof of Corollary 9.8 whilst at the MSRI. The reduction to regular unipotent elements used in Section 9 was inspired by his proof.

Part of this work was also completed while the authors were in residence at the Isaac Newton Institute for Mathematical Sciences during the Spring 2020 programme Groups, Representations, and Applications: New Perspectives, supported by EPSRC grant number EP/R014604/1. The authors thank Gunter Malle for useful discussions and for his comments on an early preprint.

Both authors thank the MSRI, INI, and the organisers of the programs for making their stays possible and providing a collaborative and productive work environment.

Finally, the authors thank Ryan Vinroot for pointing out an inaccuracy in an earlier version of this work.

References

- [AMR96] A.-M. Aubert, J. Michel, and R. Rouquier, *Correspondance de Howe pour les groupes réductifs sur les corps finis*, Duke Math. J. **83** (1996) no. 2, 353–397.
- [Bon05] C. Bonnafé, *Quasi-isolated elements in reductive groups*, Comm. Algebra **33** (2005) no. 7, 2315–2337.

- [Bon06] ———, *Sur les caractères des groupes réductifs finis à centre non connexe: applications aux groupes spéciaux linéaires et unitaires*, Astérisque (2006) no. 306.
- [Bou02] N. Bourbaki, *Lie groups and Lie algebras. Chapters 4–6*, Elements of Mathematics (Berlin), Translated from the 1968 French original by Andrew Pressley, Berlin: Springer-Verlag, 2002.
- [BDT19] O. Brunat, O. Dudas, and J. Taylor, *Unitriangular shape of decomposition matrices of unipotent blocks*, preprint (2019), arXiv:910.08830 [math.RT].
- [CS17] M. Cabanes and B. Späth, *Inductive McKay condition for finite simple groups of type C*, Represent. Theory **21** (2017), 61–81.
- [CSSF20] M. Cabanes, B. Späth, and A. A. Schaeffer Fry, *On the inductive Alperin-McKay conditions in the maximally split case*, (2020), preprint.
- [Car93] R. W. Carter, *Finite groups of Lie type, Conjugacy classes and complex characters*, Wiley Classics Library, Chichester: John Wiley & Sons Ltd., 1993.
- [DM90] F. Digne and J. Michel, *On Lusztig's parametrization of characters of finite groups of Lie type*, Astérisque (1990) no. 181-182, 6, 113–156.
- [DM20] ———, *Representations of finite groups of Lie type*, 2nd edition, London Mathematical Society Student Texts, Cambridge: Cambridge University Press, 2020.
- [Gec93] M. Geck, *A note on Harish-Chandra induction*, Manuscripta Math. **80** (1993) no. 4, 393–401.
- [Gec03] ———, *Character values, Schur indices and character sheaves*, Represent. Theory **7** (2003), 19–55 (electronic).
- [GM00] M. Geck and G. Malle, *On the existence of a unipotent support for the irreducible characters of a finite group of Lie type*, Trans. Amer. Math. Soc. **352** (2000) no. 1, 429–456.
- [GM20] ———, *The character theory of finite groups of Lie type: A guided tour*, Cambridge Studies in Advanced Mathematics 187, Cambridge University Press, 2020.
- [GP00] M. Geck and G. Pfeiffer, *Characters of finite Coxeter groups and Iwahori-Hecke algebras*, vol. 21, London Mathematical Society Monographs. New Series, New York: The Clarendon Press Oxford University Press, 2000.
- [GLS98] D. Gorenstein, R. Lyons, and R. Solomon, *The classification of the finite simple groups. Number 3. Part I. Chapter A*, vol. 40, Mathematical Surveys and Monographs, Almost simple K-groups, American Mathematical Society, Providence, RI, 1998.
- [Isa18] I. M. Isaacs, *Characters of solvable groups*, vol. 189, Graduate Studies in Mathematics, American Mathematical Society, Providence, RI, 2018.
- [Li19] C. Li, *An equivariant bijection between irreducible Brauer characters and weights for $\mathrm{Sp}(2n, q)$* , J. Algebra **539** (2019), 84–117.
- [LS12] M. W. Liebeck and G. M. Seitz, *Unipotent and nilpotent classes in simple algebraic groups and Lie algebras*, vol. 180, Mathematical Surveys and Monographs, American Mathematical Society, Providence, RI, 2012.
- [Lus78] G. Lusztig, *Representations of finite Chevalley groups*, vol. 39, CBMS Regional Conference Series in Mathematics, Expository lectures from the CBMS Regional Conference held at Madison, Wis., August 8–12, 1977, Providence, R.I.: American Mathematical Society, 1978.
- [Lus84] ———, *Characters of reductive groups over a finite field*, vol. 107, Annals of Mathematics Studies, Princeton, NJ: Princeton University Press, 1984.
- [Lus09a] ———, *Remarks on Springer's representations*, Represent. Theory **13** (2009), 391–400.
- [Lus09b] ———, *Unipotent classes and special Weyl group representations*, J. Algebra **321** (2009) no. 11, 3418–3449.
- [LS85] G. Lusztig and N. Spaltenstein, *On the generalized Springer correspondence for classical groups*, in: *Algebraic groups and related topics (Kyoto/Nagoya, 1983)*, vol. 6, Adv. Stud. Pure Math. Amsterdam: North-Holland, 1985, 289–316.

- [Mal08] G. Malle, *Extensions of unipotent characters and the inductive McKay condition*, J. Algebra **320** (2008) no. 7, 2963–2980.
- [Mal17] ———, *Cuspidal characters and automorphisms*, Adv. Math. **320** (2017), 887–903.
- [MS16] G. Malle and B. Späth, *Characters of odd degree*, Ann. of Math. (2) **184** (2016) no. 3, 869–908.
- [NRSFV20] G. Navarro, N. Rizo, A. A. Schaeffer Fry, and C. Vallejo, *Characters and generation of Sylow 2-subgroups*, (2020), preprint.
- [NSV20] G. Navarro, B. Späth, and C. Vallejo, *A reduction theorem for the Galois-McKay conjecture*, Trans. Amer. Math. Soc. (2020), available online.
- [NT13] G. Navarro and P. H. Tiep, *Characters of relative p' -degree over normal subgroups*, Ann. of Math. (2) **178** (2013) no. 3, 1135–1171.
- [NT19] ———, *Sylow subgroups, exponents, and character values*, Trans. Amer. Math. Soc. **372** (2019) no. 6, 4263–4291.
- [NTT08] G. Navarro, P. H. Tiep, and A. Turull, *Brauer characters with cyclotomic field of values*, J. Pure Appl. Algebra **212** (2008) no. 3, 628–635.
- [RSFV20] N. Rizo, A. A. Schaeffer Fry, and C. Vallejo, *Galois action on the principal block and cyclic Sylow subgroups*, Algebra & Number Theory (2020), to appear.
- [SF19] A. A. Schaeffer Fry, *Galois automorphisms on Harish-Chandra series and Navarro’s self-normalizing Sylow 2-subgroup conjecture*, Trans. Amer. Math. Soc. **372** (2019) no. 1, 457–483.
- [SFT18] A. A. Schaeffer Fry and J. Taylor, *On self-normalising Sylow 2-subgroups in type A*, J. Lie Theory **28** (2018) no. 1, 139–168.
- [SFV19] A. A. Schaeffer Fry and C. R. Vinroot, *Fields of character values for finite special unitary groups*, Pacific J. Math. **300** (2019) no. 2, 473–489.
- [Spa85] N. Spaltenstein, *On the generalized Springer correspondence for exceptional groups*, in: *Algebraic groups and related topics (Kyoto/Nagoya, 1983)*, vol. 6, Adv. Stud. Pure Math. Amsterdam: North-Holland, 1985, 317–338.
- [Spä12] B. Späth, *Inductive McKay condition in defining characteristic*, Bull. Lond. Math. Soc. **44** (2012) no. 3, 426–438.
- [SV19] B. Srinivasan and C. R. Vinroot, *Galois group action and Jordan decomposition of characters of finite reductive groups with connected center*, J. Algebra (2019), available online.
- [Tay13] J. Taylor, *On unipotent supports of reductive groups with a disconnected centre*, J. Algebra **391** (2013), 41–61.
- [Tay18] ———, *Action of automorphisms on irreducible characters of symplectic groups*, J. Algebra **505** (2018), 211–246.
- [Tes95] D. Testerman, *A_1 -type overgroups of elements of order p in semisimple algebraic groups and the associated finite groups*, J. Algebra **177** (1995) no. 1, 34–76.
- [TZ04] P. H. Tiep and A. E. Zalesskiĭ, *Unipotent elements of finite groups of Lie type and realization fields of their complex representations*, J. Algebra **271** (2004) no. 1, 327–390.
- [Tur01] A. Turull, *The Schur indices of the irreducible characters of the special linear groups*, J. Algebra **235** (2001) no. 1, 275–314.
- [Vin20] C. R. Vinroot, *Totally orthogonal finite simple groups*, Math. Z. **294** (2020) no. 3–4, 1759–1785.
- [Wal04] J.-L. Waldspurger, *Une conjecture de Lusztig pour les groupes classiques*, Mém. Soc. Math. Fr. (N.S.) (2004) no. 96.
- [Zas62] H. Zassenhaus, *On the spinor norm*, Arch. Math. **13** (1962), 434–451.

J. TAYLOR, DEPARTMENT OF MATHEMATICS, UNIVERSITY OF SOUTHERN CALIFORNIA,
LOS ANGELES, CA 90089, USA

Email: jayt@usc.edu

A. A. SCHAEFFER FRY, DEPARTMENT OF MATHEMATICAL AND COMPUTER SCIENCES,
METROPOLITAN STATE UNIVERSITY OF DENVER, DENVER, CO 80217, USA
Email: aschaef6@msudenver.edu