

## Some Applications of Integration <sup>1</sup>

Integration can be used to obtain many properties connected with curves.

### Area Between Two Curves

Integration and area are closely related


*Example.*

The curves  $y = 2/x$  and  $y = 2 - \frac{1}{6}(x - 1)^2$  intersect at  $x = 1$  and  $x = 4$

The area between them is simply

## Area in Polar Coordinates <sup>2</sup>

Consider the curve:  $r = \sin(2\theta)$  for  $0 \leq \theta \leq \frac{1}{2}\pi$



If  $\theta$  increases by  $d\theta$   
the radial vector  $r$   
sweeps through an area  
of  $dA = \frac{1}{2}r^2 d\theta$ .

The whole area of the curve is

## Length of a Curve <sup>3</sup>

Suppose that  $y = f(x)$  is plotted as a curve

and  $s(x)$  is the 'arclength' of the curve from some point, then

if  $x$  changes by  $dx$  and

$y$  changes by  $dy$

$$s \text{ changes by } ds = \sqrt{dx^2 + dy^2} \\ = \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

so the length of the curve from  $x = a$  to  $x = b$  is

$$\int_{x=a}^{x=b} ds = \int_a^b \sqrt{1 + \left(\frac{dy}{dx}\right)^2} dx$$

## Circumference of a Circle <sup>4</sup>

A circle of radius  $a$  can be parameterised as

$$x = a \cos t \quad y = a \sin t$$

with  $dx = -a \sin t dt$   $dy = a \cos t dt$


$$ds = \sqrt{dx^2 + dy^2} = \sqrt{a^2 \sin^2 t + a^2 \cos^2 t} dt \\ = a dt$$

The arclength of the circumference is

## Surfaces and Volumes of Revolution <sup>5</sup>

Rotating a curve  $y = f(x)$  about the  $x$ -axis creates

- a surface of revolution of area  $A$ , and
- a volume of revolution,  $V$



As  $x$  increases by  $dx$ , the area and volume increase by

$$dA = 2\pi y ds$$

$$dV = \pi y^2 dx$$

so that the area and volume (from  $x = a$  to  $x = b$ ) are

$$A = \int_{x=a}^{x=b} 2\pi y ds \quad \text{and} \quad V = \int_a^b \pi y^2 dx$$

## Surface and Volume of a Bucket <sup>6</sup>

*Example.* The line  $y = \frac{1}{2}x$  rotated about the  $x$ -axis creates a bucket-shape for  $2 \leq x \leq 4$ .

## Rotation about the $y$ -axis <sup>7</sup>

Rotating the curve

$$y = f(x)$$

about the  $y$ -axis, the surface area and volume increase by

$$dA = 2\pi x ds$$

$$dV = \pi x^2 dy$$


as  $x$  increases by  $dx$ .

Hence the area and volume,

from  $y = f(a)$  to  $y = f(b)$

are

$$A = \int_{x=a}^{x=b} 2\pi x ds \quad \text{and} \quad V = \int_a^b \pi x^2 \frac{dy}{dx} dx$$



## Surface and Volume of a Paraboloid <sup>8</sup>

*Example.* The parabola  $y = \frac{1}{2}x^2$  for  $0 \leq y \leq 2$