

Real and Complex Numbers

Number Sets. there are 5 basic number sets¹:

$\mathbb{N} = \{1, 2, 3, \dots\}$ the '*natural numbers*'
(positive whole numbers as used for counting)

$\mathbb{Z} = \{\dots, -3, -2, -1, 0, 1, 2, 3, \dots\}$ '*integers*'
(include zero and negative whole numbers)

$\mathbb{Q} = \{\frac{m}{n} \mid m \in \mathbb{Z}, n \in \mathbb{N}\}$ '*rational numbers*'
(include all fractions)

\mathbb{R} the set of '*real numbers*'
(equivalent to all finite or infinite decimal Nos.)

\mathbb{C} the set of '*complex numbers*'
(include all 'imaginary' numbers)

Note that $\mathbb{N} \subset \mathbb{Z} \subset \mathbb{Q} \subset \mathbb{R} \subset \mathbb{C}$

We will examine the last two in more detail now.

¹a summary of set notation is available on the web page

Real Numbers, \mathbb{R}

a real number measures distance along a '*real line*'

(you will often hear a real number called '*a point*')

Real numbers satisfy all algebraic and inequality laws.
these can be summarised as sets of 'axioms':

field axioms for any x, y, z in \mathbb{R}

$$\begin{array}{ll} x + 0 = x & x + y = y + x \\ x - x = 0 & (x + y) + z = x + (y + z) \\ 1x = x & xy = yx \\ x/x = 1 \text{ (for } x \neq 0) & (xy)z = x(yz) \\ & x(y + z) = xy + xz \end{array}$$

ordering axioms for any x, y, z in \mathbb{R}

only one of ' $x > y$ ', ' $x = y$ ' or ' $x < y$ ' can hold
if $x < y$ and $y < z$ then $x < z$
if $x < y$ then $x + z < y + z$
if $x < y$ and $z > 0$ then $xz < yz$

(You will learn more about 'fields' in other courses)

a few points to note

All you know about the algebra and inequalities can be worked out from the axioms.

Example 1. Show that $0x = 0$ for any real number x .

Example 2. If $x > y > 0$ then $\frac{1}{y} > \frac{1}{x} > 0$.

Example 3. If $x > y$ and $z < 0$ then $zx < zy$.

Example 4. Is infinity (∞) a real number?

Note. We use x, y, z , etc. only as names or labels.


All of these definitions, formulae, etc., remain exactly the same if we use other names, such as s, t, u , etc.

Intervals

$a < x \leq b$ represents any value of x in an 'interval'

There is a special notation for intervals:

- if $a < x \leq b$ we write $x \in (a, b]$
- if an end-point is not in the interval it is called an 'open' end point, marked by a curved bracket
- if an end-point is in the interval it is called a 'closed' end point, marked by a square bracket



- if an interval extends to infinity use $(-\infty$ or $\infty)$
e.g. $x \geq a$ means $x \in [a, \infty)$

Note: infinity can only be an open end point.

Absolute Value or 'Modulus'

The absolute value $|x|$ of a real number x gives its 'size' but not its sign.

$$\text{We can define: } |x| = \begin{cases} x & \text{if } x \geq 0 \\ -x & \text{if } x < 0 \end{cases}$$

sketch:

Example 1. In what interval of x is $|x + 3| < 1$?

Example 2. In what intervals of x is $|x - 1| \geq 7$?

Squares and Square Roots

For any real number x :

$$\begin{aligned} \text{either } x > 0 & \text{ so } x^2 > 0 \\ \text{or } x = 0 & \text{ so } x^2 = 0 \\ \text{or } x < 0 & \text{ so } x^2 > 0 \end{aligned}$$

so we always have $x^2 \geq 0$

Hence we can also define

$$|x| = \sqrt{x^2}$$

Note: for any real number a :

- \sqrt{a} only exists if $a \geq 0$
- \sqrt{a} is the positive square root of a

Example 1. Solve $x^2 = 1$ for x .

Example 2. Can you solve $x^2 = -1$ for x ?

Imaginary Numbers

There is no real number x for which $x^2 < 0$

but it is (remarkably) useful to invent new numbers (imaginary numbers) for which $x^2 < 0$

Let i be defined as

$$i = \sqrt{-1}$$

Example 1. Now can you solve $x^2 = -1$ for x ?

- Note:**
- clearly $i \notin \mathbb{R}$ (i is not 'real')
 - i is an 'imaginary' square root of -1
 - if $x \in \mathbb{R}$ then xi is an imaginary number such that

$$(xi)^2 = x^2 i^2 = -x^2$$

Example 2. What are the values of i^2, i^3, i^4, i^5 , etc.?

Complex Numbers, \mathbb{C}

Complex numbers arise if we add real and imaginary numbers.

If x and y are real, then

$$z = x + iy \text{ is a complex number}$$

we can write: $x + iy \in \mathbb{C}$

Standard Form of a complex number is

'real part' + i 'imaginary part'

If x and y are real, then we define

$$\text{'real part': } \operatorname{Re}(x + iy) = x$$

$$\text{'imaginary part': } \operatorname{Im}(x + iy) = y$$

Note: the real part is a real number
the imaginary part is a real number

Example. Give the real and imaginary parts of $7 - 9i$

Complex Plane and Polar Form

We can extend the 'real line' to a 'complex plane'

A complex number $z = x + iy$ is a point with

horizontal coordinate x
vertical coordinate y

It is also useful to use polar coordinates:

$$\begin{aligned}z &= r(\cos(\theta) + i\sin(\theta)) \\ &= r\cos(\theta) + ir\sin(\theta)\end{aligned}$$

having: $x = r\cos(\theta)$ and $y = r\sin(\theta)$

Names: r is the 'modulus'. θ is the 'argument'.

Example. $-1 + i\sqrt{3} =$

Modulus and Complex Conjugate

The modulus is the distance from the origin in the complex plane

so for $z = x + iy$ we define

$$|z| = \sqrt{x^2 + y^2}$$

and for $z = r(\cos(\theta) + i\sin(\theta))$ we have, simply

$$|z| = r$$

The complex conjugate or simply **the conjugate** of a complex number has the sign of the imaginary part changed

so if $z = x + iy$ we define the complex conjugate of z as

$$\bar{z} = x - iy$$

Note

$$\begin{aligned}z\bar{z} &= (x + iy)(x - iy) \\ &= x^2 - i^2y^2 \\ &= x^2 + y^2\end{aligned}$$

So:

$$\begin{aligned}z\bar{z} &\text{ is real} \\ z\bar{z} &= |z|^2 \\ \text{or } |z| &= \sqrt{z\bar{z}}\end{aligned}$$

Algebra of Complex Numbers

Complex numbers can be added, subtracted, multiplied, divided just like real numbers (simply use $i^2 = -1$ where it appears)

For $z = x + iy$ and $w = u + iv$ give the following in 'standard form'

Example 1. $z + w$

Example 2. $z - w$

Example 3. zw

Example 4. z/w (provided $w \neq 0$)

Two Examples

Convert to standard form:

$$\frac{3 + 4i}{1 - 2i} =$$

Where is $|z - (\pi + i\sqrt{2})| \leq 1$ in the complex plane?