

MATH10222: EXAMPLE SHEET¹ IV*Questions for supervision classes*

Hand in the solutions to questions 1a,c,d and 3. Attempt all other questions too and raise any problems with your supervisor. Question 1b should be straightforward. If you're a self-respecting mathematician you should find question 2 interesting.

1. Inhomogeneous linear second-order ODEs with constant coefficients**(a) Exploiting linearity**

Find the general solutions of each of the following linear ODEs:

i.

$$\ddot{y} + 3\dot{y} + 2y = 4e^{2t}.$$

ii.

$$\ddot{y} + 3\dot{y} + 2y = 7.$$

iii. Use the results just obtained to write down the general solution of the ODE

$$\ddot{y} + 3\dot{y} + 2y = 4e^{2t} + 7.$$

(b) Using complex variables for trigonometric RHSs

Find the general solutions of

i.

$$\ddot{y} + 2\dot{y} + 2y = 10 \cos t$$

ii.

$$\ddot{y} + 2\dot{y} + 2y = 10 \sin t.$$

[**Hint:** Use complex variables to do both in one go.]

(c) Degenerate and non-degenerate cases for RHSs of exponential form

Find the general solutions of the following ODEs. Watch out for degenerate cases!

i.

$$\ddot{y} + 3\dot{y} + 2y = 2e^{-t}$$

ii.

$$\ddot{y} + 4\dot{y} + 4y = e^{-2t}$$

iii.

$$\ddot{y} + 2\dot{y} + 2y = 5 \cosh t$$

iv.

$$\ddot{y} + 3\dot{y} + 2y = 2 \cosh t$$

¹Any feedback to: M.Heil@maths.man.ac.uk

(d) **Degenerate and non-degenerate cases for polynomial RHSs**

Find the general solutions of the following ODEs. Watch out for degenerate cases!

i.

$$\ddot{y} + 3\dot{y} + 2y = 1 + t^2$$

ii.

$$\ddot{y} + 2\dot{y} = 1 + t^2$$

iii.

$$\ddot{y} = 1 + t^2$$

2. **Linear ODEs with non-constant coefficients: Euler's ODE**

(a) Homogeneous second-order Euler ODEs have the form

$$at^2\ddot{y} + bt\dot{y} + cy = 0,$$

where a , b and c are constants. Explain why looking for solutions of the form $y \sim t^n$ is a promising ansatz.

(b) Use the ansatz to obtain the general solution of

$$t^2\ddot{y} + 2t\dot{y} - 2y = 0$$

and explain why your method works.

(c) Explain why the ansatz doesn't work for the ODE

$$t^2\ddot{y} - t\dot{y} + y = 0.$$

How would you obtain a second, linearly independent solution for this ODE?

3. **Non-linear ODEs with special properties**

(a) Solve the autonomous nonlinear ODE

$$yy'' = (y')^2.$$

(b) Solve the initial value problem

$$y'' = \frac{2x^2}{(y')^2}$$

subject to

$$y'(1) = 2^{1/3} \quad \text{and} \quad y(1) = 2^{-2/3}.$$

[**Hint:** Apply the initial conditions as soon as possible, otherwise the algebra gets messy...]