

# MT3261: EXAMPLE SHEET<sup>1</sup> II

- 1.) A 2D flow field is given by  $u_1 = ax_2$  and  $u_2 = ax_1$ .
- a) Sketch the flow field (begin by considering the velocity vectors on the coordinate axes and on the main diagonals).
  - b) Determine the trajectories  $x_i^p(t)$  of particles which are at position  $x_i = X_i$  at time  $t = 0$  by integrating the equations of motion  $\partial x_i^p(t)/\partial t = u_i(x_j^p(t), t)$ .
  - c) Determine the acceleration of material particles directly from their trajectories (i.e. by evaluating  $a_i = \partial^2 x_i^p(t)/\partial t^2$ ).
  - d) Compare this to the result obtained by using the material derivative  $a_i = Du_i/Dt$ .
  - e) What is the divergence of the flow field?
- 2.) A 2D flow field is given by  $u_1 = Ux_2$  and  $u_2 = 0$ .
- a) Sketch the flow field in the region  $x_2 \in [0, 1]$ .
  - b) Determine the rate of strain and the rate of rotation tensors  $\epsilon_{ij}$  and  $\omega_{ij}$ .
  - c) Sketch the deformation of a small rectangular fluid element whose edges of lengths  $\delta x_1$  and  $\delta x_2$  are parallel to the  $x_1$  and  $x_2$  axes, respectively. (Hint: Sketch the corners of the fluid element at time  $t$  and at time  $t + \delta t$ ). Relate this to the interpretation of the rate of strain and rate of rotation tensors.

*Coursework*

Please hand in the solution to question 2 by 12 noon on Friday. Please place them into the envelope at the door of Dr. Heil's office (room 1.05 in the Lamb building).

---

<sup>1</sup>Any feedback to: [M.Heil@maths.man.ac.uk](mailto:M.Heil@maths.man.ac.uk)