


MATH35001: EXAMPLE SHEET¹ V

- 1.) The figure below shows a film of Newtonian incompressible fluid on an inclined belt which is moving with constant velocity U . The no-slip condition applies on the surface of the belt, i.e. the fluid particles on the belt move with velocity $\mathbf{u}(y=0) = -U\mathbf{e}_x$. Gravity acts vertically downwards and a strong wind exerts a tangential shear stress τ_0 in the negative x -direction onto the surface of the fluid. You can assume that the flow is steady and unidirectional and that the film thickness h is constant along the belt (and given). The air pressure at the free surface is constant and given by $p = 0$.
- Determine the velocity field and the pressure distribution in the fluid.
 - Determine the volume flux Q (per unit width of the belt) in the positive x -direction, i.e. evaluate $Q = \int_0^h u(y) dy$.
 - Now consider the case $U = 0$, i.e. a stationary belt. Determine the critical value τ_0^{crit} of the shear stress τ_0 for which the volume flux becomes negative (in other words, for $\tau_0 > \tau_0^{crit}$ overall the fluid flows 'up the hill'). Sketch the velocity distributions for the cases $\tau_0 = 0$ and $\tau_0 = \tau_0^{crit}$ (still assuming that $U = 0$).



- 2.) Fluid is confined between two infinite parallel plates at $y = 0$ and $y = h$. An externally applied pressure gradient $\nabla p = G\mathbf{e}_x$ drives the fluid in the x -direction. The plates are porous and fluid is driven through the top surface at $y = h$ with a uniform normal velocity V and leaves the bottom wall at the same uniform velocity such that $\mathbf{u} = -V\mathbf{e}_y$ at $y = 0$ and $y = h$. You can assume that there is no motion in the z -direction (i.e. $w = 0$) and that all quantities are independent of z .
- Explain why $\mathbf{u}(x, y, t) = (u(y), -V)$ is a plausible guess for the velocity field.
 - Show that the velocity field assumed in (i) is consistent with the 2D Navier Stokes equations and the equation of continuity and that the only non-trivial equation is given by

$$\nu \frac{\partial^2 u}{\partial y^2} + V \frac{\partial u}{\partial y} = \frac{G}{\rho}.$$

- Solve this equation subject to the no-slip condition for u on the top and bottom walls [Hint: The constant term on the RHS is a singular form since a constant function is already contained in the solution of the homogeneous equation. Therefore, the particular solution must have the form 'constant $\times y$ '].

Coursework

Please hand in the solutions to question 1 by 12 noon on Thursday. Please place them into the envelope at the door of Prof. Heil's office (room 2.224 in the Alan Turing building).

¹Any feedback to: M.Heil@maths.man.ac.uk