

Mathematics 0C2/MATH19832  
Workbook  
Spring 2018

Dr Tuomas Sahlsten

Office 2.120, Alan Turing Building

Email: [tuomas.sahlsten@manchester.ac.uk](mailto:tuomas.sahlsten@manchester.ac.uk)

This workbook is heavily based on Dr Marianne Johnson's workbook from Spring 2017

Including examples and exercises to be completed during class

Website:

<http://personalpages.manchester.ac.uk/staff/tuomas.sahlsten/courses/0c2>

School of Mathematics, University of Manchester

# Contents

<b>1</b>	<b>Integration</b>	<b>1</b>
1.1	Derivatives and antiderivatives . . . . .	2
1.2	Indefinite integrals and inspection . . . . .	6
1.3	Definite integrals and the area under a graph . . . . .	9
1.4	Integration by substitution . . . . .	18
<b>2</b>	<b>Trigonometry</b>	<b>22</b>
2.1	Trigonometric functions . . . . .	23
2.2	Trigonometric identities and integration . . . . .	26
2.3	Inverse trigonometric functions . . . . .	32
<b>3</b>	<b>Sequences, series and summations</b>	<b>40</b>
3.1	Arithmetic progressions . . . . .	41
3.2	Geometric progressions . . . . .	44
3.3	The binomial theorem . . . . .	50
<b>4</b>	<b>Further calculus</b>	<b>56</b>
4.1	Differentiation revisited . . . . .	57
4.2	Implicit differentiation . . . . .	63
4.3	Integration revisited . . . . .	68
4.4	Integration by parts . . . . .	69
4.5	Taylor series . . . . .	72
<b>5</b>	<b>Rational functions</b>	<b>77</b>
5.1	Polynomials and long division . . . . .	78
5.2	Introducing partial fractions . . . . .	82
5.3	The method of partial fractions . . . . .	85
5.4	Integration by partial fractions . . . . .	90

# Chapter 1

## Integration

- 1.1 Derivatives and antiderivatives
- 1.2 Indefinite integrals and inspection
- 1.3 Definite integrals and the area under a graph
- 1.4 Integration by substitution

### Revision from last semester..

The **derivative** of a function  $f(x)$  is given by

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$$

Sometimes we write

$$f'(x) = \frac{df}{dx}$$

Function	Derivative	Notes
$m$	0	$m$ any constant
$x^n$	$nx^{n-1}$	$n$ any (non-zero) real number
$e^x$	$e^x$	
$\ln(x)$	$\frac{1}{x}$	
$\sin(x)$	$\cos(x)$	for $x$ in radians
$\cos(x)$	$-\sin(x)$	for $x$ in radians
$\tan(x)$	$\sec^2(x)$	for $x$ in radians
$\lambda f(x)$	$\lambda f'(x)$	(constant $\times$ function)
$f(x) + g(x)$	$f'(x) + g'(x)$	(sum of functions)
$f(u(x))$	$f'(u(x))u'(x)$	(function of a function; 'chain rule')
$u(x)v(x)$	$u'(x)v(x) + u(x)v'(x)$	(product of functions; 'product rule')
$\frac{u(x)}{v(x)}$	$\frac{u'(x)v(x) - u(x)v'(x)}{v(x)^2}$	(quotient of functions; 'quotient rule')



**Exercise**

If  $f'(x) = 3x^2 + 2$ , what could  $f(x)$  be?

→→→→→→→→→→ **Reversing differentiation** →→→→→→→→→→

$f'(x) = 3x^2 + 2$	$f(x) =$
--------------------	----------

There is more than one correct answer! (See below for an explanation.)

**Definition (Antiderivative)**

An **antiderivative** of a function  $f(x)$  is a function  $F(x)$  whose derivative is equal to  $f(x)$ .

**Example**

The function  $F(x) = x^2$  is an antiderivative of  $f(x) = 2x$ , because

$$F'(x) = 2x = f(x).$$

The function  $G(x) = x^2 - 12$  is another antiderivative of  $f(x) = 2x$ , because

$$G'(x) = 2x = f(x).$$

In fact, it is easy to see that if  $C$  is any constant then the function  $H(x) = x^2 + C$  will also be an antiderivative of  $f(x) = 2x$ .

**Note:** The example above illustrates that once one antiderivative is known, infinitely many can be found by adding on different constant terms. In fact, two antiderivatives of the same function can *only* differ by addition of a constant term, as we shall now show.

**Theorem 1**

If  $F(x)$  and  $G(x)$  are two antiderivatives of  $f(x)$ , then  $F(x) = G(x) + C$  for some constant  $C$ .

**Reason:** If  $F(x)$  and  $G(x)$  are two antiderivatives of  $f(x)$ , then  $F'(x) = f(x)$  and  $G'(x) = f(x)$ , which gives

$$F'(x) - G'(x) = f(x) - f(x) = 0, \text{ and hence } \frac{d}{dx}(F(x) - G(x)) = 0.$$

Since only constants differentiate to give zero, it follows that  $F(x) - G(x) = C$  for some constant  $C$ . Rearranging this gives  $F(x) = G(x) + C$ .

**Note:** It follows from the above that if we can find just one antiderivative, then we can describe all the others by adding on constants.

**Exercise**

For each of the following functions  $f(x)$ , find *all* antiderivatives:

→→→→→→→→→→→→ <b>Reversing differentiation</b> →→→→→→→→→→→→	
$f(x) = \frac{1}{x}$ is the derivative of $\boxed{\ln(x)}$	Every antiderivative of $f(x) = \frac{1}{x}$ has the form $F(x) = \boxed{\ln(x) + C}$ , where $C$ is a constant.
$f(x) = e^x$ is the derivative of	Every antiderivative of $f(x) = e^x$ has the form $F(x) =$
$f(x) = \cos(x)$ is the derivative of	Every antiderivative of $f(x) = \cos(x)$ has the form $F(x) =$
$f(x) = -\sin(x)$ is the derivative of	Every antiderivative of $f(x) = -\sin(x)$ has the form $F(x) =$
$f(x) = \sec^2(x)$ is the derivative of	Every antiderivative of $f(x) = \sec^2(x)$ has the form $F(x) =$
$f(x) = nx^{n-1}$ is the derivative of	Every antiderivative of $f(x) = nx^{n-1}$ has the form $F(x) =$
$f(x) = m$ , a constant is the derivative of	Every antiderivative of $f(x) = m$ has the form $F(x) =$

**Note:** Sometimes we may wish to find a *particular* antiderivative, satisfying an extra condition.

**Example**

Find the antiderivative  $F(x)$  of  $f(x) = 2x$ , satisfying  $F(1) = 2$ .

**Solution:**

- $f(x) = 2x$  is the derivative of  $x^2$ .
- Every antiderivative of  $f(x)$  has the form  $F(x) = x^2 + C$  where  $C$  is a constant.
- To find the correct value of  $C$ , substitute  $x = 1$  and rearrange.  
We obtain  $F(1) = 1^2 + C$ . Since  $F(1) = 2$  this gives  $2 = 1 + C$  and hence  $C = 1$ .
- The required antiderivative is therefore  $F(x) = x^2 + 1$

**Exercise**

Find the antiderivative  $F(x)$  of  $f(x) = \cos(2x)$  satisfying  $F\left(\frac{\pi}{4}\right) = 0$ .

**Solution:**

- $f(x) = \cos(2x)$  is the derivative of
- Every antiderivative of  $f(x)$  has the form  $F(x) =$  , where  $C$  is a constant.
- To find the correct value of  $C$ , substitute  $x = \frac{\pi}{4}$  and rearrange.  
We obtain  $F\left(\frac{\pi}{4}\right) =$   
Since  $F\left(\frac{\pi}{4}\right) = 0$  this gives and hence  $C =$
- The required antiderivative is therefore  $F(x) =$

## 1.2 Indefinite integrals and inspection

### Definition (Indefinite integral)

The **indefinite integral**  $\int f(x) dx$  represents all possible antiderivatives of  $f(x)$ . In other words, if  $F(x)$  is some particular antiderivative for  $f(x)$ , then

$$\int f(x) dx = F(x) + C,$$

where  $C$  is an *undetermined* constant term.

**Note:** The mathematical expression above is read aloud as

*The (indefinite) integral of the function “f of x” with respect to x is equal to “capital F of x” plus a constant term, C.*

### Example

Find  $\int 3x^2 dx$ .

**Solution:** We have seen that any function of the form  $x^3 + C$  (where  $C$  is a constant) is an antiderivative of  $3x^2$ . Hence,

$$\int 3x^2 dx = x^3 + C.$$

You can apply **your knowledge of derivatives from last semester** to find indefinite integrals. By looking at (or ‘inspecting’) the function  $f(x)$  you can try to recognise it as the derivative of some other function. This process is called **integration by inspection**. We have already seen some basic examples of this:

Function	Indefinite integral
$m$ (a constant)	$\int m dx = mx + C$
$nx^{n-1}$	$\int nx^{n-1} dx = x^n + C$
$e^x$	$\int e^x dx = e^x + C$
$\frac{1}{x}$	$\int \frac{1}{x} dx = \ln x  + C$
$\sin(x)$	$\int \sin(x) dx = -\cos(x) + C$
$\cos(x)$	$\int \cos(x) dx = \sin(x) + C$
$\sec^2(x)$	$\int \sec^2(x) dx = \tan(x) + C$



When asked to integrate a function, the general strategy is to try to *manipulate* the function into a well-known form (such as the forms in the table above) which can be easily integrated. These forms are called “standard forms” or “table integrals”.

Keep in mind that integration by inspection may involve a bit of guess work on your part at first - as the course progresses you will learn some more techniques from basic algebra and trigonometry to help, as well as more advanced methods of integration. In general a calculation may contain several steps. Thus when finding indefinite integrals, we shall agree to add a constant only at the end of the calculation.

**Example**

Let  $n$  be an integer with  $n \neq -1$ . Find  $\int x^n dx$ .

**Solution:** We know that

$$\frac{d}{dx} (x^{n+1}) = (n+1)x^n.$$

Since  $n \neq -1$  we can divide both sides by  $n+1$  to find that

$$\frac{d}{dx} \left( \frac{1}{n+1} x^{n+1} \right) = x^n.$$

Thus

$$\int x^n dx = \frac{1}{n+1} x^{n+1} + C.$$

**Example**

Let  $a, b$  be constants with  $a \neq 0$ . Find  $\int e^{ax+b} dx$ .

**Solution:** We know that

$$\frac{d}{dx} (e^{ax+b}) = ae^{ax+b}.$$

Since  $a \neq 0$  we can divide both sides by  $a$  to find that

$$\frac{d}{dx} \left( \frac{1}{a} e^{ax+b} \right) = e^{ax+b}.$$

Thus

$$\int e^{ax+b} dx = \frac{1}{a} e^{ax+b} + C.$$

**Example**

Find  $\int 3x^2 + 4x \, dx$ .

**Solution:** We note that

$$3x^2 + 4x = \frac{d}{dx}(x^3) + 2 \times \frac{d}{dx}(x^2) = \frac{d}{dx}(x^3 + 2x^2).$$

Thus

$$\int 3x^2 + 4x \, dx = x^3 + 2x^2 + C.$$

The previous examples illustrate some general principles:

**Theorem 2**

Let  $f(x)$ ,  $g(x)$  be functions of  $x$ , and let  $\lambda$  be a constant. Then

$$\int [f(x) + g(x)] \, dx = \int f(x) \, dx + \int g(x) \, dx + C$$

$$\int \lambda f(x) \, dx = \lambda \int f(x) \, dx + C$$

**Example**

Find  $I = \int (x + \frac{1}{x})^2 \, dx$ .

**Solution:** By expanding the brackets we find

$$(x + \frac{1}{x})^2 = (x + \frac{1}{x})(x + \frac{1}{x}) = (x^2 + 1 + 1 + \frac{1}{x^2}) = x^2 + 2 + x^{-2}$$

Thus

$$\begin{aligned} I &= \int x^2 \, dx + \int 2 \, dx + \int x^{-2} \, dx \\ &= \frac{1}{3}x^3 + 2x - \frac{1}{x} + C. \end{aligned}$$

**Check:**

$$\frac{d}{dx} \left( \frac{1}{3}x^3 + 2x - x^{-1} + C \right) = x^2 + 2 + x^{-2}$$

**Example**

Find the indefinite integral

$$\int \frac{1}{1+2x} dx.$$

**Solution:** Recalling that  $\int \frac{1}{x} dx = \ln|x| + C$ , we decide to inspect the function  $\ln|1+2x|$ . By the chain rule we have

$$\frac{d}{dx} (\ln|1+2x|) = \frac{1}{1+2x} \times 2 = \frac{2}{1+2x}.$$

Therefore

$$\int \frac{1}{1+2x} dx = \frac{1}{2} \times \int \frac{2}{1+2x} dx = \frac{1}{2} \ln|1+2x| + C.$$

**Check:**

$$\frac{d}{dx} \left( \frac{1}{2} \ln|1+2x| + C \right) = \frac{1}{2} \times \frac{1}{1+2x} \times 2 = \frac{1}{1+2x}$$

**Remark:** All of the examples we have considered so far concern functions of a variable  $x$ . There is nothing special about the letter  $x$ . Depending on the nature of the problem, we may consider, for example, functions  $f(t)$  of an argument denoted by  $t$ . (The letter  $t$  is a traditional notation for the time variable.)

### 1.3 Definite integrals and the area under a graph

**Definition** (Definite integral)

For constants  $a$  and  $b$ , the **definite integral**  $\int_a^b f(x) dx$  is given by

$$\int_a^b f(x) dx = F(b) - F(a)$$

where  $F(x)$  is an antiderivative of  $f(x)$ .

(The result is **independent** of choice of antiderivative  $F(x)$ .)

**Example**

Find  $\int_2^3 3x^2 dx$ .

**Solution:** Since  $F(x) = x^3$  is an antiderivative of  $3x^2$  we find that

$$\int_2^3 3x^2 dx = F(3) - F(2) = (3^3) - (2^3) = 27 - 8 = 19.$$

Notice that if  $C$  is a constant then  $F(x) = x^3 + C$  is another antiderivative of  $3x^2$ , also giving

$$\int_2^3 3x^2 dx = F(3) - F(2) = (3^3 + C) - (2^3 + C) = 27 - 8 = 19.$$

**Notation:** It will often be convenient to write  $[F(x)]_a^b$  to denote  $F(b) - F(a)$  in our calculations.

**Example**

Find  $\int_0^1 \sqrt{x} dx$ .

**Solution:**

$$\int_0^1 \sqrt{x} dx = \int_0^1 x^{1/2} dx = \left[ \frac{x^{3/2}}{3/2} \right]_0^1 = \left[ \frac{2}{3} x^{3/2} \right]_0^1 = \left( \frac{2}{3} 1^{3/2} \right) - \left( \frac{2}{3} 0^{3/2} \right) = \frac{2}{3}.$$

The following properties can be easily deduced from the above definition:

**Theorem 3**

Let  $f(x)$ ,  $g(x)$  be functions of  $x$ , and let  $\lambda$  be a constant. Then

$$\int_a^b (f(x) + g(x)) dx = \int_a^b f(x) dx + \int_a^b g(x) dx.$$

$$\int_a^b \lambda f(x) dx = \lambda \int_a^b f(x) dx.$$

**Example**

Find  $\int_{-3}^{-1} \frac{1}{x} + x \, dx$ .

**Solution:**

$$\begin{aligned}\int_{-3}^{-1} \frac{1}{x} + x \, dx &= \int_{-3}^{-1} \frac{1}{x} \, dx + \int_{-3}^{-1} x \, dx \\ &= [\ln |x|]_{-3}^{-1} + \left[ \frac{1}{2} x^2 \right]_{-3}^{-1} \\ &= (\ln(1) - \ln(3)) + \left( \frac{1}{2}(-1)^2 - \frac{1}{2}(-3)^2 \right) \\ &= \ln(1) - \ln(3) + \frac{1}{2} - \frac{9}{2} = -\ln(3) - 4 \approx -5.10.\end{aligned}$$

**Example**


Find  $\int_0^{\pi/2} 5 \sin(x) \, dx$ .

**Solution:**

$$\begin{aligned}\int_0^{\pi/2} 5 \sin(x) \, dx &= 5 \int_0^{\pi/2} \sin(x) \, dx \\ &= 5 [-\cos(x)]_0^{\pi/2} \\ &= 5((-\cos(\pi/2)) - (-\cos(0))) \\ &= 5(0 + 1) = 5.\end{aligned}$$


So far we have concentrated on *how to calculate* the value of a given definite integral, without considering *the meaning* behind this value. This turns out to be closely related to the area under a curve, as we shall now explain.

Consider the area under a curve  $y = f(x)$  between a fixed point  $a$  and a variable point  $x$ :



Note that this area depends on both the lower limit (which we have fixed) and the variable upper limit,  $x$ . In other words it is a *function of  $x$* , which we shall denote by  $A(x)$ . We shall show that  $A(x)$  is an *antiderivative* of  $f(x)$ .

If we change the upper limit by a small amount  $h$ ...



...the change in area  $A(x + h) - A(x)$  can be approximated by the area of the thin rectangle of width  $h$  and height  $f(x)$ :

$$A(x + h) - A(x) \approx f(x)h.$$

Note that rearranging gives:

$$f(x) \approx \frac{A(x + h) - A(x)}{h}.$$

**Key idea: The thinner the rectangle, the better the approximation.**

Thus as  $h$  tends to zero we obtain

$$f(x) = \lim_{h \rightarrow 0} \frac{A(x + h) - A(x)}{h} = A'(x),$$

where the second equality follows directly from the **definition of the derivative**. Notice that this means that  $A(x)$  is an **antiderivative** of  $f(x)$ .

Since the area between the curve  $y = f(x)$ , the  $x$ -axis and the vertical lines  $x = a$  and  $x = b$  is equal to  $A(b) - A(a)$ , we note that this is given by the definite integral  $\int_a^b f(x) dx$ .

The relation between the two seemingly unrelated geometric problems, finding the area under the graph and finding the tangent, which links differentiation and integration, was Newton's extraordinary insight.

**Theorem 4**

If the curve  $y = f(x)$  lies entirely above the  $x$ -axis between  $x = a$  and  $x = b$ , then the area between the curve, the  $x$ -axis and the two vertical limits  $x = a$  and  $x = b$  is given by  $\int_a^b f(x) dx$ .

**Exercise**

Consider the constant function  $f(x) = 3$ . Make a sketch of the function  $y = f(x)$ .

For any constants  $a$  and  $b$ , it is clear that the area *between* the graph, the  $x$ -axis and the vertical lines  $x = a$  and  $x = b$  is equal to  $(b - a) \times 3 = 3b - 3a$ , since this is the area of a rectangle of width  $b - a$  and height 3, agreeing with the formula above:

$$\int_a^b 3 \cdot dx = [3x]_a^b = 3b - 3a.$$

If the curve  $y = f(x)$  lies entirely *below* the  $x$ -axis between  $x = a$  and  $x = b$ , then  $\int_a^b f(x) dx$  instead yields a negative value equal to minus 1 times the area between the curve, the  $x$ -axis and the two vertical limits  $x = a$  and  $x = b$ .

**Exercise**

Consider the constant function  $f(x) = -3$ . Make a sketch of the function  $y = f(x)$ .

For any constants  $a$  and  $b$ , it is clear that the area *between* the graph, the  $x$ -axis and the vertical lines  $x = a$  and  $x = b$  is again equal to  $3b - 3a$ , whilst the definite integral is equal to:

$$\int_a^b -3 \cdot dx = [-3x]_a^b = (-3b) - (-3a) = -3b + 3a = -(3b - 3a).$$



**Exercise**

Calculate the integral  $\int_{-2}^5 x \, dx$  and interpret your answer in terms of the areas between the curve and the  $x$ -axis.

**Solution:** It is straightforward to calculate that

$$\int_{-2}^5 x \, dx = \left[ \frac{1}{2}x^2 \right]_{-2}^5 = \frac{1}{2}(5^2 - (-2)^2) = 10.5$$


Make a sketch of the curve  $y = x$ .

The graph  $y = x$  is a straight line with slope 1. It lies below the  $x$ -axis for values of  $x$  between  $-2$  and  $0$ . It lies above the  $x$ -axis for values of  $x$  between  $0$  and  $5$ . The given indefinite integral therefore represents the difference of the area beneath the line  $y = x$  between  $x = 0$  and  $x = 5$ , and the area above the line  $y = x$  between  $x = -2$  and  $x = 0$ . Notice that this area can also be found (using your sketch) as the difference of the areas of the two right-angled triangles of heights  $5$  and  $2$  respectively and of the same bases, again giving  $\frac{1}{2}(5^2) - \frac{1}{2}(2^2) = 10.5$ .

**Example**

Find the area under the curve  $y = \sin(x)$  between  $x = 0$  and  $x = \pi$ .

**Solution:** The curve lies above the  $x$ -axis between  $x = 0$  and  $x = \pi$



Thus the area can be found as follows:

$$\int_0^{\pi} \sin(x) \, dx = [-\cos(x)]_0^{\pi} = -\cos(\pi) + \cos(0) = 1 + 1 = 2,$$

showing that the area is 2.

**Example**

Find the two places where the curve  $y = 2x - x^2$  crosses the  $x$ -axis, and hence calculate the enclosed area.

**Solution:** First make a sketch of  $y = 2x - x^2$ . Solving  $2x - x^2 = 0$  we find that the curve crosses the  $x$ -axis at  $x = 0$  and  $x = 2$ . Note that when  $x = 1$ ,  $y = 1 > 0$ , so the graph *peaks* somewhere between  $x = 0$  and  $x = 2$ .

Now integrating between these values:

$$\int_0^2 (2x - x^2) \, dx = \left[ x^2 - \frac{1}{3}x^3 \right]_0^2 = \left( 4 - \frac{8}{3} \right) - (0 - 0) = \frac{4}{3}$$

The following properties (which can be easily deduced from the definition) may be useful when calculating areas.

**Theorem 5**

For constants  $a$ ,  $b$  and  $c$ ,


$$\int_a^b f(x) dx + \int_b^c f(x) dx = \int_a^c f(x) dx.$$

$$\int_b^a f(x) dx = - \int_a^b f(x) dx.$$

$$\int_a^a f(x) dx = 0.$$

**Example**

Find the area of the shaded region.



**Solution:** The shaded area can be divided into two regions:

(1) the region above the  $x$ -axis;

(2) the region below the  $x$ -axis.

Notice that area of the first region can be found by computing

$$\int_0^1 x dx - \int_0^1 x^3 dx = \left[ \frac{1}{2}x^2 \right]_0^1 - \left[ \frac{1}{4}x^4 \right]_0^1 = \left( \frac{1}{2} - 0 \right) - \left( \frac{1}{4} - 0 \right) = \frac{1}{4}.$$

The area of the second region is also equal to  $\frac{1}{4}$  (by symmetry). It can be found by computing

$$\int_0^{-1} x dx - \int_0^{-1} x^3 dx = \left[ \frac{1}{2}x^2 \right]_0^{-1} - \left[ \frac{1}{4}x^4 \right]_0^{-1} = \left( \frac{1}{2} - 0 \right) - \left( \frac{1}{4} - 0 \right) = \frac{1}{4}.$$

## 1.4 Integration by substitution

If an integral appears difficult on first inspection, progress can often be made using a *substitution*. This method of solving integrals is known as the *change of variable* method or **integration by substitution** and in a sense can be thought of as the ‘reverse’ process to differentiating using the chain rule. The key idea is to choose a substitution so that the new integral (in terms of a new variable  $u$ ) is simpler than the original integral. After solving it and obtaining a function of  $u$ , we can then back-substitute the expression for  $u$  (given as a function of  $x$ ) in order to obtain the final answer as a function of  $x$ . This is best understood by means of a few examples.

### Example

To find  $\int e^{7x} dx$  we can make the substitution  $u = 7x$ . Then we have  $\frac{du}{dx} = 7$ , giving  $dx = \frac{1}{7} du$  and hence

$$\int e^{7x} dx = \int e^u \frac{1}{7} du = \frac{1}{7} \int e^u du = \frac{1}{7} e^u + C = \frac{1}{7} e^{7x} + C.$$

### Example

To find  $\int \sin(3x + 5) dx$  we can make the substitution  $u = 3x + 5$ . Then we have  $\frac{du}{dx} = 3$  or  $dx = \frac{1}{3} du$  and hence

$$\begin{aligned} \int \sin(3x + 5) dx &= \int \sin(u) \frac{1}{3} du = \frac{1}{3} \int \sin(u) du \\ &= -\frac{1}{3} \cos(u) + C \\ &= -\frac{1}{3} \cos(3x + 5) + C. \end{aligned}$$

The previous two examples illustrate a general principle:

### Theorem 6

Let  $f(x)$  be a function of  $x$  and let  $u = ax + b$  be a (non-constant) linear function of  $x$ . Then

$$\int f(ax + b) dx = \frac{1}{a} \int f(u) du.$$

(This can be seen by using the substitution  $u = ax + b$ .)

**Example**

Find  $\int \tan(x) dx$ .

**Solution:** By definition of the tangent function we have:

$$\int \tan(x) dx = \int \frac{\sin(x)}{\cos(x)} dx.$$

Consider the substitution  $u = \cos(x)$ . Then  $\frac{du}{dx} = -\sin(x)$ , giving  $\sin(x) dx = -du$  and so

$$\int \tan(x) dx = \int \frac{\sin(x) dx}{\cos(x)} = \int \frac{-du}{u} = -\int \frac{1}{u} du = -\ln |u| + C = -\ln |\cos(x)| + C.$$

**Example**

Find  $\int \frac{2x}{1+x^2} dx$ .

**Solution:** Let  $u = 1 + x^2$ . Then  $\frac{du}{dx} = 2x$  giving  $du = 2x dx$  and so

$$\int \frac{2x}{1+x^2} dx = \int \frac{2x dx}{1+x^2} = \int \frac{du}{u} = \ln |u| + C = \ln(1+x^2) + C.$$

The previous two examples illustrate another general principle:

**Theorem 7**

Let  $f(x)$  be a function of  $x$ . Then

$$\int \frac{f'(x)}{f(x)} dx = \int \frac{f'(x) dx}{f(x)} = \ln |f(x)| + C.$$

(This can be seen by using the substitution  $u = f(x)$ .)

**Example**

To find  $\int 2xe^{x^2} dx$  we can make the substitution  $u = x^2$ . Then  $\frac{du}{dx} = 2x$  giving  $du = 2x dx$  and hence

$$\int 2xe^{x^2} dx = \int e^{x^2} 2x dx = \int e^u du = e^u + C = e^{x^2} + C.$$

**Example**

Find

$$\int \frac{x}{\sqrt{1+x^2}} dx.$$

**Solution:** Let  $u = 1 + x^2$ , then  $\frac{du}{dx} = 2x$  giving  $du = 2x dx$  and so

$$\begin{aligned} \int \frac{x}{\sqrt{1+x^2}} dx &= \int \frac{x dx}{\sqrt{1+x^2}} = \int \frac{\frac{1}{2} du}{\sqrt{u}} \\ &= \frac{1}{2} \int u^{-1/2} du = \frac{1}{2} \frac{1}{(1/2)} u^{1/2} + C \\ &= u^{1/2} + C = \sqrt{1+x^2} + C. \end{aligned}$$

**Warning:** When we have to evaluate a definite integral such as  $\int_a^b f(x) dx$  by using a substitution, we need to take care of the limits of integration. Two approaches are possible:

(1) First solve the indefinite integral working as before (by substitution and then returning to the original variable  $x$ ), obtaining

$$\int f(x) dx = F(x) + C,$$

and then evaluate the definite integral as usual

$$\int_{x=a}^{x=b} f(x) dx = F(b) - F(a). ;$$

(2) Work with the definite integral directly and *change the limits of integration accordingly* when we introduce the new variable  $u$ . In the second approach, the formula for a change of variable has the form

$$\int_{x=a}^{x=b} f(x) dx = \int_{u=\alpha}^{u=\beta} f(x(u)) \frac{dx}{du} du \quad \text{where } \alpha = u(a), \beta = u(b).$$

(To avoid any confusion, we can write the limits showing the variable explicitly, as  $x = a$  and  $x = b$  in the original integral, and  $u = \alpha$  and  $u = \beta$  after the substitution.)

**Example**

Find

$$\int_1^5 (2x + 3)^3 dx.$$

**Solution:** Let  $u = 2x + 3$ . Then  $\frac{du}{dx} = 2$ , so  $dx = \frac{1}{2} du$ . Finding the indefinite integral and eliminating  $u$  in favor of  $x$  gives

$$\int (2x + 3)^3 dx = \frac{1}{2} \int u^3 du = \frac{1}{8} u^4 + C = \frac{1}{8} (2x + 3)^4 + C.$$

Evaluating using the limits of  $x$  now gives

$$\int_{x=1}^{x=5} (2x + 3)^3 dx = \left[ \frac{1}{8} (2x + 3)^4 \right]_1^5 = \frac{1}{8} (13^4 - 5^4) = 3492$$

---

Alternatively, in making the substitution and changing the variable of integration, we must *change the limits* accordingly. If  $x = 1$ , then  $u = 5$ , and if  $x = 5$ , then  $u = 13$ . The integral becomes

$$\int_{x=1}^{x=5} (2x + 3)^3 dx = \frac{1}{2} \int_{u=5}^{u=13} u^3 du = \left[ \frac{1}{8} u^4 \right]_5^{13} = \frac{1}{8} (28561 - 625) = 3492.$$

giving the same answer.

# Chapter 2

## Trigonometry

2.1 Trigonometric functions

2.2 Trigonometric identities and integration

2.3 Inverse trigonometric functions


### Revision from last semester..

1 radian = the angle which makes an arc length of  $r$  on a circle of radius  $r$

Circumference of a circle of radius  $r$  is  $2\pi r$

Area of a circle of radius  $r$  is  $\pi r^2$

Equation of a circle of radius  $r$   
centred on the origin is  $x^2 + y^2 = r^2$



Angles can be measured in **degrees** (e.g. a full circle is  $360^\circ$ , a half circle is  $180^\circ$  and a right-angle or quarter circle is  $90^\circ$ ) or in **radians** (e.g. a full circle is  $2\pi$  rad, a half circle is  $\pi$  rad and a right-angle is  $\frac{\pi}{2}$  rad).

In this course we will use most often use **radians**, as this is required by the formulas for derivatives and integrals of trigonometric functions. The notation for the unit "rad" is commonly omitted. Thus if no units are indicated, you can assume that we are working in radians. Make sure that you understand the difference between these two units and that you know how to convert one into the other.



E.g. An angle of  $135^\circ$  represents  $\frac{135}{360} = \frac{3}{8}$  of a full circle.

The corresponding angle in radians is  $\frac{3}{8}$  of  $2\pi$ , or  $\frac{3\pi}{4}$ .



## 2.1 Trigonometric functions

A point on the unit circle (that is the circle of radius 1 centred on the origin) can be specified by an angle  $\theta$ , measured anti-clockwise from the  $x$ -axis as indicated in the diagram below. Consider the co-ordinates of such a point:







$\cos(\theta) =$ the (horizontal) $x$ -co-ordinate of the point on the unit circle making an angle of $\theta$	$\sec(\theta) = \frac{1}{\cos(\theta)}$
$\sin(\theta) =$ the (vertical) $y$ -co-ordinate of the point on the unit circle making an angle of $\theta$	$\operatorname{cosec}(\theta) = \frac{1}{\sin(\theta)}$
$\tan(\theta) = \frac{\sin(\theta)}{\cos(\theta)}$	$\cot(\theta) = \frac{\cos(\theta)}{\sin(\theta)}$

These functions are called **trigonometric functions**. Using the above definitions, it is easy to check that these functions are periodic satisfying the following relations:

After one full turn we arrive at the same point on the circle	
$\cos(\theta + 2\pi) = \cos(\theta)$ (same $x$ -co-ordinate)	$\sin(\theta + 2\pi) = \sin(\theta)$ (same $y$ -co-ordinate)
After one half turn we arrive at the opposite point on the circle	
$\cos(\theta + \pi) = -\cos(\theta)$ ( $x$ -co-ordinate changes sign)	$\sin(\theta + \pi) = -\sin(\theta)$ ( $y$ -co-ordinate changes sign)
Reflecting in the (horizontal) $x$ -axis negates the $y$ -co-ordinate	
$\cos(2\pi - \theta) = \cos(\theta)$ (same $x$ -co-ordinate)	$\sin(2\pi - \theta) = -\sin(\theta)$ ( $y$ -co-ordinate changes sign)
Reflecting in the (vertical) $y$ -axis negates the $x$ -co-ordinate	
$\cos(\pi - \theta) = -\cos(\theta)$ ( $x$ -co-ordinate changes sign)	$\sin(\pi - \theta) = \sin(\theta)$ (same $y$ -co-ordinate)

The graphs of  $\sin(\theta)$ ,  $\cos(\theta)$  and  $\tan(\theta)$  are shown below:





The diagram above gives the following additional relations

$$\cos\left(\frac{\pi}{2} - \theta\right) = \sin(\theta) \text{ and } \sin\left(\frac{\pi}{2} - \theta\right) = \cos(\theta).$$

Notice that


$$\cos\left(\theta + \frac{\pi}{2}\right) = \cos\left(\pi - \left(\frac{\pi}{2} - \theta\right)\right) = -\cos\left(\frac{\pi}{2} - \theta\right) = -\sin(\theta)$$

and


$$\sin\left(\theta + \frac{\pi}{2}\right) = \sin\left(\pi - \left(\frac{\pi}{2} - \theta\right)\right) = \sin\left(\frac{\pi}{2} - \theta\right) = \cos(\theta)$$

### Remark


Three useful right-angled triangles are:



$$\begin{aligned} 30^\circ &= \frac{\pi}{6} \\ \sin\left(\frac{\pi}{6}\right) &= \frac{1}{2} \\ \cos\left(\frac{\pi}{6}\right) &= \frac{\sqrt{3}}{2} \\ \tan\left(\frac{\pi}{6}\right) &= \frac{1}{\sqrt{3}} \end{aligned}$$



$$\begin{aligned} 45^\circ &= \frac{\pi}{4} \\ \sin\left(\frac{\pi}{4}\right) &= \frac{1}{\sqrt{2}} \\ \cos\left(\frac{\pi}{4}\right) &= \frac{1}{\sqrt{2}} \\ \tan\left(\frac{\pi}{4}\right) &= 1 \end{aligned}$$



$$\begin{aligned} 60^\circ &= \frac{\pi}{3} \\ \sin\left(\frac{\pi}{3}\right) &= \frac{\sqrt{3}}{2} \\ \cos\left(\frac{\pi}{3}\right) &= \frac{1}{2} \\ \tan\left(\frac{\pi}{3}\right) &= \sqrt{3} \end{aligned}$$

## 2.2 Trigonometric identities and integration

### Remark

We use the following notation for **positive powers** of trigonometric functions:

$$\begin{aligned}\sin^2(\theta) &= \sin(\theta)\sin(\theta) = (\sin(\theta))^2 \\ \sin^3(\theta) &= \sin(\theta)\sin(\theta)\sin(\theta) = (\sin(\theta))^3 \\ &\vdots \\ \sin^n(\theta) &= \underbrace{\sin(\theta)\sin(\theta)\cdots\sin(\theta)}_{n \text{ times}} = (\sin(\theta))^n\end{aligned}$$

The notations  $\cos^n(\theta)$ ,  $\tan^n(\theta)$ ,  $\operatorname{cosec}^n(\theta)$ ,  $\sec^n(\theta)$  and  $\cot^n(\theta)$  are defined similarly.

We recall the following fundamental identity:

### Theorem 8

For all angles  $\theta$ ,  $\cos^2(\theta) + \sin^2(\theta) = 1$ .

### Proof

Each point on the unit circle has co-ordinates  $(\cos(\theta), \sin(\theta))$  for some angle  $\theta$ , so the result follows immediately from the equation of the unit circle:  $x^2 + y^2 = 1^2$ .  $\square$

Trigonometric identities can help with integration:

### Example

Find  $\int \cos^2(x) + \sin^2(x) dx$ .

**Solution:** Since  $\cos^2(x) + \sin^2(x) = 1$ , we may write the integral as:

$$\int \cos^2(x) + \sin^2(x) dx = \int 1 dx = x + C$$

### Corollary 9

For all values of  $\theta$  for which the functions are defined:

$$\begin{aligned}1 + \tan^2(\theta) &= \sec^2(\theta) \\ \cot^2(\theta) + 1 &= \operatorname{cosec}^2(\theta)\end{aligned}$$

**Proof**

These two formulas are obtained by dividing the previous result through by  $\cos^2\theta$  and  $\sin^2\theta$  respectively.  $\square$

**Example**

Find  $\int 1 + \tan^2(2x) dx$ .

**Solution:** Since  $1 + \tan^2(u) = \sec^2(u)$ , we may write the integral as:

$$\int 1 + \tan^2(2x) dx = \int \sec^2(2x) dx = \frac{1}{2} \int \sec^2(u) du = \frac{1}{2} \tan(u) + C = \frac{1}{2} \tan(2x) + C,$$

by using the substitution  $u = 2x$ .

The following results can also be derived geometrically (see 0C1 notes for details):

**Theorem 10** (Addition formulas)

For any angles  $A$  and  $B$ ,

$$\sin(A + B) = \sin(A) \cos(B) + \cos(A) \sin(B)$$

$$\cos(A + B) = \cos(A) \cos(B) - \sin(A) \sin(B)$$

$$\sin(A - B) = \sin(A) \cos(B) - \cos(A) \sin(B)$$

$$\cos(A - B) = \cos(A) \cos(B) + \sin(A) \sin(B)$$

**Example**

Find  $\int \sin(x + 5) \cos(2x) + \cos(x + 5) \sin(2x) dx$ .

**Solution:** Using the addition formula for sine above we see that  $\sin(x + 5) \cos(2x) + \cos(x + 5) \sin(2x) = \sin(x + 5 + 2x)$ . Thus we may write the integral as:

$$\int \sin(3x + 5) dx = -\frac{1}{3} \cos(3x + 5) + C,$$

by using the substitution  $u = 3x + 5$ , or by inspection.

We have seen that it is useful (e.g. for integration) to be able to work with identities in order to simplify expressions involving trigonometric functions. We can deduce further useful identities from the formulas  $\sin(A + B)$  and  $\cos(A + B)$ .

**Theorem 11** (Addition formulas for tangent)

$$\tan(A + B) = \frac{\tan(A) + \tan(B)}{1 - \tan(A) \tan(B)}$$

$$\tan(A - B) = \frac{\tan(A) - \tan(B)}{1 + \tan(A) \tan(B)}$$

**Proof**

Use the addition formula for sine and cosine:

$$\begin{aligned} \tan(A + B) &= \frac{\sin(A + B)}{\cos(A + B)} = \frac{\sin(A)\cos(B) + \sin(B)\cos(A)}{\cos(A)\cos(B) - \sin(A)\sin(B)} \\ &= \frac{\cos(A)\cos(B) \left( \frac{\sin(A)}{\cos(A)} + \frac{\sin(B)}{\cos(B)} \right)}{\cos(A)\cos(B) \left( 1 - \frac{\sin(A)}{\cos(A)} \frac{\sin(B)}{\cos(B)} \right)} = \frac{\tan(A) + \tan(B)}{1 - \tan(A) \tan(B)} \end{aligned}$$

The second formula follows in the same way. □

**Theorem 12** (Double angle formulas)

$$\begin{aligned} \sin(2\theta) &= 2 \sin(\theta) \cos(\theta) \\ \cos(2\theta) &= \cos^2(\theta) - \sin^2(\theta) \\ &= 2\cos^2(\theta) - 1 \\ &= 1 - 2\sin^2(\theta). \end{aligned}$$

**Proof**

Let  $\theta = A = B$  in the sum identities. This gives:

$$\begin{aligned} \sin(2\theta) &= \sin(\theta + \theta) = \sin(\theta) \cos(\theta) + \cos(\theta) \sin(\theta) = 2\sin(\theta)\cos(\theta) \\ \cos(2\theta) &= \cos(\theta + \theta) = \cos(\theta) \cos(\theta) - \sin(\theta) \sin(\theta) = \cos^2(\theta) - \sin^2(\theta) \end{aligned}$$

Finally we note that using the identity  $\cos^2(\theta) + \sin^2(\theta) = 1$  gives

$$\begin{aligned} 2\cos^2(\theta) - 1 &= 2\cos^2(\theta) - (\cos^2(\theta) + \sin^2(\theta)) = \cos^2(\theta) - \sin^2(\theta) = \cos(2\theta) \\ 1 - 2\sin^2(\theta) &= (\cos^2(\theta) + \sin^2(\theta)) - 2\sin^2(\theta) = \cos^2(\theta) - \sin^2(\theta) = \cos(2\theta) \end{aligned}$$

□

**Example**

Find  $\int \sin(x) \cos(x) dx$ .

**Solution:** Using the double angle formula for sine given above we see that

$$\sin(x) \cos(x) = \frac{1}{2} \sin(2x)$$

Thus

$$\int \sin(x) \cos(x) dx = \int \frac{1}{2} \sin(2x) dx = -\frac{1}{4} \cos(2x) + C$$

**Example**

Find  $\int \cos^2(x) dx$ .

**Solution:** Using the double angle formula for cosine given above we see that

$$\cos^2(x) = \frac{1}{2} \cos(2x) + \frac{1}{2}$$

Thus

$$\int \cos^2(x) dx = \int \frac{1}{2} \cos(2x) dx + \int \frac{1}{2} dx = \frac{1}{4} \sin(2x) + \frac{1}{2}x + C$$

From the formulas for multiple angles (such as  $\cos(nx)$  and  $\sin(nx)$ ), we can deduce formulas for powers (such as  $\cos^n(x)$  and  $\sin^n(x)$ ). We consider some particular examples to illustrate this idea.

**Example**

Express  $\cos(3x)$  as a polynomial in  $\cos(x)$ .

**Solution:**

$$\begin{aligned} \cos(3x) &= \cos(2x + x) = \cos(2x) \cos(x) - \sin(2x) \sin(x) \\ &= (2\cos^2(x) - 1)\cos(x) - (2\sin(x)\cos(x))\sin(x) \\ &= 2\cos^3(x) - \cos(x) - 2\sin^2(x)\cos(x) \\ &= 2\cos^3(x) - \cos(x) - 2(1 - \cos^2(x))\cos(x) \\ &= 4\cos^3(x) - 3\cos(x). \end{aligned}$$

**Example**

Find  $\int \cos^3(x) dx$ .

**Solution:** We have seen that  $\cos(3x) = 4\cos^3(x) - 3\cos(x)$ .

Thus

$$\cos^3(x) = \frac{1}{4} (\cos(3x) + 3\cos(x)) .$$

Since we know how to integrate each of the functions on the right-hand side:

$$\begin{aligned} \int \cos^3(x) dx &= \int \frac{1}{4} (\cos(3x) + 3\cos(x)) dx = \frac{1}{4} \int \cos(3x) dx + \frac{3}{4} \int \cos(x) dx \\ &= \frac{1}{12} \sin(3x) + \frac{3}{4} \sin(x) + C. \end{aligned}$$

**Example**

Show that  $\sin(3x) = 3 \cos^2(x) \sin(x) - \sin^3(x)$  and hence find  $\int \sin^3(x) dx$ .

**Solution:**

$$\begin{aligned} \sin(3x) &= \sin(2x + x) = \sin(2x) \cos(x) + \cos(2x) \sin(x) \\ &= 2 \sin(x) \cos^2(x) + (\cos^2(x) - \sin^2(x)) \sin(x) \\ &= 3 \sin(x) \cos^2(x) - \sin^3(x) \end{aligned}$$

Rearranging gives  $\sin^3(x) = 3 \sin(x) \cos^2(x) - \sin(3x)$  and hence

$$\int \sin^3(x) dx = 3 \int \sin(x) \cos^2(x) dx - \int \sin(3x) dx = -\cos^3(x) + \frac{1}{3} \cos(3x) + C.$$

(The first integral can be found using the substitution  $u = \cos(x)$ . The second integral can be found using the substitution  $u = 3x$ .)

**Theorem 13**

$$\sin(X) + \sin(Y) = 2 \sin\left(\frac{X+Y}{2}\right) \cos\left(\frac{X-Y}{2}\right)$$

$$\sin(X) - \sin(Y) = 2 \cos\left(\frac{X+Y}{2}\right) \sin\left(\frac{X-Y}{2}\right)$$

$$\cos(X) + \cos(Y) = 2 \cos\left(\frac{X+Y}{2}\right) \cos\left(\frac{X-Y}{2}\right)$$

$$\cos(X) - \cos(Y) = -2 \sin\left(\frac{X+Y}{2}\right) \sin\left(\frac{X-Y}{2}\right)$$

**Proof**



Using the sum and difference formulas for sine we find that:

$$\begin{aligned}\sin(A+B) + \sin(A-B) &= 2 \sin(A) \cos(B) \\ \sin(A+B) - \sin(A-B) &= 2 \cos(A) \sin(B).\end{aligned}$$

Now setting  $X = A + B$  and  $Y = A - B$ , so that  $A = \frac{1}{2}(X + Y)$  and  $B = \frac{1}{2}(X - Y)$ , gives our first two identities:

$$\begin{aligned}\sin(X) + \sin(Y) &= 2 \sin\left(\frac{X+Y}{2}\right) \cos\left(\frac{X-Y}{2}\right) \\ \sin(X) - \sin(Y) &= 2 \cos\left(\frac{X+Y}{2}\right) \sin\left(\frac{X-Y}{2}\right)\end{aligned}$$

Similarly, using the the sum and difference formulas for cosine gives:

$$\begin{aligned}\cos(A+B) + \cos(A-B) &= 2 \cos A \cos B \\ \cos(A+B) - \cos(A-B) &= -2 \sin A \sin B.\end{aligned}$$

and setting  $X = A + B$  and  $Y = A - B$  gives the second two identities. □

### Example

Find  $\int 2 \sin(x) \cos(x + \frac{\pi}{6}) dx$

**Solution:** Let  $x = \frac{X+Y}{2}$  and  $x + \frac{\pi}{6} = \frac{X-Y}{2}$ , so that the integrand resembles the identity for  $\sin(X) + \sin(Y)$  given above. Now find expressions for  $X$  and  $Y$  in terms of  $x$ . Since

$$\frac{X+Y}{2} = x \text{ and } \frac{X-Y}{2} = x + \frac{\pi}{6}$$

we can solve for  $X$  by adding together the two equations to obtain  $X = 2x + \frac{\pi}{6}$ . Similarly we can solve for  $Y$  by subtracting the two equations to obtain  $Y = -\frac{\pi}{6}$ . Now the expression we wish to integrate becomes

$$\begin{aligned}2 \sin(x) \cos(x + \frac{\pi}{6}) &= 2 \sin\left(\frac{X+Y}{2}\right) \cos\left(\frac{X-Y}{2}\right) = \sin(X) + \sin(Y) \\ &= \sin(2x + \frac{\pi}{6}) + \sin(-\frac{\pi}{6}) \\ &= \sin(2x + \frac{\pi}{6}) - \frac{1}{2}.\end{aligned}$$

Hence

$$\int 2 \sin(x) \cos(x + \frac{\pi}{6}) dx = \int \sin(2x + \frac{\pi}{6}) dx - \int \frac{1}{2} dx = -\frac{1}{2} \cos(2x + \frac{\pi}{6}) - \frac{1}{2}x + C.$$

We conclude this section with two more useful identities:

**Theorem 14**

$$\sin(2\theta) = \frac{2\tan(\theta)}{1 + \tan^2(\theta)}, \quad \cos(2\theta) = \frac{1 - \tan^2(\theta)}{1 + \tan^2(\theta)}.$$

**Proof**

Dividing the double angle formulas through by  $1 = \cos^2(\theta) + \sin^2(\theta)$  gives:

$$\begin{aligned} \sin(2\theta) &= 2\sin(\theta)\cos(\theta) = \frac{2\sin(\theta)\cos(\theta)}{\cos^2(\theta) + \sin^2(\theta)} = \frac{2\cos^2(\theta)\frac{\sin\theta}{\cos\theta}}{\cos^2(\theta)\left(1 + \frac{\sin^2\theta}{\cos^2\theta}\right)} = \frac{2\tan(\theta)}{1 + \tan^2(\theta)}. \\ \cos(2\theta) &= \cos^2(\theta) - \sin^2(\theta) = \frac{\cos^2(\theta) - \sin^2(\theta)}{\cos^2(\theta) + \sin^2(\theta)} = \frac{1 - \tan^2(\theta)}{1 + \tan^2(\theta)}. \end{aligned}$$

□

**Remark**

The above group of formulas is often re-written in terms of  $x = 2\theta$  and referred to as “tangent half-angle formulas”:

$$\sin(x) = \frac{2\tan\left(\frac{x}{2}\right)}{1 + \tan^2\left(\frac{x}{2}\right)}, \quad \cos(x) = \frac{1 - \tan^2\left(\frac{x}{2}\right)}{1 + \tan^2\left(\frac{x}{2}\right)}.$$

## 2.3 Inverse trigonometric functions

We begin by recalling the idea of an **inverse function**.

**Example**

The functions  $y = e^x$  and  $y = \ln(x)$  are **mutually inverse functions**.

This means that:

(1) For all  $x > 0$ , if we first take  $\ln(x)$  and then apply the exponential, we return to the original value of  $x$ :

$$e^{(\ln x)} = x.$$

(2) For all  $x$ , if we first take  $e^x$  and then apply the logarithm, we return to the original value of  $x$ :

$$\ln(e^x) = x.$$

Each function “undoes” the other.

“Undoing” a function it is not always so simple.

**Example**

Consider the functions  $y = x^2$  and  $y = \sqrt{x}$ .

It is natural (but incorrect!) to think that these are mutually inverse functions. On the one hand, squaring “undoes” the square root function:

$$(\sqrt{x})^2 = x \text{ for all } x > 0.$$

But what about the other way around? Is it true that  $\sqrt{x^2} = x$ ?

**No!** Every positive number has *two* square roots, one positive and one negative. (The symbol  $\sqrt{x}$  is usually interpreted as the *positive* square root and we write  $\pm\sqrt{x}$  to denote the *pair* of square roots.) Since  $\sqrt{x} \geq 0$  for all  $x \geq 0$  we see that

$$\sqrt{x^2} = \begin{cases} x & \text{for } x \geq 0 \\ -x & \text{for } x \leq 0 \end{cases}$$

E.g.  $(\sqrt{((-1)^2)}) = 1 = -(-1)$ .

To summarize, given a function  $f(x)$ , the function which “undoes”  $f$  sends  $y$  to  $x$  if  $f(x) = y$ . In general there may or may not be such a function. When we solve an equation of the form  $f(x) = y$  for  $x$  (where  $y$  is given), two things may happen:

1. Solutions may not exist for some values of  $y$ ;
2. For a given value of  $y$ , there may be many solutions  $x$ .

The first case tells us that we need to think a bit carefully about which values we can apply our “undoing function” to. For example, since  $e^x = y > 0$  for all  $x$ , this tells us that  $\ln(x)$  only makes sense for  $x > 0$ .

The second case causes more severe problems when trying to define the inverse, because for some values of  $y$  there are many choices for  $x$ . Such functions are called “many-valued”. Typically, one particular solution  $x$  is chosen as the *principal value* and other solutions are expressed in terms of the principal value.

**Example**

We have seen that the function which “undoes”  $f(x) = x^2$  is many-valued. All solutions of the equation

$$x^2 = y$$

are expressed as

$$x = \pm\sqrt{y}$$

Thus we consider the positive square roots  $\sqrt{y}$  to be the *principal values*.

**Definition** (Inverse trigonometric functions)

Consider the graphs  $y = \sin(x)$ ,  $y = \cos(x)$  and  $y = \tan(x)$ .

The functions which “undo” these trigonometric functions are many-valued.

We write:

$x = \text{Arcsin}(y)$  to denote the angles  $x$  such that  $\sin(x) = y$ .

$x = \text{Arccos}(y)$  to denote the angles  $x$  such that  $\cos(x) = y$ .


$x = \text{Arctan}(y)$  to denote the angles  $x$  such that  $\tan(x) = y$ .

Notice that the input vales  $y$  for  $\text{Arcsin}(y)$  and  $\text{Arccos}(y)$  must be between  $-1$  and  $1$  (since the output values for sine and cosine do not lie outside this range) , whilst the argument  $y$  for  $\text{Arctan}(y)$  can be any number (since the output of the tangent function covers all real numbers).

**Example**

Find  $\text{Arcsin}(\frac{\sqrt{3}}{2})$ .

**Solution:** We need to find **all values** of  $\theta$  satisfying  $\sin(\theta) = \frac{\sqrt{3}}{2}$ .



There are two points on the circle with vertical co-ordinate equal to  $\frac{\sqrt{3}}{2}$ ; one corresponding to an angle of  $\theta = \frac{\pi}{3}$ , and one corresponding to an angle of  $\pi - \theta = \frac{2\pi}{3}$ .

Therefore  $\text{Arcsin}(\frac{\sqrt{3}}{2}) = \frac{\pi}{3} + 2k\pi, \frac{2\pi}{3} + 2k\pi, \quad \text{where } k = 0, \pm 1, \pm 2, \pm 3, \dots$

In general, we note that  $\text{Arcsin}(c)$  (for  $-1 \leq c \leq 1$ ) can be found by considering the points on the unit circle with **vertical** co-ordinate equal to  $c$ .


If  $\theta$  is one such value, then

$$\text{Arcsin}(c) = \theta + 2k\pi, \quad \pi - \theta + 2k\pi, \quad \text{where } k = 0, \pm 1, \pm 2, \pm 3, \dots$$

**Example**

Find  $\text{Arccos}(\frac{\sqrt{3}}{2})$ .

**Solution:** We need to find **all values** of  $\theta$  satisfying  $\cos(\theta) = \frac{\sqrt{3}}{2}$ .



There are two points on the circle with horizontal co-ordinate equal to  $\frac{\sqrt{3}}{2}$ ; one corresponding to an angle of  $\theta = \frac{\pi}{6}$ , and one corresponding to an angle of  $-\theta = -\frac{\pi}{6}$ .

$$\text{Therefore } \text{Arccos}(\frac{\sqrt{3}}{2}) = \pm\frac{\pi}{6} + 2k\pi, \quad \text{where } k = 0, \pm 1, \pm 2, \pm 3, \dots$$

---

In general, we note that  $\text{Arccos}(c)$  (for  $-1 \leq c \leq 1$ ) can be found by considering the points on the unit circle with **horizontal** co-ordinate equal to  $c$ .

If  $\theta$  is one such value, then


$$\text{Arccos}(c) = \pm\theta + 2k\pi, \quad \text{where } k = 0, \pm 1, \pm 2, \pm 3, \dots$$

**Definition**

$y = \sin^{-1}(x)$  is the unique value of  $y = \text{Arcsin}(x)$  such that  $-\frac{1}{2}\pi \leq y \leq \frac{1}{2}\pi$ .

$y = \cos^{-1}(x)$  is the unique value of  $y = \text{Arccos}(x)$  such that  $0 \leq y \leq \pi$

$y = \tan^{-1}(x)$  is the unique value of  $y = \text{Arctan}(x)$  such that  $-\frac{1}{2}\pi \leq y \leq \frac{1}{2}\pi$ .



**Remark**

For all values of  $x$  we have:

$$\sin(\sin^{-1}(x)) = x, \quad \cos(\cos^{-1}(x)) = x, \quad \text{and} \quad \tan(\tan^{-1}(x)) = x.$$

**Warning:** For values of  $x$  outside the range of the given inverse function:

$$\sin^{-1}(\sin(x)) \neq x, \quad \cos^{-1}(\cos(x)) \neq x, \quad \tan^{-1}(\tan(x)) \neq x.$$

**Remark**

The inverse trigonometric functions  $\sin^{-1}(x)$ ,  $\cos^{-1}(x)$  and  $\tan^{-1}(x)$  are single-valued functions regarded as the **principal values** of the multi-valued functions  $\text{Arcsin}(x)$ ,  $\text{Arccos}(x)$  and  $\text{Arctan}(x)$ . (Their position is similar to that of the positive square root  $\sqrt{x}$ , in relation to the double-valued square root  $\pm\sqrt{x}$ .)

The multivalued functions  $\text{Arcsin}(x)$ ,  $\text{Arccos}(x)$ ,  $\text{Arctan}(x)$  can be expressed via their principal values as follows:

$$\begin{aligned}\text{Arcsin}(x) &= \sin^{-1}(x) + 2k\pi, & \pi - \sin^{-1}(x) + 2k\pi & \quad (k = 0, \pm 1, \pm 2, \pm 3, \dots) \\ \text{Arccos}(x) &= \pm \cos^{-1}(x) + 2k\pi & & \quad (k = 0, \pm 1, \pm 2, \pm 3, \dots), \\ \text{Arctan}(x) &= \tan^{-1}(x) + k\pi & & \quad (k = 0, \pm 1, \pm 2, \pm 3, \dots).\end{aligned}$$

**Warning:** Do not confuse the notation for inverse functions with the notation for positive powers!

$$(\sin(x))^{-1} \neq \sin^{-1}(x)$$

E.g.  $(\sin(\pi/2))^{-1} = 1$ , but since  $\frac{\pi}{2} > 1$  we cannot evaluate  $\sin^{-1}(\pi/2)$ .

(This is because there is no angle which gives a point on the unit circle with vertical co-ordinate equal to  $\frac{\pi}{2}$ .)

**Example**

Find all values of  $a$  such that  $\cos(a) = \frac{1}{2}$ .

**Solution:** We begin by finding the principal value  $\cos^{-1}(\frac{1}{2}) = \frac{\pi}{3}$ .

The solutions are then given by

$$a = \text{Arccos}\left(\frac{1}{2}\right) = \pm \cos^{-1}\left(\frac{1}{2}\right) + 2k\pi = \pm \frac{\pi}{3} + 2k\pi \quad (\text{where } k = 0, \pm 1, \pm 2, \dots).$$

(This can be seen by looking at our unit circle diagram, as in the previous example.)

**Example**

Find  $\theta$  in the range  $0 \leq \theta < \pi$  such that  $\tan(\theta) = 5$ .

**Solution:** The unique solution in the range between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$  is the principal value  $\theta = \tan^{-1}(5) \approx 1.37$  (this can be found using a calculator).

The general solution of the equation  $\tan(\theta) = 5$  is obtained by adding multiples of  $\pi$ . Since adding any multiple of  $\pi$  to  $\theta = \tan^{-1}(5)$  takes us out of the range  $0 \leq \theta < \pi$ , we see that there is just one solution.

Hence the answer is:  $\theta = \tan^{-1}(5) \approx 1.37$

We record some useful properties of the inverse trigonometric functions:

**Theorem 15** (Properties of inverse trigonometric functions)

$$\begin{aligned}\sin^{-1}(-x) &= -\sin^{-1}(x), \\ \cos^{-1}(-x) &= \pi - \cos^{-1}(x), \\ \tan^{-1}(-x) &= -\tan^{-1}(x).\end{aligned}$$

**Proof**

Note that both sides of the above identities take values in the same range. To prove them, it is sufficient to apply  $\sin$ ,  $\cos$  and  $\tan$  respectively to both sides.  $\square$

**Example**

Solve the equation  $\cos^2(x) = \frac{1}{7}$ .

**Solution:** Taking the square root gives two possibilities:

$$\begin{array}{l} \cos(x) = \frac{1}{\sqrt{7}} \\ x = \text{Arccos}\left(\frac{1}{\sqrt{7}}\right) \\ x = \pm\cos^{-1}\left(\frac{1}{\sqrt{7}}\right) + 2k\pi \end{array} \quad \left\| \quad \begin{array}{l} \cos(x) = -\frac{1}{\sqrt{7}} \\ x = \text{Arccos}\left(-\frac{1}{\sqrt{7}}\right) \\ x = \pm\cos^{-1}\left(-\frac{1}{\sqrt{7}}\right) + 2k\pi = \pm\left(\pi - \cos^{-1}\left(\frac{1}{\sqrt{7}}\right)\right) + 2k\pi \end{array}\right.$$

Putting this all together, the full set of solutions can also be written as:

$$x = \pm\cos^{-1}\left(\frac{1}{\sqrt{7}}\right) + k\pi \quad (\text{where } k = 0, \pm 1, \pm 2, \dots).$$

**Example**

Solve the equation:  $2\sin^2(x) + 5\sin(x) - 3 = 0$ .

**Solution:** We want to solve the equation  $2y^2 + 5y - 3 = 0$  where  $y = \sin(x)$ .

We begin by factorising the quadratic polynomial:  $2y^2 + 5y - 3 = (2y - 1)(y + 3) = 0$ .

Since the solutions to this equation are  $y = \frac{1}{2}$  and  $y = -3$ , we see that we must now solve  $\sin(x) = \frac{1}{2}$  and  $\sin(x) = -3$  for  $x$ .

It is easy to see that the second equation has no solutions, so it remains to solve  $\sin(x) = \frac{1}{2}$ .

The solutions are:

$$\begin{aligned}x &= \text{Arcsin}\left(\frac{1}{2}\right) \\ &= \sin^{-1}\left(\frac{1}{2}\right) + 2k\pi, \quad \pi - \sin^{-1}\left(\frac{1}{2}\right) + 2k\pi \\ &= \frac{\pi}{6} + 2k\pi, \quad \frac{5\pi}{6} + 2k\pi \quad \text{for } k = 0, \pm 1, \pm 2, \dots\end{aligned}$$



**Theorem 16** (Relations between inverse trigonometric functions)

$$\begin{aligned}\sin^{-1}(x) &= \frac{\pi}{2} - \cos^{-1}(x), \\ \sin^{-1}(x) &= \tan^{-1}\left(\frac{x}{\sqrt{1-x^2}}\right).\end{aligned}$$

**Proof**

Consider the first identity. Since  $\cos^{-1}(x)$  takes values between 0 and  $\pi$ , the expression  $\frac{\pi}{2} - \cos^{-1}(x)$  takes values between  $-\frac{\pi}{2}$  and  $\frac{\pi}{2}$  as is required for  $\sin^{-1}(x)$ . Therefore it is sufficient to check that  $\sin$  applied to the r.h.s. gives  $x$ . Indeed,

$$\sin\left(\frac{\pi}{2} - \cos^{-1}(x)\right) = \cos(\cos^{-1}(x)) = x.$$

So the l.h.s. is indeed equal to the r.h.s. For the second identity, note that the l.h.s. and r.h.s. take values in the same range by the definition of  $\sin^{-1}$  and  $\tan^{-1}$ . Therefore it is sufficient to check that  $\tan$  applied to both sides gives the same number. Indeed, if  $a = \sin^{-1}(x)$ , then  $\sin(a) = x$  and  $-\frac{\pi}{2} \leq a \leq \frac{\pi}{2}$ . In this range,  $\cos(a) = \sqrt{1 - \sin^2(a)} = \sqrt{1 - x^2}$ . Hence

$$\tan(\sin^{-1}(x)) = \tan(a) = \frac{\sin(a)}{\cos(a)} = \frac{x}{\sqrt{1-x^2}},$$

which completes the proof. □

# Chapter 3

## Sequences, series and summations

3.1 Arithmetic progressions

3.2 Geometric progressions

3.3 The binomial theorem

### Revision from last semester..

#### Powers:

For  $a, b$  real numbers and  $r, s$  rational numbers such that  $a^r, a^s$  etc. are defined:

$$a^0 = 1 \quad a^1 = a$$

$$a^r a^s = a^{r+s} \quad a^r / a^s = a^{r-s} \quad a^{-s} = 1/a^s$$

$$(a^r)^s = a^{rs} \quad (ab)^r = a^r b^r \quad (a/b)^r = a^r / b^r$$

#### Logarithms:

For  $a > 0, a \neq 1, a^x = y \Leftrightarrow x = \log_a(y)$ .

$$\log_a(1) = 0 \quad \log_a(a) = 1$$

$$\log_a(a^x) = x \quad a^{\log_a(b)} = b$$

$$\log_a(bc) = \log_a(b) + \log_a(c) \quad \log_a(b/c) = \log_a(b) - \log_a(c)$$

$$\log_a(b^y) = y \log_a(b) \quad \log_b(c) = \frac{\log_a(c)}{\log_a(b)}$$

## 3.1 Arithmetic progressions

### Definition (Sequences)

A **sequence** is an ordered list of numbers  $a_1, a_2, a_3, \dots$ , each of which is called a **term**. The  $k$ th term of the sequence is  $a_k$ .

A sequence may have finitely many terms or infinitely many terms.

### Example

$$2, 5, 10, 17, 26, \dots, k^2 + 1, \dots, 10001$$

is a finite sequence of 100 numbers.

If  $k$  is a whole number between 1 and 100, then the  $k$ th term of this sequence is  $a_k = k^2 + 1$ .

Given a sequence

$$a_1, a_2, a_3, a_4, \dots, a_k, a_{k+1}, \dots,$$

(finite or infinite), we can associate to it the corresponding sequence of *differences*:

$$d_1 = a_2 - a_1, \quad d_2 = a_3 - a_2, \quad d_3 = a_4 - a_3, \quad \dots, \quad d_k = a_{k+1} - a_k, \quad \dots$$

The sequence  $d_1, d_2, d_3, \dots$  describes the “growth” or “rate of change” of the original sequence  $a_1, a_2, a_3, \dots$ . Notice that this is a similar idea to taking the derivative<sup>1</sup> of a function. You may find it useful to think of a sequence as a function of  $k$  where  $k$  only takes discrete values ( $k = 1, 2, 3, \dots$ ).

### Example

If  $d_k = 0$  for all  $k$ , this means that  $a_{k+1} - a_k = 0$ , or in other words, all the terms  $a_k$  are the same,  $a_k = a$ . This is the analog of a constant function.

(Recall that if a function  $f(x)$  has derivative 0 everywhere, then it must be a constant.)

In this section we shall be interested in sequences whose differences are not necessarily all zero, but are the *same for all*  $k$ . This means that the sequence has “uniform growth”. Such sequences are analogous to linear functions and there is a special name for sequences of this type.

---

<sup>1</sup>Recall from last semester that the derivative is defined via the differences  $f(x+h) - f(x)$ ; we take the limit of  $(f(x+h) - f(x))/h$  when  $h \rightarrow 0$ . Unlike this situation, for sequences, the “step size” is fixed (it is always 1) so there is no way of taking such a limit, and we have to deal with the differences  $a_{k+1} - a_k$  directly.

**Definition** (Arithmetic progression)

An **arithmetic progression** is a sequence  $a_1, a_2, a_3, \dots$  in which the corresponding differences  $d_k = a_{k+1} - a_k$  are all the same.

We call this value the **common difference**  $d$ .

An arithmetic progression with common difference  $d$  and first term  $a$  has the form

$$a, \quad a + d, \quad a + 2d, \quad a + 3d, \quad \dots, \quad a + (k - 1)d, \dots$$

Each term is obtained by adding  $d$  to the previous term.

The  $k$ th term of this sequence is  $a_k = a + (k - 1)d$ .

**Example**

Each pair of consecutive terms in the sequence

$$5, 7, 9, 11, 13, 15, \dots, 77$$

differs by 2.

This is an arithmetic progression with first term  $a = 5$  and common difference  $d = 2$ .

We can calculate the number of terms in the sequence as follows:

Since  $a = 5$  and  $d = 2$ , we know that the last term of this sequence can be written as  $77 = 5 + (k - 1) \times 2$ , where  $k$  is the number of terms.

Rearranging this equation gives  $k - 1 = 72/2$ , and hence  $k = 37$ .

**Theorem 17**

**The sum of the first  $n$  terms of an arithmetic progression** with first term  $a$  and common difference  $d$  is

$$S_n = \frac{1}{2}n(2a + (n - 1)d).$$

**Proof**

Write out the sum twice: once forwards, once backwards, and then add them together:

$$\begin{array}{rcccccccc} S_n & = & a & & + & (a + d) & & + \cdots + & (a + (n - 2)d) & + & (a + (n - 1)d) \\ S_n & = & (a + (n - 1)d) & + & (a + (n - 2)d) & + \cdots + & (a + d) & + & a \end{array}$$

---


$$2S_n = (2a + (n - 1)d) + (2a + (n - 1)d) + \cdots + (2a + (n - 1)d) + (2a + (n - 1)d)$$

Note that each term on the right-hand side of the last line is equal to  $2a + (n - 1)d$ . Since there are  $n$  of these we see that  $2S_n = n(2a + (n - 1)d)$ , and dividing this by two gives the result.  $\square$

**Remark**

We can rewrite the formula for the sum of the first  $n$  terms of an arithmetic progression in the following easy to remember form:

$$S_n = \frac{1}{2}n(a_1 + a_n).$$

*We add together the first term and the last term, multiply the result by the number of terms and then divide by 2.*

(The first term is  $a$ , whilst the last ( $n$ th) term is  $a_n = a + (n - 1)d$ . So their sum is equal to the bracketed expression in the previous result.)

**Example**

Find the sum:

$$5 + 7 + 9 + 11 + 13 + 15 + \cdots + 77.$$

**Solution:** We have seen that this is an arithmetic progression with  $n = 37$  terms whose first term is 5 whose last term is 77, so  $S_{37} = \frac{1}{2}37(5 + 77) = 1517$ .

**Example**

Find the sum of the first  $n$  natural numbers:

$$1 + 2 + 3 + 4 + \cdots + (n - 1) + n$$

**Solution:** The terms in this sum form an arithmetic progression with common difference 1. The first term is 1 and the last term is  $n$ . Therefore the sum of the first  $n$  terms will be  $S_n = \frac{1}{2}n(n + 1)$ .

**Example**

The 8<sup>th</sup> term of an arithmetic progression is  $-\frac{1}{2}$ , and the 13<sup>th</sup> term is  $-8$ . Find the first term, the common difference, and the sum of the first 15 terms.

**Solution:** We can solve for  $a$  and  $d$ :

$$\begin{array}{rclclcl} 13^{\text{th}} \text{ term} & = & a_{13} & = & a + (13 - 1)d & = & a + 12d & = & -8 \\ 8^{\text{th}} \text{ term} & = & a_8 & = & a + (8 - 1)d & = & a + 7d & = & -\frac{1}{2} \\ \hline & & & & & & 5d & = & -\frac{15}{2}. \end{array}$$

Dividing the last equation through by 5 gives  $d = -\frac{3}{2}$ .

Substituting this value back into the first equation and rearranging gives  $a = 10$ .

The sum of the first 15 terms is  $S_{15} = \frac{15}{2}(2a + 14d) = \frac{15}{2}\left(20 + 14\left(-\frac{3}{2}\right)\right) = -\frac{15}{2}$ .

## 3.2 Geometric progressions

Consider a sequence

$$a_1, a_2, a_3, a_4, \dots, a_k, a_{k+1}, \dots,$$

(finite or infinite), such that all the terms are non-zero ( $a_k \neq 0$  for all  $k$ ). Then, instead of considering the differences  $d_k = a_{k+1} - a_k$ , as we did in the previous section, let's consider the *ratios*:

$$r_1 = a_2/a_1, \quad r_2 = a_3/a_2, \quad r_3 = a_4/a_3, \quad \dots, \quad r_k = a_{k+1}/a_k, \quad \dots$$

(Note that we need all of the terms in our original sequence to be non-zero so that we can divide by these numbers.)

If all the ratios are 1, then  $a_1 = a_2 = a_3 = \dots$ , so we again arrive at a constant sequence. The next most interesting case is when the ratios  $r_k$  are all the same. There is a special name for sequences of this type.

**Definition** (Geometric progression)

A **geometric progression** is a sequence of non-zero terms  $a_1, a_2, a_3, \dots$  in which the corresponding ratios  $r_k = a_{k+1}/a_k$  are all the same.

We call this value the **common ratio**  $r$ .

A geometric progression with common ratio  $r$  and first term  $a$  has the form

$$a, \quad ar, \quad ar^2, \quad ar^3, \quad \dots \quad ar^{k-1}, \quad \dots$$

Each term is obtained by multiplying the previous term by  $r$ .

The  $k$ th term of this sequence is  $a_k = ar^{k-1}$ .

(Notice that this definition is similar to that of an arithmetic progression considered before. Here we measure the “growth” of a sequence in a multiplicative way, rather than additively.)

**Example**

$$3, 6, 12, 24, 48, 96, \dots$$

is a geometric progression with first term  $a = 3$  and common ratio  $r = 2$ .

**Example**

$$1, \frac{1}{2}, \frac{1}{4}, \frac{1}{8}, \frac{1}{16}, \dots$$

is a geometric progression with first term  $a = 1$  and common ratio  $r = \frac{1}{2}$ .

**Theorem 18**

The sum of the first  $n$  terms of a geometric progression is

$$S_n = \begin{cases} \frac{a(1-r^n)}{1-r} & \text{if } r \neq 1, \\ na & \text{if } r = 1. \end{cases}$$

**Proof**

$$\begin{aligned} S_n &= a + ar + ar^2 + \dots + ar^{n-2} + ar^{n-1}. \\ rS_n &= ar + ar^2 + ar^3 + \dots + ar^{n-1} + ar^n. \end{aligned}$$

---

$$(1-r)S_n = a - ar^n = a(1-r^n).$$

If  $r \neq 1$  then we can divide both sides through by  $1-r$  to obtain the result. If  $r = 1$  then all terms of this geometric progression are equal, and the sum of the first  $n$  terms is  $n$  times the first term  $a$ . □

**Example**

In the following geometric progression

$$3, 6, 12, 24, 48, 96, \dots$$

after how many terms will the **sum** be greater than 9999 ?

**Solution:** We have  $a = 3$  and  $r = 2$ . The formula for the sum of the first  $n$  terms is

$$S_n = \frac{3 \times (1 - 2^n)}{1 - 2} = 3(2^n - 1).$$

We want to find the smallest number  $n$  for which  $3(2^n - 1) > 9999$ .

Dividing both sides through by 3 and adding 1 to both sides, we see that this is equivalent to finding the smallest value of  $n$  for which  $2^n > 3334$ .

We can use logarithms to solve this:

$$2^n > 3334 \Leftrightarrow \ln(2^n) > \ln(3334) \Leftrightarrow n \ln(2) > \ln(3334) \Leftrightarrow n > \frac{\ln(3334)}{\ln(2)}.$$

Using a calculator we find that

$$n > \frac{\ln(3334)}{\ln(2)} \approx \frac{8.112}{0.693} \approx 11.703.$$

Taking  $n = 12$  gives  $2^{12} = 4096 > 3334$ . Note that  $2^{11} = 2048 < 3334$ , so  $n = 12$  is indeed the smallest value of  $n$  for which  $2^n > 3334$ .

Thus after exactly 12 terms, the sum will be greater than 9999.

**Example**

Calculate the sum of the first six terms of a geometric progression with  $a = 1$  and  $r = 10$ .

**Solution:** Using the formula:

$$S_6 = \frac{1 \times (1 - 10^6)}{1 - 10} = \frac{-999999}{-9} = 111111.$$

Alternatively, we may notice that this sum is just

$$1 + 10 + 100 + 1000 + 10000 + 100000 = 111111.$$



**Example**

Calculate the sum of the first 8 terms of a geometric progression with  $a = 1$  and  $r = \frac{1}{2}$ .

**Solution:** Using the formula:

$$S_8 = \frac{1 \times \left(1 - \left(\frac{1}{2}\right)^8\right)}{1 - \frac{1}{2}} = \frac{\frac{2^8-1}{2^8}}{\frac{1}{2}} = \frac{2 \times (2^8 - 1)}{2^8} = \frac{510}{256} \approx 1.992 \text{ (3d.p.)}.$$

In fact, if we continue adding terms of the above progression, we get values closer and closer to 2. This gives an example of a *convergent series*, as we shall explain below.

**Definition** (Convergent and divergent series)

A **series** is an expression of the form  $a_1 + a_2 + \cdots + a_k + \cdots$ , representing the **sum** of terms in a sequence.

The notation

$$\sum_{k=1}^{\infty} a_k = a_1 + a_2 + \cdots + a_k + \dots$$

is also used.

We say that the series  $\sum_{k=1}^{\infty} a_k$  is **convergent** if the terms of the sequence of partial sums

$$a_1, \quad (a_1 + a_2), \quad (a_1 + a_2 + a_3), \quad \dots, \quad (a_1 + a_2 + \cdots + a_n), \dots$$

approach a fixed value as  $n \rightarrow \infty$ .

Otherwise we say that the sequence is **divergent**.

**Remark**

The notation

$$\sum_{k=1}^n a_k = a_1 + a_2 + \cdots + a_n$$

is also used for finite sums.

We give some examples of the compact notation for series and finite sums, using the examples we have seen thus far.

**Example**

The finite sequence 2, 5, 10, 17, 26, ...,  $k^2 + 1$ , ..., 10001 has 100 terms. Since the  $k$ th term of this sequence is  $a_k = k^2 + 1$ , we can use the notation

$$\sum_{k=1}^{100} (k^2 + 1)$$

to represent the sum of the terms in this sequence.

**Example**

The finite arithmetic progression 5, 7, 9, 11, 13, 15, ..., 77 has 37 terms. Since the  $k$ th term of this sequence is  $a_k = 5 + 2(k - 1)$ , we can use the notation

$$\sum_{k=1}^{37} (5 + 2(k - 1))$$

to represent the sum of the terms in this sequence.

A general arithmetic progression has the form  $a, a + d, a + 2d, \dots$ , with  $k$ th term  $a_k = a + (k - 1)d$ . We use the notation

$$\sum_{k=1}^n (a + (k - 1)d)$$

to represent the sum of the first  $n$  terms. The corresponding infinite series

$$\sum_{k=1}^{\infty} (a + (k - 1)d)$$

is **divergent** unless  $a = d = 0$ .

**Example**

Consider the infinite arithmetic progression

$$1, 2, 3, 4, 5, 6, \dots$$

with first term 1 and common difference 1. The corresponding sequence of partial sums is:

$$1, 3, 6, 10, 15, 21, \dots, \frac{1}{2}n(n + 1), \dots$$

These values do not approach a fixed value as  $n \rightarrow \infty$ . Thus the series  $\sum_{k=1}^{\infty} k$  is divergent.

**Example**

The infinite geometric progression with first term 1 and common difference 10 has  $k$ th term  $a_k = 10^{(k-1)}$ . We use the notation

$$\sum_{k=1}^n 10^{k-1} = \frac{1 - 10^n}{1 - 10}$$

to represent the sum of the first  $n$  terms in this sequence. The sequence of partial sums is

$$1, 11, 111, 1111, 11111, 111111, \dots, \frac{1 - 10^n}{1 - 10}, \dots$$

Since the terms do not approach a fixed value for  $n \rightarrow \infty$ , the infinite series

$$\sum_{k=1}^{\infty} 10^{k-1}$$

is **divergent**

Recall that a general geometric progression has the form  $a, ar, ar^2, \dots$ , with  $k$ th term  $a_k = ar^{k-1}$ . We use the notation  $\sum_{k=1}^n ar^{k-1}$  to represent the sum of the first  $n$  terms. The corresponding infinite series is  $\sum_{k=1}^{\infty} ar^{k-1}$ .

**Theorem 19**

The sum of an infinite geometric progression with first term  $a$  and common ratio  $r$  satisfying  $|r| < 1$  is

$$\sum_{k=1}^{\infty} ar^{k-1} = \frac{a}{1 - r}.$$

**Proof**

We have  $S_n = \frac{a(1 - r^n)}{1 - r}$ , and we note that as  $n$  gets larger, the value of  $r^n$  becomes smaller, since  $|r| < 1$ . We can say that as  $n \rightarrow \infty$ ,  $r^n \rightarrow 0$ . Thus, letting  $n \rightarrow \infty$  in the formula for  $S_n$  gives

$$\sum_{k=1}^{\infty} ar^{k-1} = \lim_{n \rightarrow \infty} S_n = \frac{a(1 - 0)}{1 - r} = \frac{a}{1 - r}.$$

□

**Example**

Using the previous result we find that the sum of the infinite geometric series with first term  $a = 1$  and common ratio  $r = \frac{1}{2}$  is

$$\sum_{k=1}^{\infty} \left(\frac{1}{2}\right)^{k-1} = 1 + \frac{1}{2} + \left(\frac{1}{2}\right)^2 + \left(\frac{1}{2}\right)^3 + \left(\frac{1}{2}\right)^4 + \dots = \frac{a}{1-r} = \frac{1}{0.5} = 2.$$

### 3.3 The binomial theorem

The binomial theorem is used when we want to multiply out expressions of the form  $(a + b)^n$ . We shall limit ourselves to the case where  $n$  is a positive integer, but there are analogues for negative, and non-integer values of  $n$ .

We can obtain the following directly (by multiplying out the brackets):

$$\begin{aligned} (a + b)^1 &= a + b \\ (a + b)^2 &= a^2 + 2ab + b^2 \\ (a + b)^3 &= a^3 + 3a^2b + 3ab^2 + b^3 \\ (a + b)^4 &= a^4 + 4a^3b + 6a^2b^2 + 4ab^3 + b^4 \\ (a + b)^5 &= a^5 + 5a^4b + 10a^3b^2 + 10a^2b^3 + 5ab^4 + b^5 \\ &\dots \end{aligned}$$

Each term of the above expansions is of the form  $a^{n-r}b^r$ , where  $r$  takes the values  $0, 1, \dots, n$  successively. Notice that the sum of the two exponents is  $(n - r) + r = n$ . The coefficients of the terms  $a^{n-r}b^r$  occurring in the expansion of  $(a + b)^n$  are called **binomial coefficients**. These coefficients may be obtained using Pascal's Triangle:

Pascal's Triangle																	
Row 0										1							
Row 1									1	1							
Row 2									1	2	1						
Row 3									1	3	3	1					
Row 4									1	4	6	4	1				
Row 5									1	5	10	10	5	1			
Row 6									1	6	15	20	15	6	1		
Row 7									1	7	21	35	35	21	7	1	
Row 8									1	8	28	56	70	56	28	8	1
...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...	...

The rule for forming the numbers in Pascal's triangle is:  
**Each number is the sum of the two nearest neighbors in the previous row.**

Notice that the  $n$ th row of Pascal's triangle contains  $n + 1$  numbers. We label the entries of the  $n$ th row as follows:

$$\binom{n}{0}, \binom{n}{1}, \binom{n}{2}, \dots, \binom{n}{n-1}, \binom{n}{n}.$$

For example the sixth row of Pascal's triangle contains the numbers:  $\binom{6}{0} = 1, \binom{6}{1} = 6, \binom{6}{2} = 15, \binom{6}{3} = 20, \binom{6}{4} = 15, \binom{6}{5} = 6, \binom{6}{6} = 1.$

Notice that:

$$\binom{n}{0} = \binom{n}{n} = 1, \quad \text{and} \quad \binom{n}{r} = \binom{n-1}{r-1} + \binom{n-1}{r} \quad \text{for all } 0 < r < n.$$

The mathematical notation  $\binom{n}{r}$  is pronounced “ $n$  choose  $r$ ” because these numbers count the different possible ways of choosing  $r$  objects from a choice of  $n$ . You may come across other notations for these numbers (e.g.  $\binom{n}{r}$  or  $nCr$  or  ${}^nC_r$ ) on your calculator.

**Theorem 20** (The Binomial Theorem)

If  $n$  is a positive integer, and  $a$  and  $b$  are real numbers then

$$(a + b)^n = \sum_{r=0}^n \binom{n}{r} a^{n-r} b^r,$$

where the coefficients  $\binom{n}{r}$  are the numbers in Pascal's triangle defined above.

**Proof**

We have seen that this is true for some small values of  $n$ .

Since  $(a + b)^{n+1} = (a + b)^n(a + b)$ , we shall first assume that the formula holds for some value of  $n$  and by multiplying through by  $a + b$  deduce that the corresponding formula must also hold for  $n + 1$ .

If

$$(a + b)^n = \binom{n}{0} a^n + \binom{n}{1} a^{n-1}b + \dots + \binom{n}{r} a^{n-r}b^r + \dots + \binom{n}{n} b^n,$$

then multiplying both sides by  $(a + b)$  gives

$$(a + b)^{n+1} = (a + b) \left[ \binom{n}{0} a^n + \binom{n}{1} a^{n-1}b + \dots + \binom{n}{r} a^{n-r}b^r + \dots + \binom{n}{n} b^n \right].$$

The right hand side can be rewritten as

$$\begin{aligned} & \left( \binom{n}{0} a^{n+1} + \binom{n}{1} a^n b + \dots + \binom{n}{r} a^{n+1-r} b^r + \dots + \binom{n}{n} a b^n \right) \\ & + \left( \binom{n}{0} a^n b + \binom{n}{1} a^{n-1} b^2 + \dots + \binom{n}{r} a^{n-r} b^{r+1} + \dots + \binom{n}{n} b^{n+1} \right) \end{aligned}$$

Gathering together terms involving the same powers of  $a$  and  $b$  gives

$$\begin{aligned} & \binom{n}{0} a^{n+1} + \left[ \binom{n}{1} + \binom{n}{0} \right] a^n b + \left[ \binom{n}{2} + \binom{n}{1} \right] a^{n-1} b^2 + \dots \\ & + \left[ \binom{n}{r} + \binom{n}{r-1} \right] a^{n+1-r} b^r + \dots + \binom{n}{n} b^{n+1} \\ = & a^{n+1} + \binom{n+1}{1} a^n b + \binom{n+1}{2} a^{n-1} b^2 + \dots + \binom{n+1}{r} a^{n+1-r} b^r + \dots + b^{n+1}, \end{aligned}$$

which is the desired formula for  $n + 1$ . Therefore by starting from the known cases and increasing the power by one at each step, we can prove the formula for all  $n = 1, 2, 3, 4, \dots$  using the above method.  $\square$

The previous result confirms our observation that the binomial coefficients are the numbers occurring in Pascal's triangle. Using Pascal's triangle is the quickest way of finding the binomial coefficients for small values of  $n$ . However, it is useful to have a non-recursive formula for binomial coefficients more generally. We need one more definition.

**Definition** (Factorial)

The number denoted  $k!$ , pronounced  $k$  **factorial**, is the product of all integers from  $k$  down to 1, i.e.,

$$k! = k \times (k - 1) \times (k - 2) \times \dots \times 3 \times 2 \times 1.$$

**Convention:** We also define  $0! = 1$ .

**Example**

$$\begin{aligned} 1! &= 1, & 2! &= 2 \times 1 = 2, & 3! &= 3 \times 2 \times 1 = 6, & 4! &= 4 \times 3 \times 2 \times 1 = 24, \\ 5! &= 5 \times 4! = 120. \end{aligned}$$

**Theorem 21**

For every positive integer  $n$  and for all  $r = 0, 1, \dots, n$ ,

$$\binom{n}{r} = \frac{n!}{r!(n-r)!}.$$

**Proof**

We check that the RHS, i.e., the number  $\frac{n!}{r!(n-r)!}$  satisfies the relations for the numbers in Pascal's Triangle. First note that

$$\frac{n!}{0!(n-0)!} = 1 \text{ and } \frac{n!}{n!(n-n)!} = 1.$$

Now for all positive  $n$  and  $r = 1, 2, \dots, n - 1$ , we have

$$\begin{aligned} \frac{n!}{r!(n-r)!} + \frac{n!}{(r-1)!(n-r+1)!} &= \frac{n!(n-r+1)}{r!(n-r+1)!} + \frac{n!r}{r!(n-r+1)!} \\ &= \frac{(n+1)n!}{r!(n-r+1)!} \\ &= \frac{(n+1)!}{r!(n+1-r)!}. \end{aligned}$$

Therefore we have shown that these numbers are calculated by exactly the same rules as the numbers in Pascal's triangle, i.e.,  $\binom{n}{r} = \frac{n!}{r!(n-r)!}$ .  $\square$

**Corollary 22**

It follows from the symmetry of the previous result that

$$\binom{n}{r} = \binom{n}{n-r}.$$

**Example**

In the case  $n = 5$ , we can compute the coefficients:

$$\begin{array}{llll} \binom{5}{0} & = \frac{5!}{5!0!} & = \frac{120}{120 \times 1} & = 1 \\ \binom{5}{1} & = \frac{5!}{4!1!} & = \frac{120}{24 \times 1} & = 5 \\ \binom{5}{2} & = \frac{5!}{3!2!} & = \frac{120}{6 \times 2} & = 10 \\ \binom{5}{3} & = \frac{5!}{2!3!} & = \frac{120}{2 \times 6} & = 10 \\ \binom{5}{4} & = \frac{5!}{1!4!} & = \frac{120}{1 \times 24} & = 5 \\ \binom{5}{5} & = \frac{5!}{0!5!} & = \frac{120}{1 \times 120} & = 1 \end{array}$$

giving the fifth row of Pascal's triangle.

**Example**

$$\begin{aligned}
(a+b)^5 &= \binom{5}{0}a^5b^0 + \binom{5}{1}a^4b^1 + \binom{5}{2}a^3b^2 + \binom{5}{3}a^2b^3 + \binom{5}{4}a^1b^4 + \binom{5}{5}a^0b^5 \\
&= a^5 + 5a^4b + 10a^3b^2 + 10a^2b^3 + 5ab^4 + b^5.
\end{aligned}$$

When expressing the binomial coefficients using factorials, it can be convenient to get rid of common factors in the numerator and denominator of the fraction:

$$\binom{n}{r} = \frac{n!}{r!(n-r)!} = \frac{n(n-1)\cdots(n-r+1)(n-r)\cdots 3\cdot 2\cdot 1}{r!(n-r)\cdots 3\cdot 2\cdot 1} = \frac{n(n-1)\cdots(n-r+1)}{r!}.$$

Therefore the first few terms in the binomial expansion (for general  $n$ ) are as follows:

$$(a+b)^n = a^n + n a^{n-1}b + \frac{n(n-1)}{2} a^{n-2}b^2 + \frac{n(n-1)(n-2)}{6} a^{n-3}b^3 + \dots$$

We may substitute any expression for the values of  $a$  and  $b$ .

**Example**

To expand  $(1-2x)^5$ , we set  $a = 1$  and  $b = -2x$ , which gives:

$$\begin{aligned}
(a+b)^5 &= a^5 + 5a^4b + 10a^3b^2 + 10a^2b^3 + 5ab^4 + b^5 \\
(1-2x)^5 &= 1 + 5(-2x) + 10(-2x)^2 + 10(-2x)^3 + 5(-2x)^4 + (-2x)^5 \\
&= 1 - 10x + 40x^2 - 80x^3 + 80x^4 - 32x^5.
\end{aligned}$$

**Example**

Approximate  $(0.98)^5$  to 4 decimal places.

**Solution:** We can use the previous example, by substituting  $x = 0.01$

$$(0.98)^5 = (1-2x)^5 = 1 - 0.1 + 0.004 - 0.00008 + 0.0000008 - 0.0000000032$$

Notice that the last two terms can be ignored as they are clearly insignificant to 4 decimal places. Thus

$$\begin{aligned}
(0.98)^5 &\approx 1 - 0.1 + 0.004 - 0.00008 \\
&\approx 0.9039 \text{ ( to four decimal places).}
\end{aligned}$$



**Example**

Use the binomial theorem to expand out the brackets in the expression  $(x + 3)^8$

**Solution:**

$$\begin{aligned}
 (x + 3)^8 &= \sum_{r=0}^8 \binom{8}{r} x^{8-r} (3)^r \\
 &= \binom{8}{0} x^8 (3)^0 + \binom{8}{1} x^7 (3)^1 + \binom{8}{2} x^6 (3)^2 + \binom{8}{3} x^5 (3)^3 + \binom{8}{4} x^4 (3)^4 \\
 &\quad + \binom{8}{5} x^3 (3)^5 + \binom{8}{6} x^2 (3)^6 + \binom{8}{7} x^1 (3)^7 + \binom{8}{8} x^0 (3)^8 \\
 &= (1 \times 1)x^8 + (8 \times 3)x^7 + (28 \times 9)x^6 + (56 \times 27)x^5 + (70 \times 81)x^4 \\
 &\quad + (56 \times 243)x^3 + (28 \times 729)x^2 + (8 \times 2187)x + (1 \times 6561) \\
 &= x^8 + 24x^7 + 252x^6 + 1512x^5 + 5670x^4 + 13608x^3 + 20412x^2 \\
 &\quad + 17496x + 6561
 \end{aligned}$$

**Example**

Find the coefficient of  $x^2$  in the expansion of  $(3x - \frac{1}{x})^8$ .

**Solution:** Each term of the expansion will be of the form  $\binom{n}{r} (3x)^r (\frac{-1}{x})^{8-r}$ . Note that we can write this term as some coefficient times  $x^{r-(8-r)}$ . Since we want to find the coefficient of the  $x^2$  term we must look for the value of  $r$  such that  $r - (8 - r) = 2r - 8 = 2$ ; it is easy to see that this is when  $r = 5$ . The corresponding binomial coefficient is

$$\binom{8}{5} = \frac{8!}{3!5!} = \frac{8 \cdot 7 \cdot 6}{3!} = 8 \cdot 7 = 56,$$

and so we deduce that the  $x^2$  term of the expansion will be

$$\binom{8}{5} (3x)^5 \left(\frac{-1}{x}\right)^3 = -56 \times 3^5 x^2 = -13608x^2.$$

Thus the required coefficient is  $-13608$ .

# Chapter 4

## Further calculus

4.1 Differentiation revisited

4.2 Implicit differentiation

4.3 Integration revisited

4.4 Integration by parts

4.5 Taylor series

### Revision from last semester..


The **derivative** of a function  $f(x)$  is given by

$$f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}, \text{ where } h \text{ represents a small change in } x.$$

Function	Derivative	Notes
$m$	0	$m$ any constant
$x^n$	$nx^{n-1}$	$n$ any (non-zero) real number
$e^x$	$e^x$	
$\ln(x)$	$\frac{1}{x}$	
$\sin(x)$	$\cos(x)$	for $x$ in radians
$\cos(x)$	$-\sin(x)$	for $x$ in radians
$\tan(x)$	$\sec^2(x)$	for $x$ in radians
$\lambda f(x)$	$\lambda f'(x)$	(constant $\times$ function)
$f(x) + g(x)$	$f'(x) + g'(x)$	(sum of functions)
$f(u(x))$	$f'(u(x))u'(x)$	(function of a function; 'chain rule' )
$u(x)v(x)$	$u'(x)v(x) + u(x)v'(x)$	(product of functions; 'product rule')
$\frac{u(x)}{v(x)}$	$\frac{u'(x)v(x) - u(x)v'(x)}{v(x)^2}$	(quotient of functions; 'quotient rule')

## 4.1 Differentiation revisited

The derivative  $f'(x)$  of a (continuous) function  $f(x)$  represents the gradient of the *tangent line* to  $f$  at a point  $x$ . We can make an approximation using a chord over a small change in  $x$ , denoted  $h$ :




The gradient of the chord (being a straight line with known end points  $P$  and  $Q$ ) is:

$$\text{gradient of chord} = \frac{f(x+h) - f(x)}{(x+h) - x} = \frac{f(x+h) - f(x)}{h}.$$

The gradient of tangent,  $f'(x)$ , is approximately equal to the gradient of the chord:

$$f'(x) \approx \frac{f(x+h) - f(x)}{h}.$$

**Key idea:** The smaller the change in  $x$ , the better the approximation.



The error  $\alpha(x, h)$  in our approximation depends upon both  $x$  and  $h$ :

$$\alpha(x, h) = \frac{f(x+h) - f(x)}{h} - f'(x)$$

**Definition** (Derivative)

A function  $f(x)$  is called **differentiable** if there are functions  $f'(x)$  and  $\alpha(x, h)$  such that

$$\alpha(x, h) = \frac{f(x+h) - f(x)}{h} - f'(x) \quad (4.1)$$

where  $\alpha(x, h) \rightarrow 0$  when  $h \rightarrow 0$ .

In this case

- $f'(x) = \lim_{h \rightarrow 0} \frac{f(x+h) - f(x)}{h}$  is called the **derivative** of  $f(x)$  at  $x$ .
- For small changes in  $x$ , the corresponding change in  $f$  can be approximated by  $f'(x)h$ . This linear function in the change in  $x$  is called the **differential** of  $f(x)$  at  $x$ .

The following notation is commonly used for the derivative of  $y = f(x)$ :

$$y'(x) = f'(x) = \frac{dy}{dx}.$$

We shall also use the following notation for the differential of  $y = f(x)$ :

$$dy = df = f'(x)dx$$

Differentials therefore have similar properties to derivatives:


Function	Derivative	Differential
$m$ (constant)	0	0
$x^n$	$nx^{n-1}$	$nx^{n-1}dx$
$e^x$	$e^x$	$e^x dx$
$\ln(x)$	$\frac{1}{x}$	$\frac{1}{x}dx$
$\sin(x)$	$\cos(x)$	$\cos(x)dx$
$\cos(x)$	$-\sin(x)$	$-\sin(x)dx$
$\tan(x)$	$\sec^2(x)$	$\sec^2(x)dx$
$\lambda f(x)$	$\lambda \frac{df}{dx}$	$\lambda df$
$f(x) + g(x)$	$\frac{df}{dx} + \frac{dg}{dx}$	$df + dg$
$f(u(x))$	$\frac{df}{du} \frac{du}{dx}$	$\frac{df}{du} du$
$u(x)v(x)$	$\frac{du}{dx}v + u \frac{dv}{dx}$	$duv + u dv$
$\frac{u(x)}{v(x)}$	$\frac{\frac{du}{dx}v - u \frac{dv}{dx}}{v^2}$	$\frac{duv - u dv}{v^2}$

The process of finding the differential (or equivalently the derivative) of a function is called **differentiation**.

In general, we shall only be interested in differentiable functions (i.e., those functions for which the differential and derivative make sense everywhere). We give two examples of non-differentiable functions. Geometrically this means that either there is no tangent for the graph  $y = f(x)$  or that the slope of the tangent is not defined, which happens if the tangent is vertical.

**Example**


The modulus function  $y = |x|$  is not differentiable at  $x = 0$ :



At  $x = 0$  there is no well-defined tangent to the graph. The graph is not “smooth” at this point. If we approach zero from the left, the tangent coincides with  $y = -x$ ; if we approach zero from the right, the tangent coincides with  $y = x$ .  
*At  $x = 0$  there is no tangent.*

**Example**

The positive square root function  $y = \sqrt{x}$  (defined for  $x \geq 0$ ) is not differentiable at  $x = 0$ :



The tangent line at  $x = 0$  is vertical, which means that the derivative tends to infinity as we approach zero from the right.  
*At  $x = 0$  the slope of the tangent is not defined.*

We show how to find the derivative of some well-known functions using the definition alone. This is called **differentiation from first principles**.

**Example**

Let  $f(x) = x^2$ . Then we have

$$\begin{aligned} f'(x) &= \lim_{h \rightarrow 0} \left( \frac{f(x+h) - f(x)}{h} \right) = \lim_{h \rightarrow 0} \left( \frac{(x+h)^2 - x^2}{h} \right) \\ &= \lim_{h \rightarrow 0} \left( \frac{x^2 + 2xh + h^2 - x^2}{h} \right) = \lim_{h \rightarrow 0} \left( \frac{2xh + h^2}{h} \right) = \lim_{h \rightarrow 0} (2x + h) = 2x \end{aligned}$$

We conclude that the derivative is  $f'(x) = 2x$  (as expected!).

The differential is  $df = 2x dx$ .

The previous example generalizes to higher powers of  $x$ :

**Example**

Let  $f(x) = x^n$ , where  $n$  is a positive integer. Then

$$f'(x) = \lim_{h \rightarrow 0} \left( \frac{f(x+h) - f(x)}{h} \right) = \lim_{h \rightarrow 0} \left( \frac{(x+h)^n - x^n}{h} \right)$$

Expanding  $(x+h)^n$  using the **Binomial theorem** and canceling out the  $x^n$  terms gives:

$$\begin{aligned} \frac{dy}{dx} &= \lim_{h \rightarrow 0} \left( \frac{nx^{n-1}h + \frac{1}{2}n(n-1)x^{n-2}(h)^2 + \cdots + (h)^n}{h} \right) \\ &= \lim_{h \rightarrow 0} \left( nx^{n-1} + \frac{1}{2}n(n-1)x^{n-2}h + \cdots + (h)^{n-1} \right) \end{aligned}$$

As  $h \rightarrow 0$  all but the first term tend to zero, so

$$f'(x) = nx^{n-1}.$$

The differential is  $df = nx^{n-1} dx$ .

Differentiation from first principles can be time-consuming! It also requires a pretty good understanding of limits. Luckily tables of derivatives of many well-known functions are widely available - we have seen some of these already. However, having a bit of knowledge about where the idea of derivative really comes from is useful.

Using our table of standard derivatives together with the fundamental properties of differentiation (such as the product rule and the chain rule) it is possible to differentiate a wide range of functions.

**Example**

Find the differential of  $y = x^2 \sin(x)$ .

**Solution:** Let  $u = x^2$  and  $v = \sin(x)$ , so that  $du = 2x dx$  and  $dv = \cos(x) dx$ , giving

$$\begin{aligned} dy = d(uv) &= du v + u dv \\ &= (2x dx) \sin(x) + x^2 (\cos(x) dx) \\ &= (2x \sin(x) + x^2 \cos(x)) dx. \end{aligned}$$

**Example**

Find the derivative of  $y = x \ln(x)$ .

**Solution:** Let  $u = x$  and  $v = \ln(x)$ , so that  $u' = 1$  and  $v' = \frac{1}{x}$  giving

$$y' = (uv)' = u'v + uv' = 1 \ln(x) + x \frac{1}{x} = \ln(x) + 1.$$

**Example**

Differentiate  $y = \sin(x^2)$ .

**Solution:** Let  $u(x) = x^2$  so that  $y(u) = \sin(u)$  and

$$\frac{dy}{du} = \cos(u), \quad du = 2x dx.$$

Using the chain rule gives

$$dy = \frac{dy}{du} du = (\cos(u)) (2x dx) = 2x \cos(x^2) dx.$$

**Example**

Find the derivative of  $y = e^{\sin(x)}$ .

**Solution:** Let  $u = \sin(x)$ , so that  $y = e^u$ . Applying the chain rule gives

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = e^u \cos(x) = e^{\sin(x)} \cos(x).$$

**Example**

Differentiate  $y = \frac{e^{x^2}}{\sin(x)}$ .

**Solution:** Let  $u = e^{x^2}$  and  $v = \sin(x)$ .

$$\begin{aligned} \frac{dy}{dx} &= \frac{v \frac{du}{dx} - u \frac{dv}{dx}}{v^2} = \frac{(\sin(x))(2xe^{x^2}) - (e^{x^2})(\cos(x))}{\sin^2(x)} \\ &= (2x\sin(x) - \cos(x)) \frac{e^{x^2}}{\sin^2(x)} \end{aligned}$$

**Example**

Find the differential of  $y = \cot(x) = \frac{\cos(x)}{\sin(x)}$ .

**Solution:**

$$\begin{aligned} dy &= \frac{d(\cos(x)) \sin(x) - \cos(x) d(\sin(x))}{\sin^2(x)} && \text{(By the quotient rule)} \\ &= \frac{(-\sin(x) dx) \sin(x) - \cos(x) (\cos(x) dx)}{\sin^2(x)} && \text{(Using tables.)} \\ &= \frac{(-\sin^2(x) - \cos^2(x)) dx}{\sin^2(x)} = \frac{-dx}{\sin^2(x)} && \text{(Using } \cos^2(x) + \sin^2(x) = 1) \\ &= -\operatorname{cosec}^2(x) dx. \end{aligned}$$

Since the quotient rule is a consequence of the product rule and the chain rule, if we prefer, we can always apply the product rule and then chain rule separately:

**Example**

Let  $y = \cot(x)$ . We have:

$$\begin{aligned} dy &= d\left(\cos(x) \frac{1}{\sin(x)}\right) = d(\cos(x)) \frac{1}{\sin(x)} + \cos(x) d\left(\frac{1}{\sin(x)}\right) && \text{(Product rule)} \\ &= (-\sin(x) dx) \frac{1}{\sin(x)} + \cos(x) \left(-\frac{1}{\sin^2(x)} (\cos(x) dx)\right) && \text{(Tables + Chain rule)} \\ &= \left(-1 - \frac{\cos^2(x)}{\sin^2(x)}\right) dx \\ &= -(1 + \cot^2(x)) dx = -\operatorname{cosec}^2(x) dx && \text{(Trig. identities)} \end{aligned}$$



### Example

Let  $y = \frac{1}{w}$  for any function  $w$  of  $x$ . Now, applying the chain rule:

$$\frac{dy}{dx} = \frac{dy}{dw} \frac{dw}{dx} = -w^{-2} \frac{dw}{dx} = -\frac{1}{w^2} \frac{dw}{dx}$$

### Example

Differentiate  $y = (1 + \cos(x^2))^6$ .

**Solution:** Let  $u = 1 + \cos(x^2)$ . Then  $y = u^6$  and by the chain rule

$$\frac{dy}{dx} = \frac{dy}{du} \frac{du}{dx} = 6u^5 \frac{du}{dx}.$$

Notice that  $u = 1 + \cos(x^2)$  is also a “function of a function”. So in order to differentiate  $u$  with respect to  $x$  we may use the chain rule *again*.

Let  $v = x^2$ . Then  $u = 1 + \cos(v)$  and by the chain rule:

$$\frac{du}{dx} = \frac{du}{dv} \frac{dv}{dx} = (-\sin(v)) 2x.$$

Putting this all together gives:

$$\begin{aligned} \frac{dy}{dx} &= \frac{dy}{du} \frac{du}{dv} \frac{dv}{dx} \\ &= 6u^5 (-\sin(v)) 2x \\ &= 6(1 + \cos(x^2))^5 (-\sin(x^2))2x \\ &= -12x \sin(x^2)(1 + \cos(x^2))^5 \end{aligned}$$

## 4.2 Implicit differentiation

An expression of the form  $y = f(x)$  describes  $y$  as an *explicit* function of  $x$ .

e.g.,  $y = x^2$ ,  $y = \cos(x) + \sin(x)$ ,  $y = \frac{1}{x+2}$  etc.

The dependence of  $y$  upon the variable  $x$  is given *explicitly* by an equation with  $y$  on one side, and an expression *in terms of the variable  $x$  only* on the other side.

We may encounter more complicated relationships between the variables  $x$  and  $y$ .

e.g.  $x^2 + y^2 = 1$ ,  $e^y = x^2 + 3$ ,  $\ln(2y + 1) = \sin(x)$  etc.

Such an equation still describes a dependence of variable  $y$  upon variable  $x$ , and so defines  $y$

as a function of  $x$  'implicitly'. We obtain an explicit function if we can solve the equation for  $y$ . However it is possible to find the derivative  $\frac{dy}{dx}$  **without** explicitly solving the equation.

**Example**

Find  $\frac{dy}{dx}$  when  $x^2 + y^2 = 1$ .

**Solution:** By treating  $y$  as an implicit function of  $x$ , we can differentiate the equation of the circle directly to obtain:

$$2x + 2y \frac{dy}{dx} = 0.$$

Rearranging this gives:

$$\frac{dy}{dx} = -\frac{2x}{2y} = -\frac{x}{y}.$$

Note that the right-hand side contains  $y$ ; this may be eliminated from the formula by solving the equation for  $y$ , if required.

For example, when  $y \geq 0$  we obtain

$$\frac{dy}{dx} = -\frac{x}{\sqrt{1-x^2}}.$$

**Example**

Find  $\frac{dy}{dx}$  when  $e^y = x^2 + 3$

**Solution:** By treating  $y$  as an implicit function of  $x$ , we obtain:

$$e^y \frac{dy}{dx} = 2x,$$

which rearranges to give

$$\frac{dy}{dx} = \frac{2x}{e^y} = \frac{2x}{x^2 + 3}$$

**Example**

Find  $\frac{dy}{dx}$  when  $\ln(2y - 1) = \sin(x)$

**Solution:** By treating  $y$  as an implicit function of  $x$ , we obtain:

$$\frac{1}{2y - 1} 2 \frac{dy}{dx} = \cos(x),$$

which rearranges to give

$$\frac{dy}{dx} = \frac{1}{2}(2y - 1) \cos(x) = \frac{1}{2} e^{\sin(x)} \cos(x)$$

Implicit differentiation is useful in finding the derivatives of inverse functions.

**Example**

Use implicit differentiation to find the derivative of  $y = \sin^{-1}(x)$ .

**Solution:** We have  $\sin(y) = x$ . Now treating  $y$  as an implicit function of  $x$  gives

$$\begin{aligned}\cos(y) \frac{dy}{dx} &= 1 \\ \frac{dy}{dx} &= \frac{1}{\cos(y)} = \frac{1}{\cos(\sin^{-1}(x))}\end{aligned}$$

This expression can be further simplified.

Using the identity  $\cos^2(y) + \sin^2(y) = 1$ , we note that  $\cos(y) = \pm\sqrt{1 - \sin^2(y)}$ .

Since  $-\frac{\pi}{2} \leq \sin^{-1}(x) \leq \frac{\pi}{2}$ , it follows that  $\cos(\sin^{-1}(x))$  is **positive**.

Thus  $\cos(\sin^{-1}(x)) = \sqrt{1 - \sin^2(\sin^{-1}(x))} = \sqrt{1 - x^2}$ , giving

$$\frac{dy}{dx} = \frac{1}{\sqrt{1 - x^2}}$$

**Example**

Use implicit differentiation to find the derivative of  $y = \cos^{-1}(x)$ .

**Solution:** We have  $\cos(y) = x$ . Now treating  $y$  as an implicit function of  $x$  gives

$$\begin{aligned}-\sin(y) \frac{dy}{dx} &= 1 \\ \frac{dy}{dx} &= -\frac{1}{\sin(y)} = -\frac{1}{\sin(\cos^{-1}(x))}\end{aligned}$$

This expression can be further simplified.

Using the identity  $\cos^2(y) + \sin^2(y) = 1$ , we note that  $\sin(y) = \pm\sqrt{1 - \cos^2(y)}$ .

Since  $0 \leq \cos^{-1}(x) \leq \pi$ , it follows that  $\sin(\cos^{-1}(x))$  is **positive**.

Thus  $\sin(\cos^{-1}(x)) = \sqrt{1 - \cos^2(\cos^{-1}(x))} = \sqrt{1 - x^2}$ , giving

$$\frac{dy}{dx} = -\frac{1}{\sqrt{1 - x^2}}$$

**Example**

Use implicit differentiation to find the derivative of  $y = \tan^{-1}(x)$ .

**Solution:** We have  $\tan(y) = x$ . Now treating  $y$  as an implicit function of  $x$  gives

$$\begin{aligned}\sec^2(y) \frac{dy}{dx} &= 1 \\ \frac{dy}{dx} &= \frac{1}{\sec^2(y)} = \cos^2(y) = \cos^2(\tan^{-1}(x))\end{aligned}$$

This expression can be simplified using the identity  $1 + \tan^2(y) = \sec^2(y)$  which rearranges to  $\cos^2(y) = \frac{1}{1 + \tan^2(y)}$ . Therefore

$$\frac{dy}{dx} = \frac{1}{1 + \tan^2(\tan^{-1}(x))} = \frac{1}{1 + x^2}.$$

**Example**

Find the derivative of  $y = a^x$  where  $a > 0$  and  $a \neq 1$ .

**Solution:** First note that we may write  $\ln(y) = \ln(a^x) = x \ln(a)$ . Now treating  $y$  as an implicit function of  $x$  gives

$$\begin{aligned}\frac{1}{y} \frac{dy}{dx} &= \ln(a) \\ \frac{dy}{dx} &= \ln(a)y = \ln(a)a^x.\end{aligned}$$

**Example**

Find the differential of  $y = x^\alpha$  where  $x > 0$  and  $\alpha$  is a real number.

**Solution:** First note that we may write  $\ln(y) = \alpha \ln(x)$ . Now treating  $y$  as an implicit function of  $x$  gives

$$\begin{aligned}\frac{1}{y} \frac{dy}{dx} &= \alpha \frac{1}{x} \\ \frac{dy}{dx} &= \alpha \frac{y}{x} = \alpha \frac{x^\alpha}{x} = \alpha x^{\alpha-1}.\end{aligned}$$

Notice that the formula is the same as that for the powers  $x^n$ , where  $n$  is an integer.

We add the more fundamental examples to our table of standard derivatives:

Function	Derivative	Differential
$\cot(x)$	$-\operatorname{cosec}^2(x)$	$-\operatorname{cosec}^2(x) dx$
$\sin^{-1}(x)$	$\frac{1}{\sqrt{1-x^2}}$	$\frac{dx}{\sqrt{1-x^2}}$
$\cos^{-1}(x)$	$\frac{-1}{\sqrt{1-x^2}}$	$\frac{-dx}{\sqrt{1-x^2}}$
$\tan^{-1}(x)$	$\frac{1}{1+x^2}$	$\frac{dx}{1+x^2}$
$x^\alpha$ (where $x > 0$ and $\alpha$ is a real number)	$\alpha x^{\alpha-1}$	$\alpha x^{\alpha-1} dx$
$a^x$ (where $a > 0$ )	$a^x \ln(a)$	$a^x \ln(a) dx$

### 4.3 Integration revisited

Recall the relationship between integration and differentiation that we established at the beginning of the course. For  $F'(x) = f(x)$ , we defined

**Indefinite integrals:**  $\int f(x) dx = F(x) + C$  where  $C$  is a constant.

**Definite integrals:**  $\int_a^b f(x) dx = F(b) - F(a)$ .

Since the derivative and differential of a function  $F(x)$  are related by

$$dF = F'(x) dx,$$

the above formulas can be conveniently expressed in terms of differentials as

$$\int dF = F(x) + C \quad \text{and} \quad \int_a^b dF = F(b) - F(a)$$

In this way the integral can easily be seen as the ‘inverse operator’ to the differential. By combining our knowledge of derivatives with other techniques we’ve learned so far, we can now integrate many functions.

For example, we can use the table of differentials found in the previous chapter to add to our table of standard indefinite integrals:

Function	Indefinite integral
$m$ (a constant)	$\int m dx = mx + C$
$nx^{n-1}$	$\int nx^{n-1} dx = x^n + C$
$e^x$	$\int e^x dx = e^x + C$
$\frac{1}{x}$	$\int \frac{1}{x} dx = \ln x  + C$
$\sin(x)$	$\int \sin(x) dx = -\cos(x) + C$
$\cos(x)$	$\int \cos(x) dx = \sin(x) + C$
$\sec^2(x)$	$\int \sec^2(x) dx = \tan(x) + C$
$\operatorname{cosec}^2(x)$	$\int \frac{dx}{\sin^2(x)} = -\cot(x) + C$
$\frac{1}{\sqrt{1-x^2}}$	$\int \frac{dx}{\sqrt{1-x^2}} = \sin^{-1}x + C$
$\frac{1}{1+x^2}$	$\int \frac{dx}{1+x^2} = \tan^{-1}x + C$
$x^\alpha$	$\int x^\alpha dx = \frac{x^{\alpha+1}}{\alpha+1} + C \quad (\text{for } \alpha \neq -1)$
$a^x$	$\int a^x dx = \frac{a^x}{\ln a} + C \quad (\text{for } a \neq 1)$

**Example**

Use the substitution  $x = 2\sin(v)$  to calculate:

$$\int_0^2 \sqrt{4-x^2} \, dx.$$

**Solution:** Let  $x = 2\sin(v)$ , then  $dx = 2\cos(v) \, dv$ . The new limits are as follows: If  $x = 0$ , then  $v = 0$ , and if  $x = 2$ , then  $v = \frac{\pi}{2}$ . Hence

$$\begin{aligned} \int_0^2 \sqrt{4-x^2} \, dx &= \int_0^{\frac{\pi}{2}} \sqrt{4-(2\sin(v))^2} \, 2\cos(v) \, dv \\ &= \int_0^{\frac{\pi}{2}} \sqrt{4(1-\sin^2(v))} \, 2\cos(v) \, dv \\ &= \int_0^{\frac{\pi}{2}} \sqrt{4\cos^2(v)} \, 2\cos(v) \, dv \end{aligned}$$

Since  $\cos(v)$  is **positive** when  $0 \leq v \leq \frac{\pi}{2}$ , we note that  $\sqrt{\cos^2(v)} = \cos(v)$ :

$$\begin{aligned} \int_0^2 \sqrt{4-x^2} \, dx &= \int_0^{\frac{\pi}{2}} (2\cos(v))(2\cos(v)) \, dv \\ &= \int_0^{\frac{\pi}{2}} 4\cos^2(v) \, dv \\ &= \int_0^{\frac{\pi}{2}} 2(1+\cos(2v)) \, dv \\ &= [2v + \sin(2v)]_0^{\frac{\pi}{2}} = (\pi + 0) - (0 + 0) = \pi. \end{aligned}$$

Integration by substitution can be thought of as the “reverse” process to differentiation using the chain rule.

## 4.4 Integration by parts

The product rule for differentiating the product of functions  $u = u(x)$  and  $v = v(x)$  is:

$$d(uv) = du \cdot v + u \cdot dv.$$

Integrating both sides, we obtain:

$$\int d(uv) = \int du \, v + \int u \, dv.$$

Since integrating the differential, gives back the original function (in this case  $uv$ ) we obtain:

$$uv = \int du \, v + \int u \, dv.$$

Rearranging, we arrive at the following **integration by parts** formula:

$$\int u \, dv = uv - \int v \, du.$$

This method is called ‘integration by parts’ because it requires us to re-write  $f(x) \, dx$  in the form  $u \, dv$ , which has two ‘parts’.

- The first part  $u(x)$  must be a factor of  $f(x)dx$  which we feel able to differentiate (since to apply the formula we have to first find  $du$ ).
- The second part  $dv$  will be what remains of  $f(x)dx$  once we have factored out  $u(x)$ . Note that to apply the formula we must be able to integrate  $dv$ .

There is the corresponding integration by parts formula for definite integrals:

$$\int_a^b u \, dv = [uv]_a^b - \int_a^b v \, du.$$

Here the limits refer to the independent variable  $x$ , where  $u = u(x)$  and  $v = v(x)$ .

Integration by parts reduces the original integral to another integral  $\int v \, du$ , which may or may not be easier to calculate than the original.

**The method is successful if this second integral is easier.**

### Example

Find  $\int x \sin(x) \, dx$ .

**Solution:** Let  $u = x$  and  $dv = \sin(x) \, dx$ . Differentiating  $u$ , gives  $du = dx$ . Integrating  $dv$  gives  $v = -\cos(x)$ . Now using the integration by parts formula:

$$\begin{aligned} \int x \sin(x) \, dx &= -x\cos(x) - \int (-\cos(x)) \, dx \\ &= -x\cos(x) + \sin(x) + C. \end{aligned}$$

(The second integral  $\int (-\cos(x)) \, dx$  is easier than the original integral so, in this case, the method works successfully.)

Notice that whilst we already know how to differentiate  $\ln(x)$ , we have not yet learned how to integrate this function. We can use integration by parts to do so:



**Example**

Find  $\int \ln(x) \, dx$ .

**Solution:** We take  $u = \ln(x)$  and  $dv = dx$  so that  $du = \frac{1}{x} dx$  and  $v = x$ . Applying the integration by parts formula gives

$$\int \ln(x) \, dx = x \ln(x) - \int x \frac{dx}{x} = x \ln(x) - \int dx = x \ln(x) - x + C.$$

**Example**

Find  $\int \sqrt{x} \ln x \, dx$ .

**Solution:** Setting  $u = \ln(x)$  gives  $du = \frac{dx}{x}$ . Then  $dv = \sqrt{x} \, dx = x^{\frac{1}{2}} \, dx$ , giving  $v = \frac{2}{3} x^{\frac{3}{2}}$ . Now, integrating by parts

$$\begin{aligned} \int \sqrt{x} \ln(x) \, dx &= \frac{2}{3} x^{\frac{3}{2}} \ln(x) - \int \frac{2}{3} x^{\frac{3}{2}} \frac{dx}{x} = \frac{2}{3} x^{\frac{3}{2}} \ln(x) - \frac{2}{3} \int x^{\frac{1}{2}} \, dx \\ &= \frac{2}{3} x^{\frac{3}{2}} \ln(x) - \frac{4}{9} x^{\frac{3}{2}} + C \\ &= \frac{2}{3} x^{\frac{3}{2}} \left( \ln(x) - \frac{2}{3} \right) + C. \end{aligned}$$

It may be necessary to use integration by parts more than once, as in the following example.

**Example**

Find  $\int x^2 e^x \, dx$ .

**Solution:** Let  $u = x^2$  and  $dv = e^x \, dx$ . Then  $du = 2x \, dx$  and  $v = e^x$ . So

$$\int x^2 e^x \, dx = x^2 e^x - \int 2x e^x \, dx.$$

The second integral may be found by applying integration by parts *again*. Let  $u = 2x$  and let  $dv = e^x \, dx$  so that  $du = 2 \, dx$  and  $v = e^x$ . Then

$$\int 2x e^x \, dx = 2x e^x - \int 2 e^x \, dx = 2x e^x - 2e^x + C$$

and so

$$\int x^2 e^x \, dx = x^2 e^x - 2x e^x + 2e^x + C = (x^2 - 2x + 2) e^x + C.$$

## 4.5 Taylor series

Write  $f^{(r)}(x)$  to denote the function obtained from  $f(x)$  after differentiating  $r$  times.

### Example

Find  $f^{(3)}(x)$  for  $f(x) = x^2 + \sin(x)$ .

**Solution:**

$$f^{(1)}(x) = 2x + \cos(x), \quad f^{(2)}(x) = 2 - \sin(x), \quad f^{(3)}(x) = -\cos(x)$$

The main statement of this section is as follows.

### Theorem 23 (Taylor series about the point $x = a$ )

Let  $f(x)$  be a function which can be differentiated infinitely many times at  $x = a$ .

$$\begin{aligned} \text{For } x \rightarrow a, \quad f(x) &= \sum_{r=0}^{\infty} \frac{f^{(r)}(a)}{r!} (x-a)^r \\ &= f(a) + f^{(1)}(a)(x-a) + \frac{1}{2} f^{(2)}(a)(x-a)^2 + \frac{1}{3!} f^{(3)}(a)(x-a)^3 \\ &\quad + \dots + \frac{1}{n!} f^{(n)}(a)(x-a)^n + \dots \end{aligned}$$

Taylor series about the point  $x = 0$  (that is setting  $a = 0$  in the above), are sometimes called **Maclaurin series**.

**Example**

Let  $\alpha$  be a real number. Find the Taylor series of  $f(x) = (1+x)^\alpha$  at zero.

**Solution:** We need to evaluate the function and its derivatives at  $x = 0$ .

$$\begin{array}{l|l} f(x) = (1+x)^\alpha, & f(0) = 1 \\ f^{(1)}(x) = \alpha(1+x)^{\alpha-1}, & f^{(1)}(0) = \alpha \\ f^{(2)}(x) = \alpha(\alpha-1)(1+x)^{\alpha-2} & f^{(2)}(0) = \alpha(\alpha-1) \\ \vdots & \vdots \\ f^{(n)}(x) = \alpha(\alpha-1)\dots(\alpha-n+1)(1+x)^{\alpha-n} & f^{(n)}(0) = \alpha(\alpha-1)\dots(\alpha-n+1) \end{array}$$

Substituting these values into the Taylor formula we find that for  $x \rightarrow 0$

$$\begin{aligned} (1+x)^\alpha = & 1 + \alpha x + \frac{\alpha(\alpha-1)}{2} x^2 + \frac{\alpha(\alpha-1)(\alpha-2)}{3!} x^3 + \dots \\ & + \frac{\alpha(\alpha-1)\dots(\alpha-n+1)}{n!} x^n + \dots \end{aligned}$$

Note the similarity between the formula above and the expansion of  $(1+x)^n$  given by the **Binomial theorem**, where  $n$  is a positive integer.

**Example**

Find the Taylor series about  $x = 0$  of  $f(x) = (1+x)^{-1}$ .

**Solution:** Setting  $\alpha = -1$  in the previous example we see that the coefficient of  $x^n$  will be  $(-1)(-2)\dots(-1-n+1)/n! = (-1)(-2)\dots(-n)/n! = (-1)^n$ . Therefore for  $x \rightarrow 0$

$$(1+x)^{-1} = 1 - x + x^2 - x^3 + \dots + (-1)^n x^n + \dots = \sum_{k=0}^{\infty} (-x)^k$$

Notice that the terms on the right-hand side can be viewed as a geometric progression with first term 1 and common ratio  $-x$ . Using the formula for the sum of a geometric series (which holds for all  $|x| < 1$ ) we obtain the left hand side.

Notice that when  $x$  is very close to  $a$ , or in other words, when the difference  $x - a$  is very small, the powers  $(x - a)^2$ ,  $(x - a)^3$ , etc. get smaller and smaller and smaller. If we are interested in finding an *approximation* to the function  $f(x)$  in an interval around the point  $x = a$ , we can therefore do so by ignoring all terms in the expansion involving powers  $(x - a)^{n+1}$  and higher.

**Example**

If we ignore all terms involving powers  $(x - a)^2$  and higher we obtain

$$f(x) \approx f(a) + f'(a)(x - a) \quad \text{for } x \rightarrow a$$

(Writing  $x = a + h$ , we recover the definition of the derivative at  $a$ .)

In general by ignoring all terms in the expansion involving  $(x - a)^{n+1}$  and higher powers, the Taylor formula allows us to approximate a function by a polynomial of degree  $n$ .

**Definition** (Taylor expansion)

The  $n$ th order Taylor expansion of a function  $f(x)$  at  $a$  is written as

$$\text{For } x \rightarrow a, \quad f(x) \approx f(a) + f'(a)(x - a) + \frac{1}{2} f^{(2)}(a)(x - a)^2 + \cdots + \frac{1}{n!} f^{(n)}(a)(x - a)^n.$$

**Example**

Find the third order Taylor expansion of  $f(x) = \sqrt{1 + x}$  at zero.

**Solution:** This is another special case of  $(1 + x)^\alpha$ , where  $\alpha = \frac{1}{2}$ . Note that this time we were only asked to find the expansion up to and including the  $x^3$  term. Setting  $\alpha = \frac{1}{2}$  in the Taylor expansion of  $(1 + x)^\alpha$  we find that for  $x \rightarrow 0$ ,

$$\begin{aligned} \sqrt{1 + x} &\approx 1 + \frac{1}{2}x + \frac{1}{2!} \frac{1}{2} \left(\frac{1}{2} - 1\right) x^2 + \frac{1}{3!} \frac{1}{2} \left(\frac{1}{2} - 1\right) \left(\frac{1}{2} - 2\right) x^3 \\ &\approx 1 + \frac{1}{2}x - \frac{1}{8}x^2 + \frac{1}{16}x^3 \end{aligned}$$

**Example**

Find the Taylor series about  $x = 0$  for the function  $f(x) = e^x$ .

**Solution:** We have  $f^{(1)}(x) = e^x$ ,  $f^{(2)}(x) = e^x$ , etc.

Evaluating at  $x = 0$  gives,  $f(0) = f^{(1)}(0) = f^{(2)}(0) = \cdots = 1$ . Therefore for  $x \rightarrow 0$

$$e^x = 1 + x + \frac{1}{2!}x^2 + \frac{1}{3!}x^3 + \frac{1}{4!}x^4 + \cdots + \frac{1}{n!}x^n + \cdots = \sum_{k=0}^{\infty} \frac{1}{k!}x^k.$$

**Example**

Find the Taylor series about  $x = 0$  for the function  $f(x) = \sin(x)$ .

**Solution:** We have

$f^{(1)}(x) = \cos(x)$ ,  $f^{(2)}(x) = -\sin(x)$ ,  $f^{(3)}(x) = -\cos(x)$  and  $f^{(4)}(x) = \sin(x) = f(x)$ . Evaluating at  $x = 0$  gives  $f(0) = 0$ ,  $f^{(1)}(0) = 1$ ,  $f^{(2)}(0) = 0$ ,  $f^{(3)}(0) = -1$ , etc. Notice that we get a value of 0 for all derivatives of even orders, whilst the values for the derivatives of odd orders alternate between +1 and -1. Therefore for  $x \rightarrow 0$

$$\sin(x) = x - \frac{1}{3!}x^3 + \frac{1}{5!}x^5 - \frac{1}{7!}x^7 + \dots + (-1)^n \frac{1}{(2n+1)!}x^{2n+1} + \dots$$

**Example**

Find the Taylor series about  $x = 0$  for the function  $f(x) = \cos(x)$ .

**Solution:** This is very similar to the previous example.

We have  $f^{(1)}(x) = -\sin(x)$ ,  $f^{(2)}(x) = -\cos(x)$ ,  $f^{(3)}(x) = \sin(x)$ , etc.

Evaluating at  $x = 0$  gives  $f(0) = 1$ ; we get a value of 0 for all derivatives of odd orders and values alternating between +1 and -1 for the derivatives of even orders. Therefore using the Taylor formula we find:

$$\cos(x) = 1 - \frac{1}{2!}x^2 + \frac{1}{4!}x^4 - \frac{1}{6!}x^6 + \dots + (-1)^n \frac{1}{(2n)!}x^{2n} + \dots$$

**Example**

Find the Taylor series about  $x = 0$  for the function  $f(x) = \ln(1+x)$ .

**Solution:**

$$f(x) = \ln(1+x)$$

$$f(0) = \ln(1) = 0$$

$$f^{(1)}(x) = \frac{1}{1+x} = (1+x)^{-1}$$

$$f^{(1)}(0) = 1$$

$$f^{(2)}(x) = (-1)(1+x)^{-2}$$

$$f^{(2)}(0) = -1 \times 1!$$

$$f^{(3)}(x) = (-1)(-2)(1+x)^{-3}$$

$$f^{(3)}(0) = (-1)^2 \times 2!$$

$\vdots$

$\vdots$

$$f^{(n)}(x) = (-1)(2) \dots (-(n-1))(1+x)^{-n} \quad f^{(n)}(0) = (-1)^{n-1} (n-1)!$$

and hence

$$\ln(1+x) = x - \frac{1}{2}x^2 + \frac{1}{3}x^3 - \frac{1}{4}x^4 + \dots + (-1)^{n-1} \frac{1}{n}x^n + \dots$$

Taylor formula has many applications. One such application is to finding limits.

**Example**

Calculate the limit:

$$\lim_{x \rightarrow 0} \frac{\sqrt{1+2x} - 1}{x}.$$

**Solution:** Notice that it is not possible to simply substitute  $x = 0$  into the top and bottom of the fraction, because we cannot evaluate “ $\frac{0}{0}$ ”.

Using the first order Taylor expansion  $\sqrt{1+2x}$  for  $x \rightarrow 0$  we have

$$\sqrt{1+2x} = 1 + \frac{1}{2}2x + \cdots = 1 + x + \text{terms involving higher powers of } x$$

and hence

$$\begin{aligned} \lim_{x \rightarrow 0} \left( \frac{\sqrt{1+2x} - 1}{x} \right) &= \lim_{x \rightarrow 0} \left( \frac{x + \text{terms involving higher powers of } x}{x} \right) \\ &= \lim_{x \rightarrow 0} \left( 1 + \text{terms involving powers of } x \right) = 1. \end{aligned}$$

# Chapter 5

## Rational functions

- 5.1 Polynomials and long division
- 5.2 Introducing partial fractions
- 5.3 The method of partial fractions
- 5.4 Integration by partial fractions

### Revision from last semester..

#### Fractions:

$$\frac{p}{q} + \frac{n}{m} = \frac{pm + nq}{qm}$$

$$\frac{p}{q} - \frac{n}{m} = \frac{pm - nq}{qm}$$

$$\frac{p}{q} \times \frac{n}{m} = \frac{pn}{qm}$$

$$\frac{p}{q} \div \frac{n}{m} = \frac{pm}{qn}$$

#### The quadratic formula:

The quadratic equation

$$ax^2 + bx + c = 0,$$

has solutions

$$x = \frac{-b \pm \sqrt{b^2 - 4ac}}{2a}$$

## 5.1 Polynomials and long division

### Definition (Polynomial)

A **polynomial**  $P(x)$  is a function of  $x$  that may be written in the form

$$P(x) = a_n x^n + a_{n-1} x^{n-1} + \cdots + a_1 x + a_0,$$

where  $n$  is a non-negative integer and  $a_n, a_{n-1}, \dots, a_1, a_0$  are given real numbers called **coefficients**.

The **degree** of the polynomial, denoted  $\deg(P)$ , is the exponent of the highest power of  $x$  occurring. If  $a_n \neq 0$  in the above expression, then  $\deg(P) = n$ .

We say that  $a_n x^n$  is the **leading term** of this polynomial and that  $a_0$  is the **constant term** of this polynomial.

### Example

We give a few examples of these definitions below:

Polynomial	Degree	Leading term	Constant term
$4x^3 + 3x^2 - 2$	3	$4x^3$	-2
$(4x + 1)(x + 3)$	2	$4x^2$	3
$17x^2 + \frac{1}{2}x^6$	6	$\frac{1}{2}x^6$	0
3	0	3	3

### Definition (Rational function)

A **rational function**  $R(x)$  is the ratio of two polynomials  $N(x)$ ,  $D(x)$

$$R(x) = \frac{N(x)}{D(x)}$$

$R(x)$  is called **proper** if  $\deg(N) < \deg(D)$  and **improper** if  $\deg(N) \geq \deg(D)$ .



**Example**

Rational function	Proper or Improper?
$\frac{x+1}{x^2+5x+6}$	Proper, since $\deg(x+1) < \deg(x^2+5x+6)$
$\frac{x^3+x^2+1}{(x-1)(x-1)}$	Improper, since $\deg(x^3+x^2+1) \geq \deg((x-1)(x-1))$
$\frac{x^5-1}{x^5+1}$	Improper, since $\deg(x^5-1) \geq \deg(x^5+1)$

**Theorem 24**

An improper rational function,  $R(x)$ , can always be expressed as the sum of a polynomial  $Q(x)$  and a proper rational function:

$$\frac{N(x)}{D(x)} = Q(x) + \frac{M(x)}{D(x)}$$

where  $\deg(M) < \deg(D)$  and  $\deg(Q) = \deg(N) - \deg(D)$

The above result tells us that there will be polynomials  $Q(x)$  and  $M(x)$  which satisfy this equation, but it does not tell us how to find them. (More on that later!)

**Exercise**

By multiplying out, check that

$$\frac{x^6 - 2x^4 + x^2 - 2}{x^2 - x - 2} = x^4 + x^3 + x^2 + 3x + 6 + \frac{12x + 10}{x^2 - x - 2}$$

**Example**

It is easy to see that

$$\frac{x}{x+1} = \frac{x+1-1}{x+1} = 1 - \frac{1}{x+1}$$

**Theorem 25** (A version of the previous one)

For every two polynomials,  $N(x)$  and  $D(x)$ , it is possible to write

$$N(x) = Q(x)D(x) + M(x)$$

where  $Q(x)$  and  $D(x)$  are polynomials and  $\deg(M) < \deg(D)$ .

This is called “division with remainder”. The polynomial  $Q(x)$  is called the **quotient** and the polynomial  $M(x)$  is called the **remainder**. If  $M(x) = 0$ , that is if there is no remainder, then  $N(x) = Q(x)D(x)$ ; in this case we say that  $D(x)$  is a factor of  $N(x)$ . If the degree of  $D(x)$  exceeds the degree of  $N(x)$ , then we may write  $N(x) = 0 \cdot D(x) + N(x)$ , so that in this case  $N(x)$  itself is the “remainder”.

An important special case occurs when the divisor  $D$  is linear,  $D(x) = x - a$ . Then

$$N(x) = Q(x)(x - a) + M$$

where  $M$  is a polynomial of degree less than 1, or in other words, a constant.

**Theorem 26** (The Remainder Theorem)

For any polynomial  $N(x)$ , the remainder after division by  $x - a$  is the value of  $N(x)$  at  $a$ :

$$N(x) = Q(x)(x - a) + N(a).$$

**Proof**

By the previous remark we know that  $N(x) = Q(x)(x - a) + M$  where  $M$  is a constant. Substituting  $x = a$  into this equation yields  $N(a) = Q(a) \cdot 0 + M = M$ .  $\square$

**Corollary 27**

The linear polynomial  $x - a$  is a factor of the polynomial  $N(x)$  if and only if  $N(a) = 0$ .

The practical process of finding the polynomials  $Q(x)$  and  $M(x)$  is by **polynomial long division**, which is similar to usual long division of numbers. This is best illustrated by means of some examples.

**Example**

Use polynomial long division to write the following improper rational function as the sum of a polynomial and a proper rational function:

$$\frac{N(x)}{D(x)} = \frac{x^4 + 2x^3 - 2x^2 - x + 4}{x^3 + 4x^2 + 5x + 2}.$$

**Solution:** We perform polynomial long division:

$$\begin{array}{r}
 \phantom{x^3 + 4x^2 + 5x + 2} \overline{x - 2} \phantom{=} = Q(x) \text{ quotient} \\
 x^3 + 4x^2 + 5x + 2 \phantom{)} \phantom{=} \\
 \underline{x^4 + 2x^3 - 2x^2 - x + 4} \phantom{=} \\
 \phantom{x^3 + 4x^2 + 5x + 2} \phantom{)} \phantom{=} x^4 + 4x^3 + 5x^2 + 2x \\
 \phantom{x^3 + 4x^2 + 5x + 2} \phantom{)} \phantom{=} \phantom{x^4 + 2x^3 - 2x^2 - x + 4} -2x^3 - 7x^2 - 3x + 4 \\
 \phantom{x^3 + 4x^2 + 5x + 2} \phantom{)} \phantom{=} \phantom{x^4 + 2x^3 - 2x^2 - x + 4} -2x^3 - 8x^2 - 10x - 4 \\
 \phantom{x^3 + 4x^2 + 5x + 2} \phantom{)} \phantom{=} \phantom{x^4 + 2x^3 - 2x^2 - x + 4} \phantom{-2x^3 - 7x^2 - 3x + 4} x^2 + 7x + 8 \phantom{=} = M(x) \text{ remainder}
 \end{array}$$

Hence

$$\frac{x^4 + 2x^3 - 2x^2 - x + 4}{x^3 + 4x^2 + 5x + 2} = x - 2 + \frac{x^2 + 7x + 8}{x^3 + 4x^2 + 5x + 2}.$$

Notice that  $\frac{x^2 + 7x + 8}{x^3 + 4x^2 + 5x + 2}$  is proper, as required.

**Example**

Show that the cubic polynomial

$$P(x) = x^3 - 3x^2 + 7x - 5$$

is divisible by  $x - 1$  and factorise it.

**Solution:** Substituting  $x = 1$  into  $P(x)$  gives  $P(1) = 1 - 3 + 7 - 5 = 0$ , therefore  $x - 1$  is a factor of  $P(x)$ . Now using polynomial long division we obtain

$$\begin{array}{r}
 x^2 - 2x + 5 \qquad \qquad \qquad = Q(x) \text{ quotient} \\
 \hline
 x-1 \ ) \ x^3 - 3x^2 + 7x - 5 \\
 \underline{x^3 - x^2} \phantom{+ 7x - 5} \\
 -2x^2 + 7x - 5 \\
 \underline{-2x^2 + 2x} \phantom{- 5} \\
 5x - 5 \\
 \underline{5x - 5} \\
 0 \qquad \qquad \qquad = M(x) \text{ remainder}
 \end{array}$$

and hence  $P(x) = (x - 1)(x^2 - 2x + 5)$ . Since  $x^2 - 2x + 5 = (x - 1)^2 + 4 > 0$  this quadratic polynomial does not have real roots, and so it does not factorise further.

## 5.2 Introducing partial fractions

Two rational functions can be added together to give another rational function. This involves bringing them to a common denominator, the simplest method being cross-multiplication.

**Example**

$$\frac{1}{x-1} + \frac{2}{x+2} = \frac{1(x+2)}{(x-1)(x+2)} + \frac{2(x-1)}{(x+2)(x-1)} = \frac{1(x+2) + 2(x-1)}{(x-1)(x+2)} = \frac{3x}{x^2 + x - 2}$$

Note that the denominator of the sum factorises as the product of the original denominators. It is sometimes necessary (e.g., for integration) to carry out the reverse procedure, i.e., to split a rational function into the sum of two (or more) 'simpler' ones. To begin with, we note that factorising the denominator may give us a clue as to how to proceed.

**Example**

Consider the rational function

$$\frac{x + 7}{x^2 - x - 6}.$$

By factorising the denominator we see that  $x^2 - x - 6 = (x - 3)(x + 2)$ . This factorisation suggests that it may be possible to write our rational function as the sum of a proper rational function with denominator  $x - 3$  and a proper rational function with denominator  $x + 2$ . That is,

$$\frac{x + 7}{x^2 - x - 6} = \frac{A}{x + 2} + \frac{B}{x - 3},$$

where  $A$  and  $B$  are constants (or in other words, polynomials of degree 1) to be determined. We can check (by cross-multiplying) that setting  $A = -1$  and  $B = 2$  gives a solution to the above equation. Indeed,

$$\frac{-1}{x + 2} + \frac{2}{x - 3} = \frac{-(x - 3) + 2(x + 2)}{(x + 2)(x - 3)} = \frac{-x + 3 + 2x + 4}{(x + 2)(x - 3)} = \frac{x + 7}{(x + 2)(x - 3)}.$$

So the desired decomposition is:

$$\frac{x + 7}{x^2 - x - 6} = \frac{-1}{x + 2} + \frac{2}{x - 3}.$$

Notice that the right hand side can be integrated!

In order to decompose a proper rational function into a sum of the ‘simplest possible’ proper rational functions, we begin by factorising the denominator as much as possible.

**Definition**

An **irreducible** polynomial is a polynomial  $P(x)$  that does not have any non-trivial factors of degree  $d$  where  $1 \leq d < \deg(P)$ .

Essentially, an irreducible polynomial is one which cannot be factorised further. Obviously any linear (degree 1) polynomial  $ax + b$  is irreducible. A quadratic (degree 2) polynomial  $ax^2 + bx + c$  is irreducible if it has no linear (degree 1) factors, or in other words, if the equation  $ax^2 + bx + c = 0$  has no real solutions.

**Example**

The quadratic polynomial  $x^2 + 1$  is irreducible.

Using the quadratic formula, we see that there are no real solutions to the equation  $x^2 + 1 = 0$ :

$$\frac{-0 \pm \sqrt{0^2 - 4}}{2} = \pm i$$

(Here  $i$  denotes the square root of  $-1$ , which is not a real number.)

**Theorem 28**

Every polynomial can be written as a product of linear factors ( $ax + b$ , with  $a \neq 0$ ) and irreducible quadratic factors ( $ax^2 + bx + c$ , with  $a \neq 0$  and  $b^2 - 4ac < 0$ ).

**Example**

Note that irreducible factors such as linear factors or irreducible quadratics may repeat in the factorisation of a polynomial:

$$P(x) = (x - 1)^2(x + 5)(x^2 - x + 9)^3.$$

**Definition**

Expressing a rational function  $\frac{N(x)}{D(x)}$  as the sum of rational functions

$$\frac{N(x)}{D(x)} = \frac{N_1(x)}{D_1(x)} + \dots + \frac{N_k(x)}{D_k(x)}$$

whose denominators  $D_1(x), \dots, D_k(x)$  are irreducible polynomials (or their powers) is called decomposition into **partial fractions**.

The “**partial fractions**” are the terms in this sum.

## 5.3 The method of partial fractions

To split a given rational function into partial fractions:

**Step 1:** Use long division to obtain a proper rational function.

**Step 2:** Factorize the denominator as much as possible.

**Step 3:** Identify the type of partial fractions involved.

**Step 4:** Multiply out.

**Step 5:** Determine the constants.

Steps 1 and 2 have already been discussed in the previous sections. To identify the type of partial fractions involved, we look at the factorisation of the denominator.

Denominator contains	Partial fraction decomposition contains
Exactly one irreducible factor $ax + b$	$\frac{A}{ax + b}$ , where $A$ is a constant to be determined.
Exactly $n$ irreducible factors $ax + b$	$\frac{A_1}{ax + b} + \frac{A_2}{(ax + b)^2} + \cdots + \frac{A_n}{(ax + b)^n}$ , where $A_1, A_2, \dots, A_n$ are constants to be determined.
Exactly one irreducible factor $ax^2 + bx + c$	$\frac{Ax + B}{ax^2 + bx + c}$ , where $A$ and $B$ are constants to be determined.
Exactly $n$ irreducible factors $ax^2 + bx + c$	$\frac{A_1x + B_1}{ax^2 + bx + c} + \frac{A_2x + B_2}{(ax^2 + bx + c)^2} + \cdots + \frac{A_nx + B_n}{(ax^2 + bx + c)^n}$ , where $A_1, A_2, \dots, A_n$ and $B_1, B_2, \dots, B_n$ are constants to be determined.

**Example**

Let  $f(x)$  be any polynomial such that each of the given rational functions is proper. Then we have the following expansions into partial fractions:

$$\begin{aligned}\frac{f(x)}{(x+1)(x-2)} &= \frac{A}{x+1} + \frac{B}{x-2}, \\ \frac{f(x)}{(x+1)^3(x-2)} &= \frac{A}{x+1} + \frac{B}{(x+1)^2} + \frac{C}{(x+1)^3} + \frac{D}{x-2}, \\ \frac{f(x)}{(x^2+1)(x-2)} &= \frac{Ax+B}{x^2+1} + \frac{C}{x-2}, \\ \frac{f(x)}{(x^2+1)^2(x-2)} &= \frac{Ax+B}{x^2+1} + \frac{Cx+D}{(x^2+1)^2} + \frac{E}{x-2}.\end{aligned}$$

**Example**

Split the rational function  $\frac{5x-1}{x^2-x-2}$  into partial fractions.

**Step 1:** Note that we already have a proper rational function, so we do not need to perform long division.

**Step 2:** Factorising the denominator gives

$$\frac{5x-1}{x^2-x-2} = \frac{5x-1}{(x+1)(x-2)},$$

**Step 3:** Since the denominator has two distinct linear factors, we see that

$$\frac{5x-1}{x^2-x-2} = \frac{5x-1}{(x+1)(x-2)} = \frac{A}{x+1} + \frac{B}{x-2},$$

**Step 4:** Multiplying both sides of this expression by  $x^2-x-2$  yields

$$5x-1 = A(x-2) + B(x+1).$$

**Step 5:** By setting  $x = -1$ , we see that  $-6 = -3A$  and hence  $A = 2$ .

By setting  $x = 2$ , we see that  $9 = 3B$  and hence  $B = 3$ .

**Conclusion:**

$$\frac{5x-1}{x^2-x-2} = \frac{2}{x+1} + \frac{3}{x-2}.$$

Notice that you can check your answer by combining the partial fractions you have found back into a single rational function:

$$\frac{2}{x+1} + \frac{3}{x-2} = \frac{2(x-2) + 3(x+1)}{(x+1)(x-2)} = \frac{5x-1}{(x+1)(x-2)}.$$



**Example**

Split the rational function  $\frac{1}{x^2(x-1)}$  into partial fractions.

**Steps 1 and 2:** Since we are given a proper rational function whose denominator has already been factorised as much as possible, there is nothing to do.

**Step 3:** Since the denominator factorises into three linear factors (one is repeated!) we have

$$\frac{1}{x^2(x-1)} = \frac{A}{x} + \frac{B}{x^2} + \frac{C}{x-1}.$$

**Step 4:** Multiplying both sides by  $x^2(x-1)$ , we obtain  $Ax(x-1) + B(x-1) + Cx^2 = 1$ .

**Step 5:** By setting  $x = 0$ , we see that  $-B = 1$  or in other words  $B = -1$ .

By setting  $x = 1$ , we see that  $C = 1$ .

Now by comparing the coefficients of  $x^2$ , we find that  $A + C = 0$ , so  $A = -1$ .

**Conclusion:**

$$\frac{1}{x^2(x-1)} = -\frac{1}{x} - \frac{1}{x^2} + \frac{1}{x-1}.$$

**Example**

Split the improper rational function

$$\frac{x^4 + 2x^3 - 2x^2 - x + 4}{x^3 + 4x^2 + 5x + 2}$$

into partial fractions.

**Step 1:** Using polynomial long division (see calculation in section 2.1) we find that

$$\frac{x^4 + 2x^3 - 2x^2 - x + 4}{x^3 + 4x^2 + 5x + 2} = (x - 2) + \frac{x^2 + 7x + 8}{x^3 + 4x^2 + 5x + 2}.$$

**Step 2:** To factorise the cubic denominator  $D(x) = x^3 + 4x^2 + 5x + 2$ , we begin by trying to spot a linear factor by checking the possible factors of the constant term. Since  $D(-1) = 0$ , we note that by the remainder theorem  $x + 1$  is a factor. Hence

$$\begin{aligned} x^3 + 4x^2 + 5x + 2 &= (x + 1)(x^2 + ax + b) \\ &= (x + 1)(x^2 + 3x + 2) \\ &= (x + 1)(x + 1)(x + 2) \\ &= (x + 1)^2(x + 2). \end{aligned}$$

**Step 3:** Using the factorisation of the denominator we find

$$\frac{x^2 + 7x + 8}{x^3 + 4x^2 + 5x + 2} = \frac{x^2 + 7x + 8}{(x + 1)^2(x + 2)} = \frac{A}{x + 1} + \frac{B}{(x + 1)^2} + \frac{C}{x + 2}.$$

**Step 4:** Multiplying both sides of this last equation by  $x^3 + 4x^2 + 5x + 2$  gives

$$\begin{aligned} x^2 + 7x + 8 &= \frac{A(x + 1)^2(x + 2)}{(x + 1)} + \frac{B(x + 1)^2(x + 2)}{(x + 1)^2} + \frac{C(x + 1)^2(x + 2)}{(x + 2)} \\ &= A(x + 1)(x + 2) + B(x + 2) + C(x + 1)^2. \end{aligned}$$

**Step 5:** Setting  $x = -1$  gives  $2 = B$ . Setting  $x = -2$  gives  $-2 = C$ . Comparing the coefficients of  $x^2$  gives  $1 = A + C$ , and hence  $A = 3$ .

Thus

$$\frac{x^2 + 7x + 8}{x^3 + 4x^2 + 5x + 2} = \frac{3}{x + 1} + \frac{2}{(x + 1)^2} - \frac{2}{x + 2}.$$

**Conclusion:**

$$\frac{x^4 + 2x^3 - 2x^2 - x + 4}{x^3 + 4x^2 + 5x + 2} = x - 2 + \frac{3}{x + 1} + \frac{2}{(x + 1)^2} - \frac{2}{x + 2}.$$

**Example**

Split the rational function  $\frac{1}{(x+1)^2(x^2+1)}$  into partial fractions.

**Steps 1 and 2:** Since we have been given a proper rational function whose denominator has already been factorised as much as possible, there is nothing to do.

**Step 3:** Using the factorisation of the denominator we obtain

$$\frac{1}{(x+1)^2(x^2+1)} = \frac{A}{x+1} + \frac{B}{(x+1)^2} + \frac{Cx+D}{x^2+1}.$$

**Step 4:** By multiplying both sides by  $(x+1)^2(x^2+1)$  we arrive at

$$\begin{aligned} 1 &= A(x+1)(x^2+1) + B(x^2+1) + (Cx+D)(x+1)^2 \\ &= A(x^3+x^2+x+1) + B(x^2+1) + C(x^3+2x^2+x) + D(x^2+2x+1). \end{aligned}$$

**Step 5:** Collecting terms with the same power of  $x$  gives

$$1 = (A+C)x^3 + (A+B+2C+D)x^2 + (A+C+2D)x + A+B+D.$$

By comparing the coefficients on each side of this equation we obtain a system of four linear equations for four variables  $A, B, C, D$ :

$$\begin{aligned} \text{coefficient of } x^3: & A+C & & = 0 \\ \text{coefficient of } x^2: & A+B+2C+D & & = 0 \\ \text{coefficient of } x^1: & A+C+2D & & = 0 \\ \text{coefficient of } x^0: & A+B+D & & = 1 \end{aligned}$$

By subtracting the first equations from the others, we can eliminate  $A$  from them:

$$\begin{aligned} A+C &= 0 \\ B+C+D &= 0 \\ 2D &= 0 \\ B-C+D &= 1 \end{aligned}$$

which is equivalent to the system

$$\begin{aligned} A+C &= 0 \\ B+C &= 0 \\ B-C &= 1 \\ D &= 0 \end{aligned}$$

By back substitution, we find  $A = \frac{1}{2}$ ,  $B = \frac{1}{2}$ ,  $C = -\frac{1}{2}$ , and  $D = 0$ .

**Conclusion:**

$$\frac{1}{(x+1)^2(x^2+1)} = \frac{1}{2(x+1)} + \frac{1}{2(x+1)^2} - \frac{x}{2(x^2+1)}.$$

**Example**

Split the rational function  $\frac{x^2 + 1}{x^3 + 7x}$  into partial fractions.

**Step 1:** Note that we already have a proper rational function, so we do not need to perform long division.

**Step 2:** Factorising the denominator gives

$$\frac{x^2 + 1}{x^3 + 7x} = \frac{x^2 + 1}{x(x^2 + 7)},$$

**Step 3:** Since the denominator has one linear factor and one irreducible quadratic factor, we see that

$$\frac{x^2 + 1}{x^3 + 7x} = \frac{x^2 + 1}{x(x^2 + 7)} = \frac{A}{x} + \frac{Bx + C}{x^2 + 7}.$$

**Step 4:** Multiplying both sides of this expression by the denominator  $x^3 + 7x$  yields

$$x^2 + 1 = A(x^2 + 7) + (Bx + C)x.$$

**Step 5:** Comparing the constant terms, we have  $1 = 7A$ , so  $A = \frac{1}{7}$ .

Comparing coefficients of  $x^2$  gives  $1 = A + B$ . Hence  $B = \frac{6}{7}$ .

Comparing coefficients of  $x$  gives  $0 = C$ , i.e.,  $C = 0$ .

**Conclusion:**

$$\frac{x^2 + 1}{x^3 + 7x} = \frac{1/7}{x} + \frac{6x/7}{x^2 + 7} = \frac{1}{7x} + \frac{6x}{7(x^2 + 7)}.$$

## 5.4 Integration by partial fractions

### Fractions with a quadratic denominator

Consider an integral of the following form:

$$\int \frac{dx}{ax^2 + bx + c}$$

assuming that  $a \neq 0$  (otherwise the denominator is not quadratic). To solve the integral, we transform the quadratic polynomial  $f(x) = ax^2 + bx + c$  as follows:

$$f(x) = a \left( x^2 + \frac{b}{a}x + \frac{c}{a} \right) = a \left[ \left( x + \frac{b}{2a} \right)^2 + \frac{c}{a} - \frac{b^2}{4a^2} \right] = a \left[ \left( x + \frac{b}{2a} \right)^2 - \frac{(b^2 - 4ac)}{4a^2} \right].$$

(This is called **completing the square**.) Now there are two cases:

1.  $b^2 - 4ac \geq 0$ , so there are real roots,  $f(x) = a(x - x_1)(x - x_2)$ ; or
2.  $b^2 - 4ac < 0$ , so the quadratic is irreducible.

In the first case, the fraction we started with decomposes into fractions with linear denominators, which we know how to integrate; the answer will contain logarithms. In the second case, by a substitution, the integral can be transformed into a table integral and the answer will involve  $\tan^{-1}$ . We consider some examples to illustrate this point.

**Example**

Solve

$$\int \frac{dx}{x^2 + x - 2}.$$

**Solution:** Here we see that

$$\begin{aligned} x^2 + x - 2 &= \left(x + \frac{1}{2}\right)^2 - \frac{1}{4} - 2 = \left(x + \frac{1}{2}\right)^2 - \frac{9}{4} = \left(x + \frac{1}{2}\right)^2 - \left(\frac{3}{2}\right)^2 \\ &= \left(\left(x + \frac{1}{2}\right) - \frac{3}{2}\right) \left(\left(x + \frac{1}{2}\right) + \frac{3}{2}\right) = (x - 1)(x + 2) \end{aligned}$$

(so completing the square gives the roots of the quadratic).

Using the method of partial fractions we now find that:

$$\frac{1}{x^2 + x - 2} = \frac{1}{3} \left( \frac{1}{x - 1} - \frac{1}{x + 2} \right),$$

and hence

$$\begin{aligned} \int \frac{dx}{x^2 + x - 2} &= \frac{1}{3} \left( \int \frac{dx}{x - 1} - \int \frac{dx}{x + 2} \right) \\ &= \frac{1}{3} \ln|x - 1| - \frac{1}{3} \ln|x + 2| + C = \frac{1}{3} \ln \left| \frac{x - 1}{x + 2} \right| + C. \end{aligned}$$

**Example**

Find

$$\int \frac{dx}{x^2 + 2x + 2}.$$

**Solution:** Completing the square:  $x^2 + 2x + 2 = (x + 1)^2 + 2 - 1 = (x + 1)^2 + 1$ . This time we find that the denominator is an irreducible quadratic. Now making the substitution  $u = x + 1$ , gives  $du = dx$  and so the integral becomes:

$$\int \frac{dx}{x^2 + 2x + 2} = \int \frac{dx}{(x + 1)^2 + 1} = \int \frac{du}{1 + u^2} = \tan^{-1}(u) + C = \tan^{-1}(x + 1) + C.$$

**Example**

Find

$$\int \frac{5 dx}{x^2 + 4x + 13}.$$

**Solution:** Completing the square,  $x^2 + 4x + 13 = (x + 2)^2 + 9 = (x + 2)^2 + 3^2$  allows us to rewrite the integral as follows:

$$\int \frac{5 dx}{x^2 + 4x + 13} = \int \frac{5 dx}{(x + 2)^2 + 9} = \frac{5}{9} \int \frac{dx}{\left(\frac{(x + 2)}{3}\right)^2 + 1}.$$

By making the substitution  $u = \frac{1}{3}(x + 2)$  the denominator can be reduced to  $(u^2 + 1)$ . We have  $du = \frac{1}{3}dx$  or  $dx = 3du$  and so

$$\begin{aligned} \frac{5}{9} \int \frac{dx}{\left(\frac{(x + 2)}{3}\right)^2 + 1} &= \frac{5}{9} \int \frac{\cdot 3 du}{u^2 + 1} \\ &= \frac{5}{3} \int \frac{du}{u^2 + 1} \\ &= \frac{5}{3} \tan^{-1}(u) + C \\ &= \frac{5}{3} \tan^{-1}\left(\frac{1}{3}(x + 2)\right) + C. \end{aligned}$$

**Example**

Find

$$\int \frac{2x + 9}{x^2 + 4x + 13} dx$$

**Solution:** In this case the numerator depends on  $x$ . Notice that:  $d(x^2 + 4x + 13) = 2x + 4$  and we may write

$$\int \frac{2x + 9}{x^2 + 4x + 13} dx = \int \frac{2x + 4}{x^2 + 4x + 13} dx + \int \frac{5}{x^2 + 4x + 13} dx$$

Now, the first integral gives rise to a  $\ln$  term, since the numerator is the derivative of the denominator, while the second integral is as in the previous example. Thus,

$$\int \frac{2x + 9}{x^2 + 4x + 13} dx = \ln(x^2 + 4x + 13) + \frac{5}{3} \tan^{-1}\left(\frac{x + 2}{3}\right) + C.$$

**In general when we integrate a proper rational function with an irreducible quadratic as the denominator, we should expect a  $\ln$  term plus a  $\tan^{-1}$  term.**

## Integration using partial fractions

The preceding examples hint that any rational function can be integrated by splitting it into partial fractions.

### Example

Find the integral

$$\int \frac{5x - 1}{x^2 - x - 2} dx.$$

**Solution:** Using partial fractions we find that

$$\frac{5x - 1}{x^2 - x - 2} = \frac{2}{x + 1} + \frac{3}{x - 2}.$$

Integrating then gives

$$\int \frac{5x - 1}{x^2 - x - 2} dx = \int \frac{2 dx}{x + 1} + \int \frac{3 dx}{x - 2} = 2 \ln |x + 1| + 3 \ln |x - 2| + C.$$

### Example

Find the integral

$$\int \frac{x^4 + 2x^3 - 2x^2 - x + 4}{x^3 + 4x^2 + 5x + 2} dx.$$

**Solution:** We met this rational function before (in chapter 2) and obtained that

$$\frac{x^4 + 2x^3 - 2x^2 - x + 4}{x^3 + 4x^2 + 5x + 2} = x - 2 + \frac{3}{x + 1} + \frac{2}{(x + 1)^2} - \frac{2}{x + 2}.$$

Altogether:

$$\begin{aligned} \int \frac{x^4 + 2x^3 - 2x^2 - x + 4}{x^3 + 4x^2 + 5x + 2} dx &= \int \left( x - 2 + \frac{3}{x + 1} + \frac{2}{(x + 1)^2} - \frac{2}{x + 2} \right) dx \\ &= \int (x - 2) dx + 3 \int \frac{dx}{(x + 1)} + 2 \int (x + 1)^{-2} dx \\ &\quad - 2 \int \frac{dx}{(x + 2)} \\ &= \frac{1}{2}x^2 - 2x + 3 \ln |x + 1| - \frac{2}{(x + 1)} \\ &\quad - 2 \ln |x + 2| + C. \end{aligned}$$

Recall that when we expand a rational function into partial fractions we may also obtain powers of irreducible quadratics in the denominator. The general strategy in this case is to transform the function we want to integrate so to reduce the power of the denominator. Integration by parts can be useful here...

**Example**

Consider the integral

$$\int \frac{dx}{1+x^2}$$

(which as we know is  $\tan^{-1}(x) + C$ ) and perform integration by parts taking  $u = \frac{1}{x^2+1}$  and  $dv = dx$ . We obtain

$$\begin{aligned} \int \frac{1}{1+x^2} dx &= \frac{x}{1+x^2} - \int x d\left(\frac{1}{1+x^2}\right) \\ &= \frac{x}{1+x^2} + \int x \frac{2x dx}{(1+x^2)^2} \\ &= \frac{x}{1+x^2} + \int \frac{2x^2 dx}{(1+x^2)^2} \\ &= \frac{x}{1+x^2} + \int \frac{(2x^2 + 2 - 2) dx}{(1+x^2)^2} \\ &= \frac{x}{1+x^2} + 2 \int \frac{dx}{1+x^2} - 2 \int \frac{dx}{(1+x^2)^2}, \end{aligned}$$

which gives a relation between the known integral  $\int \frac{dx}{1+x^2}$  and the integral  $\int \frac{dx}{(1+x^2)^2}$  that we want to find:

$$2 \int \frac{dx}{(1+x^2)^2} = \frac{x}{1+x^2} + \int \frac{dx}{1+x^2}.$$

This gives the answer:

$$\int \frac{dx}{(1+x^2)^2} = \frac{1}{2} \frac{x}{1+x^2} + \frac{1}{2} \tan^{-1}x + C.$$